

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search


Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/



HN 4NQU .

KF 24222 (2)





The Silvary of Havarelfallige will liftlenwood's respects.

		1

OBSERVATIONS

ON

METALLIFEROUS DEPOSITS,

AND ON

SUBTERRANEAN TEMPERATURE;

FORMING THE EIGHTH VOLUME OF THE TRANSACTIONS OF THE ROYAL GEOLOGICAL SOCIETY OF CORNWALL, Eng. —

PART THE SECOND.

BY

WILLIAM JORY HENWOOD, F.R.S.; F.G.S.;

MEMBER OF THE GEOLOGICAL SOCIETY OF FRANCE;

PRESIDENT OF THE ROYAL INSTITUTION OF CORNWALL;

HONORARY MEMBER OF THE YORKSHIEE PHILOSOPHICAL SOCIETY;

CORRESPONDING MEMBER OF THE IMPERIAL AGRICULTURAL AND NATURAL

HISTORY SOCIETY—LYONS, AND OF THE LYCRUM OF

NATURAL HISTORY—NEW YORK;

SOMETIME HER MAJESTY'S ASSAY-MASTER OF TIN IN THE DUCKY OF CORNWALL.

e PENZANCE:

PRINTED AND PUBLISHED BY WILLIAM CORNISH,
GREEN MARKET.

1871.

KF 24222 (2)

1871. Ock. 21,

Gift of
the Author,

Vm. G. Henwood,
of Tenzance,

Cornwall, Eng.

CONTENTS.

Observations on Subterranean Temperature. By WILLIAM JORY HENWOOD, F.R.S.; F.G.S.; Member of the Geological Society of France; President of the Royal Institution of Cornwall; Honorary Member of the Yorkshire Philosophical Society; Corresponding Member of the Imperial Agricultural and Natural History Society—Lyons, and of the Lyceum of Natural History—New York; sometime Her Majesty's Assay-Master of Tin in the Duchy of Cornwall; Member of the Society.

CHILI.

CHAÑARCILLO.

Colorada. Temperatures at the surface and in the rocks at various depths. 724—725.

BRAZIL.

MINAS GERAES.

- Morro Velho. Temperatures,—of air at the surface,—and of water issuing from the rock and the metalliferous deposits at various depths. 725—727.
- Gongo Soco. Temperatures,—of air and of wells at the surface,—and of streams issuing from the rock and the metalliferous deposit at various depths. 727—729.
- Agoa Quente. Temperatures,—of air and water at the surface,—
 of streams issuing from the rock and the metalliferous deposit,
 —and of water accumulated in the mine. 729—731.
- Fraga. Temperature of water issuing from the metalliferous dedeposit. 732.

NORTH AMERICA.

VIRGINIA.

Garnett and Moseley. Temperature of water pumped to the surface.

MICHIGAN.

Toltec and Douglass Houghton (Henwood). Formation of ice in the works during winter. 783—734.

THE CHANNEL ISLANDS,

SARK.

Windmill Hill. Temperature of water in a well at the surface, 735.

Port ès Sées.

"streams issuing from the rock and lode at different depths. 735.

Le Pot.

"streams issuing from the rock and lode at different depths. 736.

Sark's-Hope.

"streams issuing from the rock and lode at different depths. 736.

HERM.

Mins. Temperature,—of water in a well at the surface,—and of streams issuing from the rock and the *lode* at various depths. 737.

IRELAND.

WICKLOW.

Connorree. Temperatures,—of water in a well at the surface and of streams issuing from the rock and the metalliferous deposit, at different depths. 738.

Cronebane. Temperature of water flowing from the metalliferous deposit. 738.

WATERFORD.

Knockmahon. Temperatures,—of water in a well at the surface,—and of streams issuing from the rock and the lode, at different depths. 740.

CORK.

Bearhaven. Temperature of water issuing from the rock and lode, at different depths. 742,

KERRY.

Ardtully. Temperature of water issuing from the rock and lode, at different depths. 744.

ENGLAND.

CORNWALL.

	Cornwa	LL.
CARADOM Te	mperature of	water in a well at the surface. 745.
Gonamena.	>>	water flowing from the rock and lode. 746.
South Caradon.	"	water flowing from the rock the lodes, and the cross- veins, at different depths. 746.
Phonix.	27	water flowing from the rock and lode. 746.
Mark Valley.	29	water flowing from the lode. 746.
Мвинанот.	Temperatur	re of water in neighbouring wells at the surface. 748.
South Wh. Trelawny	'· ,,	water issuing from the branches at different depths. 748.
Wheal Mary Ann.	"	water issuing from the <i>lode</i> at different depths. 748.
Wheal Trelawny.	"	water issuing from the lode at different depths. 748.
Landrath & Saint Pinnock.	Temperatur	e of water in neighbouring wells at the surface. 749.
Herod's-foot.	"	water issuing from the lods at various depths. 749.

HRO PSHIRE.

Eardiston. Temperature of the water issuing from the rock and the metalliferous deposit, at various depths. 751.

On the mean depths, — temperatures, — and rates at which the temperatures increase with the depths,—of mines in each region. 751—752.

- On the mean depths, temperatures, and rates at which the temperatures increase with the depths, of mines in different rocks. 752—756.
- On the mean depths, temperatures, and rates at which the temperatures increase with the depths,—of mines yielding different metals and ores. 756—758.
- On the mean depths, temperatures, and rates at which the temperatures increase with the depths,—of mines affording similar metals and ores in different rocks, 757—758.
- On the mean depths, temperatures, and rates at which the temperatures increase with the depths, of mines wrought at different altitudes. 758—759.
- Observations at different (mean) depths, in elevated tropical regions, and at smaller elevations in temperate climates. 760—761.
- On the influence of deepening shafts and opening (levels) galleries on the normal temperatures of mines. 763—765.
- On occasional interruptions in the progressive increase of temperature with depth. 765—766.
- On the change of temperature which takes place—at the same, and at different, times,—on the surface, and at depths of three, six, and nine feet in the Canga, at Agoa Quente—Brazil. 767—780. Explanations of the Plates and Woodcuts. 781—782. Index. 783—916.

TABLES.

Relations of rocks to metalliferous deposits, and to cross-veins.

—summaries of mining and of financial operations,—proportions of copper precipitated to iron used in precipitation.—Temperatures at the surface and underground.

NORTH-WESTERN INDIA.

KUMAON AND GURHWAL.

- Relations between the rocks and the metalliferous deposit of Agur.
- II. Relations between the rocks and the metalliferous deposit of Kotelar and Khetsaree.

CHILI.

CHANARCILLO.

- III. Relations between the rocks, the metalliferous deposits, and the cross-veins, at Colorada and Desempeño.
- IV. Description of ore and (approximate) quantities of silver obtained from rich (bunches) parts of the Colorada lode.
- V. Forces employed and amounts paid for Salaries, Wages, Food, and Materials (1856) at Colorada.

BRAZIL.

MINAS GERAES.

- VI. Relations between the rocks and the metalliferous deposits at Morro Velho.
- VI.a Proportions of gold contained in, and extracted from, the vein-stone; force employed to extract it; and cost of its extraction (1855—1861), at Morro Velho.
- VII. Synopsis of Mining and Financial operations (1835—1861) at Morro Velho.
- VIII. Relations between the rocks and the metalliferous deposits at Gongo Soco.
 - IX. Synopsis of Mining and Financial operations (1826—1856) at Gongo Soco.
 - X. Synopsis of observations on the Gold-deposits of the Province.

NORTH AMERICA.

MICHIGAR

- XI. Relation between the rocks and the metalliferous deposits of the North American mine.
- XII. Relation between the rocks and the metalliferous deposits of the Cliff mine.
- XIII. Relation between the rocks and the metalliferons deposits of the South Cliff mine.
- XIV. Synopsis of the produce, working-costs, and profits of rich mines in various countries.
- XV. Relations between the rocks and metalliferous deposit of the Douglas Houghton (Henwood) mines.

FRANCE.

DAUPHINY.

XVI. Relations between the rocks the metalliferous deposits, and the cross-vein at Chalanches.

THE CHANNEL ISLANDS

SARK.

XVII. Relations between the rocks and the metalliferous deposit of Sark's-Hope.

TRELAND.

WICKLOW.

- XVIII. Relations between the rocks and the metalliferous deposit of Connorree.
- XVIII.s Quantities of copper-ore calcined,—iron used,—precipitate obtained,
 —and copper extracted at the *Mona* mine—Anglesea.

WATERFORD.

XIX. Relations between the rocks, the lodes, and the cross-veins of Knockmahon.

CORK.

XX. Relations between the rocks, the lodes, and the cross-veins at Bearhaven.

KERRY.

XXI. Relations between the rocks, the metalliferous deposits, and the cross-vein at Ardtully.

GREAT BRITAIN.

WALES.

- XXII. Proportions of gold extracted from the vein-stones of Cloque.
 - Gold extracted from detrital deposits and vein-stones in different Countries, means adopted for its extraction, and cost of extracting it.

CORNWALL.

- XXIII. Relations between the rocks, the lodes, and the cross-veins at West Caradon.
- XXIV. Relations between the rocks, the lodes, and the cross-veins at South Caradon.
- XXV. Relations between the rocks and the lode at the Phanix mines.
- XXVI. Relations between the rocks, the lodes, and the cross-vein at Marke Valley.
- XXVII. Relations between the rocks, the lode, and the cross-vein at Wheal Trelawny.

- XXVIII. Relations between the rocks, the lode, and the cross-vein at Wheal Mary Ann.
 - XXIX. Relations between the rocks, the lode, and the cross-vein at Herod's-foot.

XXX.	Temperature	(surface) of	f Gongo Soco—Brazil.
XXXI.	"	" of	Agoa Quente— " .
	Subterranean	temperatur	e in different Countries.
XXXIII.	,,	"	in different rocks.
XXXIV.	"	"	in mines yielding different metals and ores.
XXXV.	"	? ?	in mines affording similar metals and ores in different rocks.
XXXVI.	27	"	in mines of different altitudes
XXXVII.			face, and at depths of three, six, langa at Agoa Quente—Brazil.

CORRECTIONS.

```
read 19° 50'
,, 60°-3
,, Ansermo
,, preparation
,, 168
                             for 18° 50°
                                                      TABLES.
                                     33 column 11 ......
36 ,, 25 ......
41 ,, 11 .....
  VII.
  "
                     5 ...
88 ...
89 ...
134 ...
13 ...
9 ...
48 ...
102 Note b
 viiı.
                                                                                       " 1830 January 21st
  ίχ.
Χ.
                                                                                           403
Tellurium and Silver
                                                                                       , Tellurium and Silver
, 0.0000440
, is alloyed with both
Silver and Tellurium
, 0.2640
,, 0-7640
                                                        " Smith ......
                                                                                       " Smyth.
                       19
23
3
99
35
2
                                                       , N. ....
, 18° 58′ 30′ ....
, 61′ ....
                                                                                       " W.
" 19° 58′ 30″
" 19° 58′ 80″
" 61′5
XXXI
                       14 column 19 ......
```

CORRECTIONS IN OTHER VOLUMES.

Vol. III., page 208, line 1, for exudation read emission ,, V. ,, 19, ,, 8, ,, titanium ,, uranium ,, VI. ,, 995, ,, 11, ,, pailadium ,, tellurium

Observations on Subterranean Temperature.

By WILLIAM JORY HENWOOD, F.R.S.; F.G.S.;

MEMBER OF THE GEOLOGICAL SOCIETY OF FRANCE;

PRESIDENT OF THE ROYAL INSTITUTION OF CORNWALL;

SOMETIME HER MAJESTY'S ASSAY-MASTER OF TIN IN THE DUCHY OF CORNWALL;

MEMBER OF THE SOCIETY.

The first series of results* recorded in the following columns, was obtained in an absolutely dry†—though a deep—mine, by placing the thermometers in holes which had sometime before been purposely bored in the several limestones; * all others were determined in streams of water immediately as they issued ‡ from the various rocks and veins.

[•] Postea, p. 725.

^{† &}quot;It is in the solid rock that the best observations, and those most suited to the purpose of philosophical reasoning, are to be obtained."

PHILLIPS, Reports of the British Association, v. (1836), p. 292.

[&]quot;I am disposed to attach most importance to observations on springs of water, not coming from the roofs of galleries, or evidently proceeding from higher parts of the mines."—Fox, Cornwall Geol. Trans., III. p. 320.

[&]quot;After most careful consideration of the subject, and consultation with others who have also been engaged in this enquiry, it has been thought best to confine the observations, as much as possible to the temperature of the streams of water immediately issuing from the unbroken portions of the rocks and veins. The reasons for this preference are;—that the temperature of the air in mines is affected, not only by the presence of the workmen, the combustion of candles, and the explosion of gunpowder, but also by the warm or cold air which is brought to the same spot by the varying directions of the currents underground, which are more or less influenced by the changes of wind at the surface; that the rocks, forming the sides of the shafts and levels, must, to a certain extent, partake of the temperature of the air circulating through them, and, of course, be affected by its changes;—and that the water flowing through, or standing in pools in the levels, is exposed to the same modifying causes, and probably also, warmed by the workmen who frequently stand in it."

CHILL.

PROVINCE OF ATACAMA.—DEPARTMENT OF COPIAPÓ

DISTRICT OF CHAMARCILLO. Long. 70° 80' W., Lat. 27° 16' 8.

Mine of Colorada.* · Elevation of the surface about 3,650 feet above the Pacific; † 1,750 feet above the plain. ‡

The first,—third,—and fifth strata are of limestone; §

" second,——and fourth—— " " hornblendic rocks.§

The lodes yield silver and many of its ores in great abundance; beside iron-pyrites and blende in smaller proportions.

The	extremes	of temperature	during th	e year have been considered 52°68°; T
,,	mean	"	,,	has been estimated at
				about 64°.**
The	range	99	from the	9th to the 15th of June,
				1857, was 89°·5-66°·5; ††
99	mean	**	**	about 48°.5.††

[•] Ante, p. 90; Table III.; Pl. I., II.

Highest

DONEYKO, Annales des Mines, 4me Série, IX. p. 433.

^{**} Keith Johnston, Atlas of Physical Phonomena, Pl. XVIII.

†† Temperature in the shade at t	he surf	ace of	the Colo	rada m	ine:-	
Date.	7 A.M.	9 A.M.	Noor.	8 A.M.	6 P.M.	9 P.M.
1857, June 9th	58°-5	62°·8	•	۰	۰	۰
,, ,, 10th		61.	66.5	61.6	56.5	50·5 48·8
" " 11th		46.8	53.8		••	48.8
", ", 12th		45.	l	· ·		l
" " " 13th	39.5	43.8			44· 42·	42· 41·8
,, ,, 14th	39.8	42.	48.	46	42.	41.8
,, ,, 15th	42.8	43.8				

58.5

39.5

62.8

66.5

61.6 | 56.5 |

50.5

[†] Domeyko, Annales des Mines, 4me Série, 1x. p. 433. Henwood, Reports of the Royal Institution of Cornwall, xxxix. p. 15; Edin. New Phil. Journal, VII. N.S. p. 147.

[†] Henwood, Reports of the Royal Institution of Cornwall, XXXIX. p. 15; Edin. New Phil. Journal, VII. N.S. p. 147.

[§] Domeyko, Annales des Mines, 4me Série, 1x. pp. 435—40. Ante, pp. 69, 79.

B Domeyko, Annales des Mines, 4me Série, 1x. pp. 441—53. Ante pp. 86—118.

T" Le climat de cette montagne est très-doux et tempéré; mais il n'y pleut que tous 8 à 9 ans. * * Il est rare que le thermomètre y monte à plus de 20° C. [68° F.] à l'ombre, et qu'il descende au dessous de + 9° [52° F.]."

The stations at which observations were made, and the temperatures observed underground, were the following:—

		Locality.	Depths. fms.	Temperature of rock, in hole 2 feet deep.	Temperature of air circulating through the mine at the same spot.
lst	Limestone;	between Waring's ; and the Colorada lode lode on the E. ; on the W	4 6·	64'8	66·
2nd	Limestone ;	, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,	127	67.5	66.75
>9	,, ;	at the bottom of the shaft	150	67∙ •	66.
3rd	Limestone;	E. side (wall) of the an unfrequented part Colorada lode	227	72.	76.
"	99	, : a frequented part of the mine	,,	74.5	76.5

BRAZIL.

Province of Minas Geraes,—District of Rio das Velhas,— Parise of Congoneas de Sarara'; Long. 43° 50' W., Lat. 19° 58' 20" S.

Mine of Morro Velho.* Elevation of the surface about 3,250 feet above the sea.

Wrought in clay-slate.

[&]quot;If, in the absence of observations at midnight, and at 3 a.m., we assume the mean temperatures at those hours to have been 45°, which at this season cannot be wide of the truth; we have an average of about 48°.5 during the twenty-four hours.

[&]quot;On the 11th of June the thermometer stood at 53° .8 in the shade at noon.

",",66°.8 ", sunshine ", ."
"On that day and on the 15th of June much rain fell."

Henwood, Reports of the Royal Institution of Cornwall, XXXIX. p. 15; Edin. New Phil. Journal, VII. N.S. p. 148.

^{*&}quot; This observation, made at the bottom of the shaft, where the draught was very great, ought, perhaps, to be excluded from the general average."—Ibid.

[†] Von Bechwege, Pluto Brasiliensis, t xvi. Caldeleugh, Travels in Brasil,

The metalliferous deposit affords enormous quantities of auriferous iron-pyrites, beside much smaller proportions of arsenical-pyrites and copper-pyrites; * in vein-stones of quartz and quartzose slate.†

From July, 1868, to June, 1869, the temperature at the surface ranged from 40° to 86°, and averaged 66°84.‡

The temperatures in the same, and in different, parts of the mine § at various times, are shown in the following columns:—

				Local	ities.			
		Bahù. Cachosira. Periods.						
Localities.	Depth. Fathoms.	1848. • December.	1863.¶ July.	1864.¶ January.	1843.• December.	1868.¶ July.	1864.¶ January.	
		Temperatures.						
Water issuing pumps	12.	::	64.	68 [°] .75	°	65 [°] 12	69-25	

II. pp. 271—4. von Spix und von Martius, Reise en Brasilien, II. pp. 417—18. Saint Hilaire, Voyage dans le district des Diamans, I. p. 169. Gardner, Travels in Brasil, p. 496. Clauseen, Bulletins de l'Académie Royale de Bruxelles, VIII. Ire partie, p. 323. Whitney, Metallic Wealth of the United States, pp. 111—12. Burton, Exploration of the Highlands of the Brazil, 1. p. 251. Phillips (J. A.), Mining and Metallurgy of Gold and Silver, pp. 80—3, 210—20. Henwood, Cornwall Geol. Trans., VI. p. 143; London, Edinburgh, and Dublin Phil Mag., 3rd Series, xxv. p. 343; Ante, pp. 184—209.

[•] Henwood, Cornwall Geol. Trans., vi. p. 144; London, Edinburgh, and Dublin Phil. Mag., 3rd Series, xxv. p. 344; Ante, pp. 194—8.

⁺ Ibid.

[†] John Hookin, Esq., Chairman of the Saint John d'el Rey Company, M.S. § "Temperature at 7 mètres [8·8 fms.] below the surface 20°-65 C. [69°-17 F.]

,, 271·6 ,, [148·5 ,,] ,, 27°-22 ,, ." [81° ,,]

BURTON, Exploration of the Highlands of the Brazil, I. p. 251, Note.

[§] Henwood, Cornwall Geol. Trans., vi. p. 144, Pl. I.; London, Edinburgh, and Dublin Phil. Mag., 3rd Series, xxv. pp. 384; Ante, pp. 188—90, Pl. III.; Proceedings of the Royal Geological Society of Cornwall, 24th Oct., 1865.

Ter these observations the writer is indebted to the cordial co-operation of

					Local	ities.		
				Bahù. Cachoeira. Periods.				
	Localities.	Depths. Fathoms.	1843. December.	18 68. July.	1864. January.	1848. December.	18 68. July.	1864. January.
					Tempe	rature	 I.	
Waterissuin from th	g) e) rock upper side (hang- ing-wall) S. of metal- liferous deposit		•:	6 <u>4</u> .5	66°		•	•
**	metalliferous deposit .	45.	68-			69.+		
**	rock lower side (foot- toall) N. of metal- liferous deposit	58·6			67·			
"	,,	77.			••		••	60-16
"	rock upper side (hang- ing-wall) S. of metal- liferous deposit		 					72.*
21	,,	150.						70-8
**	metalliferous deposit .	155.					72.*	72.
Water collections Engine-si	ested at the bottom of the	160·		65.	69-5	<i></i>	69-05	71.5

Parish of Carthe'; Long. 43° 30' W., Lat. 19° 58' 30" S.

Mine of Gongo Soco.† Elevation of the surface about 3,360 feet above the sea.

J. N. Gordon, Esq., Resident Superintendent of the mine, and to the kindness of John Hockin, Esq., Chairman of the Saint John d'el Rey Mining Company.

e" At the celebrated gold-mine of Morro Velho, in Brazil, situate at a height of 3250 feet above the sea, and opened in clay-slate; the water issuing from the rock at 45 fathoms depth, observed in 1843, had a temperature of 69°; that at the bottom of the mine in 1863 and 1864, at 145 and 155 fathoms deep 72°. These temperatures were quite independent of the warm rains a little before and after Christmas, which make themselves felt all the way down the engine-shafts."—Smyth (Presidential Address to the Geological Society of London in 1868), Quarterly Journal of the Geological Society, xxiv, p. lxxxvi., Note.

[†] Von Bechwege, Pluto Brasiliensis, pp. 311—44. Gardner, Travels in Brasil, p. 491. Claussen, Bulletins, de l'Académis Royale de Bruxelles, VIII. 1re Partie, p. 327. Whitney, Metallic Wealth of the United States, p. 111. Phillips (J. A.), Mining and Metallurgy of Gold and Silver, p. 84. Ante, pp. 248—96, Pl. IV.

Wrought, for the most part, in (Jacotinga*) ironglance mixed with black, brown, and yellowish earthy iron-ore, as well as with friable black manganese and both buff-coloured and pearl-white talc; in some places, however, the iron-glance is replaced by quartz.

From (9,459) observations, made at intervals of three hours, it was ascertained, that during 1845, 1846, and 1847 the temperature, at the surface, ranged from 40°8 to 91°7, and averaged about 66°5.†

Streams issuing from the ground have, at different times, been found of the undermentioned temperatures.

			ratures.
Localities.	Depthr	1843. October.	1845. July.
The water issuing from a low hill S. of the valle passes through an ancient, long-abandoned, drif some 20 fathoms above the horizon of the adit fathom) level; in the mine, and supplies a (which is protected from both direct and refle	ft,— (48 well	•	•
sunshine) varied at different times			67.3-68
Water issuing from the pumps at the (48-fm.) adit le	mel‡ 34·	67 ·	
p) p) 29 29 29	‡ ··		67•
" " auriferous (Jacotinga) forma	tion		67·1
"; a large stream out of earthy brown iron and quarts, which represents the surifer (Jacotinga) formation, E.	ore rous 41	68-	66-6

[•] von Eschwege, Pluto Brasiliensis, p. 311. Hocheder, Report of the Imperial Brazilian Mining Association, xv. p. 54. Henwood Cornwall Geol. Trans., vi. pp. 227,—94; Ante, pp. 214,—19,—21,—3,—8,—42,—4,—6,—54,—6,—8,—63,—5.

[†] Henwood, London, Edinburgh, and Dublin Phil. Mag., 3rd Series, xxvIII. pp. 364-8; xxx. pp. 361-4; xxxIII. pp. 422-5; Table XXX.

[†] This drift is 48 fathoms deep at Lyon's shaft, but is only 34 ,, , Veseys (Engine) shaft. Ante, Table VIII., Note d; Pl. IV., Fig. 1, 2.

	Localities.				Temperatures.			
		1	Depths. fms.	1843. October.	1845. July.			
Wate	er, a small	l stream	from	(Itabirite)	and quarts		0	۰
				auriferous formation	(Jacotinga)	41,	67-5	67
>9	, ,,		out of	,,	••	48·	67·7—68	
,,	, a large	stream	"	"	••	,,		67•
"				of auriferou	s (Jacotinga)	,,	67.7	
23	,	,,		,,	••••	62		67.8
"	, a large	quartz	(Itabi		n-glance and d auriferous		••	66-8
••		rate stre irite), N		m iron-glan	ce and quarts	,,		67·3

DISTRICT OF VILLA RICA .-

PARISH OF CATTAS ALTAS. Long. 43° 10' W., Lat. 18° 50' S.

Mine of Agoa Quente. Elevation of the surface about 3.400 feet above the sea.

Wrought in Jacotinga†, composed of quartz in unequal—but sometimes in considerable—proportions, minute crystals of oxydulated and titaniferous iron, scales of micaceous iron-ore, flakes of talc, and small nests of felspar-clay, imbedded in earthy brown iron-ore tinged, at intervals, with earthy black manganese.

During 1848—9 the temperature at the surface ranged from 42. to 84°8 (Table XXXI.) and averaged about 60°.3.‡

[•] Von Eschwege, Pluto Brasiliensis, p. 299. De Monlevade, Annales des Mines, IV. p. 136. Caldeleugh, Travels in South America, II. p. 283.

[†] Von Eschwege, Pluto Brasiliensis, t IV. p. 299. Ante, pp. 224-36.

I Notwithstanding the monthly means at Agoa Quente, between October, 1848,

The undermentioned observations were made at times when very different quantities of rain-water *— absorbed at the surface—found their way into the mine; and when the works were opened at different depths.

	<u> </u>		Localisies			,	1844.	1847.	1849.		
		Localitie	6,			Depth.	1	Nov.	Jan.	▲pr.	June
Wate	er in a brook	at the su	rface	••••			:.	68·	7 6 ·	•	۰
"	out of the	back of the	(level) drift at	the end	4.5			73.		
,,		"	>>	2 feet i	rom "	.,			76-		
,,		,,	,,	3 "	,,	. ,			74.		
"	out of anci-	ent—& lon	g-abs	ndoned-	-works	6.		70-2		l	
,,		am out of (wall) of a formation					92.				
99	out of ancie	nt-& long	-aban	doned-w	or ks (a	8.	72.				
"		eams jett inga) form n both (wo	ation	, and $(I$	tabirite)	91.5				
"	, small stre deposi			rous (Ja		12.			80.7		
,,	, large stre	am, botto	m of a	ın Engin	e-shaft	15.		88.			
27	, ,,	"	8	second	"	. "		96.5			
,,	, ,,	,,		"	,,	. 18-			91.		
,,	, ,,	form	ation	rous (Ja at the b e-shaft .					85.		

and July, 1849 (Table XXXI.), differed somewhat from those at Gongo Soco during 1845—7 (Table XXX.); the general average, for corresponding periods, coincided within 0°.3.

[•] The rain which fell at Agoa Quente during the same period was

1848, October ... 3·28 inches. 1849, March ... 16·86 inches.

,, November .. 12·08 ,, . ,, April ... 7·98 ,, .

,, December .. 24·80 ,, . ,, May ... 8·14 ,, .

1849, January ... 15·10 ,, . ,, June ... 0·88 ,, .

,, February ... 19·86 ,, . ,, July ... - ,, .

Total ... 108·98 inches,

						Depth. fms.	1844.	1847.		1849.	
	Local	Localities.							Jan.	Apr.	June
Wate	er, large stream out o	of aurifer mation,				26.	•	•	•	90·5	۰
,,	, small stream out of for	of aurifer mation .			nga)	28.			••	••	77.
**	7 99		ithin o			,,				••	78.
,,	· · · · · · · · · · · · · · · · · · ·	of (<i>Itabir</i> v <i>all)</i> of a vga) form	arifero	us (Jaco-	,,			••	••	89-
**	, large stream out of	of aurifer rmation .				,,				••	88.
"	, " of	that last	within menti			,,				•••	92.
,,		ling up o				29.				92-5	
"	which filled the n to within 4.6 fm the surface, du	nine s. of ring {	stopps pump spot			15.		82.3			
**	,,	"	at a se	con	l spot	"		83.8			
"	drawn by pumps to the surface	at one	shaft	(b)	from	9.		81.9			
"	,,	,, a sec	ond "	(c)	,,	,,		84.			
"	**	, one	shaft	(b)	"	10.	84.				
,,	29	,, a sec	oùd "	(c)	,,	,,	87.	1			
,,	drawn by pumps to the adit (14 fathoms deep)	at one	shaft	(c)	"	18-			91.		
**	21	" anotl	her "	(b)	,,	24.			83-5		
٠,	"	•••••	••••		,,	29.				91.6	;

[•] Fish throve in this water. Ante, p. 355.

The power of fishes to bear extremes of temperature is well known.

YARRELL, History of British Fishes, z. pp. 316-19. COUCH, Fishes of the British Islands, IV. p. 33.

PARISH OF INFICIONADA; adjoining CATTAS ALTAS.

Mine of Fraga[®] or Ouro Fino. Elevation of the surface about 3,300 feet above the sea.

Wrought in that part of the talcose-slate series which overlies the (*Jacotinga*) manganesic iron-glance formation.

As Fraga is so near Agoa Quente and Gongo Soco, it, probably, differs little in climate, from them; but, inasmuch as it is less enclosed than they are by mountains and woods, its mean temperature may, perhaps, be somewhat cooler than theirs; on this, however, reliable observations have never been recorded.

The undermentioned temperatures have been observed:—

A large stream as it issues from the auriferous talc-slate into a (adit-level) drift opened from the vale 69°.

drawn by pumps to the surface from a depth of 21 fms. .. 70°.5

THE UNITED STATES.

STATE OF VIRGINIA,—COUNTY OF BUCKINGHAM. Long. 78° 30' W., Lat. 37° 35' N.

The Garnett and Moseley† mines have been wrought in chloritic, micaceous, and talcose slates, on a broad conformable bed of quartzose, felspathic, calcareous, and slaty matter, mixed with considerable quantities of earthy brown iron-ore near the surface, and of iron-pyrites at greater depths, as well as with smaller proportions of gold.

^{*} Von Eschwege, Pluto Brasiliensis, t v. Ante, pp. 301,-23.

[†] Rogers, Geological Reconnoisance of Virginia, p. 63. Ansted, Scenery Science, and Art, pp. 288-90. Ante, pp. 379-84.

The line of 55° mean annual temperature passes within a short distance of this district, if not directly through it.*

A small stream pumped to the surface from a depth of 15.5 fathoms maintained, during September, 1852, a temperature of 56°.8†

STATE OF MICHIGAN, COUNTY OF ONTONAGON (Long. 89° 30' W., Lat. 46° 50' N.).

The rains which immediately precede the first snows freeze almost as soon as they soak into the ground whilst the floods of autumn, which had been already absorbed, are—under influence of cold air, descending from the surface and circulating through the mines,

```
At Lac la Belle
the temperature of the air at the surface ..... was.....
                  ,, in the upper level .. ,, ......
                                                          47.
                                       ,, ............
                  air at 23.3 fms. deep
                                       ,, ...............
                  water ,,
                                       water ,,
 At Copper Falls-
the temperature of the air at the surface
                  water at 3.3 fms. deep .. " ........
                  air ,, 20.
                                                          49.
                                   .. ,, ..........
                  water,, ,,
                                       ,, .............
```

^{*} Keith Johnston, Atlas of Physical Phenomena, Pl. XVIII.

[†] On the 4th of September, 1852, the temperature at the surface about 6 p.m. was 76°.2.

[†] The following temperatures were observed in the Keweenaw district from (two hundred to four hundred feet above Lake Superior) eight hundred to a thousand feet above the sea; in mines wrought, in the trap formation, on lodes composed of calcareous-spar, prehnite, quartz, epidote, chlorite, and trappean matter. In most parts of the district some or others of these ingredients are more or less mixed with native copper, and this is frequently encrusted with virgin-silver (BAYFIELD, Quarterly Jour. of the Geol. Soc. 1. p. 451. BAUERMAN, Ibid, XXII. pp. 448-63. Jackson, Geological and Mineralogical Reports, Passim. Foster & Whitney, Geological Report, pp. 58—186. Ante, pp. 411—63, Tables XII.—XIV.).

during winter * -- often frozen as they issue, at considerable depths, from the rocks and lodes through which they had percolated. Thus, the streams which had entered at Toltec mine,† at 16.6 and at 23.3 fathoms, and the Douglas Houghton (Henwood) † mines at 36 fathoms from the surface, during the autumn of 1855 and become frozen during the succeeding winter, were yet unthawed in the following July.

THE CHANNEL ISLANDS.

The metalliferous rocks of Sark consist, in great measure, of felspar and hornblende, associated with

	American mine —	
the temperature	e of the air at the surface was	59∙
,,	a spring of water 16. fms. deep ,,	43·
",	,, ,, ,,	45.
,,	" ······ 26· " " " ······	44.
At The Cliff	mine—	
the temperature	e at the surface was	46·
,,		44.
,,	16.6, , ,,	43·
,,	20. ,, ,, ,,	44 ·
,,	39.3 ,, ,, ,,	45·
At (Fort Wil	kins) Copper Harbour the temperature, between June, 1844	i, and
May, 1846, ran	ged from 16°.35 to 72°.03 and averaged 41°.46.	
	eological and Mineralogical Report, pp. 443,—58,—9,—62,- k Whitney, Geological Report, p. 43.	- <i>5</i> 61.
	ter the pumps are occasionally covered with some non-condu t,—during stoppages of the machinery for needful repair,	_

influence of cold air from the surface should cause the water in them to freeze.

Daniel, Mining Journal, XXXVI. p. 390. Ante, pp. 465,-78.

[†] Whitney, Metallic Wealth of the United States, pp. 290-1, Fig. 23. Ante, p. 463.

¹ Jackson, Geological and Mineralogical Report, pp. 702,-42-3. Whitney, Geological Report, pp. 142,-50. Whitney, Metallic Wealth of the United States, pp. 289-90, Fig. 27. Ante, pp. 465-79; Table XV. Fig. 28.

[§] Mac Culloch, Geol. Trans., 1. p. 16. Prince, Cornwall Geol. Trans., VI. p.

smaller proportions of several other substances. The lodes which traverse them contain great quantities of the same ingredients; mixed, largely, with quartz and calcareous-spar, and, less plentifully, with earthy brown iron-ore, iron-pyrites, and yellow copper-ore. At the S.S.W. extremity of the island, however, the Sark's-Hope lode afforded also argentiferous and antimoniated galena, the super-sulphuret, sulphate, sulphato-tricarbonate, and carbonate of lead, together with the chloride of silver, earthy black silver-ore, as well as vitreous, red, and native silver,* where it was wrought beneath the sea.

The temperature of Sark is probably much the same as that of Guernsey, which ranges from 24°.5 to 83°,† and averages 51°.6.‡

The undermentioned temperatures were observed at different depths in various parts of the island:—

Windmill Hill.	De	pth.	Temp.
Well of fresh water (1841, January 26th)	Surface		47.
Port ès Sées.	Depth below the surface. fms.	Relation to the sea- level. fms.	
Small stream of fresh water, out of the rock	42.	16· A	<i>55</i> ·
Large ,, ,, ,, lode	54.	4. ₺	58.7

p. 101. Ansted, Channel Islands, pp. 263-6. Ante, pp. 630-2, Table XVII., Fig. 30.

^{*} Prince, Cornwall Geol. Trans., vi. p. 102. Ante, p. 535, Table XVII.

⁺ Ansted, Channel Islands, p. 140.

[‡] Ibid, p. 134.

[§] From 1843 to 1858 the temperature at Guernsey during the month of January has ranged from 24°.5 to 54°.5, and averaged 43°.6.—AMSTED, Channel Islands, pp. 134—7.

[&]quot;Some years ago, a level connected with mining operations then going on,

		L	e Pot.					Depth below the surface. fms.	Relation to the sea-level. fms.	Temp.
Modera	te stre	am of fre	sh wa	ter, c	out of	the r	ock	47.	12· A*	54.
	,,		,,	,	**	U	ode	65.	4· B*	56•
		Sark's	Норв :	Mine						
Small st	tream	of fresh w	ater,	out o	f rock	and	lode	24.	4. ₩	55· 5
Modera	te "	brackis	h ",	,,	rock	•••	••••	44.	20·B	56· 2
Large	"	"	"	,,	"	•••	••••	54.	30·B	<i>5</i> 7∙2
Small	27	"	,, ,	"	lode	•••	••••	,,	,,	57·2
,,	"	,,	"	,,	rock	w.	••••	64.	40∙ ▲	58•
		ish, pump						,,	,,	56•

HERM.

The Herm mines were wrought, to a depth of thirty fathoms, in rocks composed mostly of white, pinkish, buff-coloured, and greenish felspar, mixed with much hornblende, and sometimes with quartz and mica.†

and opening out on the Port du Moulin, on the side of the island towards Guernsey, was found to remove the water from a well in D'Ixcart Bay, on the other side of the island."—Ansted, Chonnel Islands, p. 472.

^{*} A denotes distance above the sea-level.

B .. below .. .

^{† &}quot;A beautiful white and black granite rock forms the hard back bone [of Herm]; and may be recognized at intervals, around the coast. * * * This granite is intersected by many wide veins, extremely variable in their nature, but generally either soft or readily decaying. * * * There is one at the back of the island of very large size, running across more than one projecting headland, nearly in a south-westerly direction, consisting entirely of black micaceous rock. * * There are other veins of soft clay, and some of decomposing greenstone. * * Traces of copper are said to have been found in veins in the granite of Herm; and mining operations were at one time commenced. The chief mineral product of the island is, however, its granite; [but] it is hardly equal to the best black Guernsey granite for paving and curbstones."

ANSTED, Channel Islands, pp. 63,—6, 263.

Two lodes respectively—

bear 10'-16° N. of E.-S. of W., * dip N., and measure 5-30 feet in width;
,, 24° W. of N.-E. of S., * ,, E., ,, 2-4 ,, ,,

Both these,—and the numerous (branches) veins which separate from, and re-unite with, them, in various parts of their range,—also contain great quantities of felspar, hornblende, and quartz; iron-pyrites abounds, and small (bunches) masses of yellow copper-ore occur at intervals.

As Herm and Guernsey are but six miles apart, they can scarcely differ much in climate.†

The undermentioned observations were made in February, 1841; viz.—

				Depth below the surface. fms.	Relation to the sea-level. fms.	Temp.
Well of	fresh water ‡		• • • • • • • • • • • • • • • • • • • •	Surface.	10· A	48·7
Small st	ream of fresh	water ou	t of lode	8.	2· B	49.5
Large	"	"	rock at some distance	"	,,	53·
Small	,,	,,	lode	12.	8 B	55·2
Minute	"	" jett ir	ing out of <i>lode</i> with- n a short distance	, ,	,,	56-

IRELAND.

COUNTY OF WICKLOW.

The mines of Connorree, Cronebane, Tigrony, Ballygahan, and Ballymurtagh have afforded enor-

^{*} In 1838 the Magnetic declination was about 24° W. Ross, Phil Trans., CXXXIX. p. 208. SABINE, Ibid, Pl. XIV. Ante, p. 531, Note *.

[†] Ansted, Channel Islands, pp. 134,-7. Ante, p. 735. Note §.

^{‡&}quot; Herm has good fresh water in natural springs, and in two places there is running water."—ANSTED, Channel Islands, p. 68.

mous quantities of iron-pyrites mixed with slaty matter, quartz, and various ores of copper,* from several beds of different widths which conformably interlie schistose rocks, presumed to be portions of the Silurian system.

As Ovoca lies between Dublin and Courtown, it probably differs but little from them in climate.

Localities.	Depth below the	Relations to the	Temp.	1840.
	surface. fms.	sea-level. fms.	May.	Nov.
Connorres Mine.			•	•
Water, a well	Surface	100∙ ▲	••	45.
,, a moderate stream out of clay-slate .	54.	75· A	••	49.5
,, ,, the Sulphur- course	27	,,	••	50.
,, pumped to the surface from	,,	,,	••	49.
Cronsbane Mine.				
Water flowing from a hole bored in the Sulphur-course	72·	2·B	54·5	
,, pumped to the adit (16 fms. above the sea from		22·B	55 ·5	

COUNTY OF WATERFORD.

The mine of Knockmahon has been wrought, both

^{*} Henry, Phil. Trans., XLVII. pp. 500-3. Journal des Mines, No. XVI. pp. 80-5. Weaver, Geol. Trans., v. pp. 173-8, 213-30. Haughton, Journal of the Geol. Soc. of Dublin, v. pp. 280-2. Smyth, Records of the School of Mines, 1. pp. 370-97, Mahon, The Mines of Wicklow, pp. 35-75. Ante, pp. 540-69, Table XVIII.

[†] Lloyd, Trans. Royal Irish Academy, XXII. p. 416.

¹ Ibid, p. 424.

⁶ Ibid, p. 422.

inland and beneath the sea, in greyish-green, greenish-black, and mottled fossiliferous slates, interlaid by massive rocks of felspar, quartz, and chlorite, as well as by thin beds of ferruginous conglomerate. The lodes—which have been very productive—consist, in great measure, of quartz, slaty matter, calcareous-spar, and chlorite, associated with earthy-brown iron-ore, iron-pyrites, earthy black copper-ore, vitreous copper, malachite, and copper-pyrites. The cross-veins, which intersect both the rocks and lodes, are composed, mostly, of slaty-clay, and disintegrated felspar; but, at intervals, they contain spheroidal masses of quartz.

At Dunmore, some 12 miles E. of *Knockmahon*, the mean temperature of the year 1851 was. 51.6;§ & "Waterford, "15 "N.E. ", the mean temperature from 1860 to 1868 was 50.3.

[§] Lloyd, Trans. Royal Irish Academy, XXII. pp. 416,-24.

Observat	ions made	at Nev	vtown, Wat	erford	;-	
Years.	3	<u> </u>	m. 1	Minima	m.	Mean.
1860		. 78°·		. 14°·		50°•
1		. 82·		. 25.	*******	52.2
2		. 76.		. 20		50.6
3		. 80		. 28.		50.3
4		. 81·		. 19		48.9
5		. 86		. 20		48.6
6	•••••	. 85		. 20.	•••••	49.2
7		. 79		. 16.		<i>5</i> 0·4
8	•••••	. 86.		. 28		52·3
Extrem	es	86.		14.		
Mean	••••••	_	•••••	_	•••••	50·3

R. J. GREER, Esq., of Newtown, Waterford, MSS.

^{*}Weaver, Geol. Trans., v. p. 248. Du Noyer, Explanation to accompany Sheets 167, 168, 178, 179 of the Geological Survey of Ireland, p. 57. Baily, Ibid,

⁺ Hore, Ibid, p. 81. Du Noyer, Ibid, pp. 81-2. Ante, pp. 594-8, Table XIX.

[#] Ante, pp. 598-9, Table XIX.

The undermentioned observations were made at Knockmahon; —

	Depth below the surface, fms,	Relations to the sea-level, fms.	Temp.
Water, a well (29th April, 1840)	1.6	3, ▲	48.
,, issuing from the clay-slate and the lode	16.	8. ▲	48.7
,, accumulating from several small streams out of the lode	112-	93·B	57·5
,, pumped to the adit (3 fms. above the sea) from	,,	,,	50-5

COUNTY OF CORK.

The Bearhaven mines are opened in rocks, of the

*" Thermometers were placed, in August 1843, in the deepest part of Knock-mahon Copper Mine; * * * one being sunk three feet into the rock, and another into the lode at a depth of [129 fathoms] from the surface. A thermometer * * * was hung in the gallery or level where these were placed. * * * "These mines are in lat. 52° 8' N. and the mean annual temperature at the surface calculated by the usual formula would, therefore, be 50° 026.

"The general average of the thermometers at the depth of [129 fathoms], and the maxima and minima, were as follows:—

In air Average, 57·176 .. Maximum, 58·5 .. Minimum, 56·25 , rock or country . , 57·369 .. , 58·5 .. , 56·25 , lode , 57·915 .. , 58·5 .. , 56·25

"Taking the temperature of the rock thus determined as the general average it shows an increase of 7°343 for a depth of [129 fathoms], or deducting [16·4 fathoms] for the line of no variation, we have 1° for [15·3 fathoms]. It was found necessary to fix the instruments not far from being perpendicularly under the sea, the shaft being nearly on the edge of the cliff, which is here [11·6 to 12·5 fathoms] high. If therefore we should * * * consider the sea level as the surface, we shall have a depth of [116·6 fathoms] corresponding to [7°343 or 1°=15·8 fathoms].

"It seems to be fully established • * • that there was a gradual though a slight diminution of temperature as the observations proceeded. Thus the temperatures were

during the first half; of the period; in air 57.613.. in the rock 57.718.. in the lode 58.000, the second half; of the period; , , 56.697.. , 57.044.. , 57.675

OLDHAM, Report of the British Association (for 1844), II. pp. 221-2 (Abridged). Carboniferous slate series, composed of siliceous matter mixed with chlorite, talc, or some similar mineral: at intervals, however, they contain small quantities of the carbonate of lime; whilst all parts of them are intersected by minute veins of quartz. Near rich portions of the lodes they are usually lilac, pale-buff, or dove-coloured, and of thick-lamellar structure; but elsewhere they are blue and fissile. The lodes consist mostly of quartz, but contain also chloritic or talcose matter, beside smaller proportions of the carbonates of lime and of iron; moreover, they include angular. (horses) masses of slate without number. Iron-pyrites, vitreous copper, and malachite occur now and then, but copper-pyrites is the prevailing ore. † A cross-vein -composed of slaty-clay in other parts of its rangebecomes highly quartzose as it intersects the Main (Mountain) lode. 1

At Castletownsend, some 40 miles E.S.E., the mean temperature of the year 1851 was 52.1;

" Cahirciveen, " 22 " N.N.W., " " " 52·3.§

The mean annual temperature at the Bearhaven mines, therefore, is probably much the same.

The temperatures observed at different depths have been,—

^{*} Jukes & Kinahan, Explanation to accompany Sheets 197 and 198 of the Geological Survey of Ireland, pp. 13, 20. Ante, p. 602, Table XX.

[†] Smyth, Explanations to accompany Sheets 197 and 198 of the Geological Survey of Ireland, pp. 30-3. Ante, pp. 603-8, Table XX.

¹ Ante, p. 608, Table XX.

[§] Lloyd, Trans. Royal Irish Academy, XXII. pp. 416,-23.

	Depths below the surface. fms.	Relation to the sea"-level. fms.	Temp.
Water cozing out of a rock N. of Main Lode	128	60· B	5 8∙5
••		72· B	61.2
,, pumped to the adit (20 fms. above the sea) from	,,	" i	56.

COUNTY OF KERRY.

The Ardtully mines were worked in, and between, reddish-purple, greenish-grey, yellowish-green, lead-coloured, or mottled, slates; † and greyish Carboniferous limestone in which crinoidal remains are not uncommon.‡ Small beds of slaty-clay, quartz, and earthy brown iron-ore, slightly sprinkled with iron-pyrites and yellow copper-ore, occur in the slate; § whilst irregular bands of calcareous-spar—here and there charged with grey and purple copper, slightly mixed with copper-pyrites, and enclosing, at intervals, small bodies of peculiar (? organic) character—merge, after short courses, in the limestone.

The principal

^{*} Kinahan, Explanations to accompany Sheets 197 and 198 of the Geological Survey of Ireland, p. 20, Fig. 3.

[†] Jukes, Du Noyer, & Willson, Explanations to accompany Sheet 184 of the Geological Survey of Ireland, pp. 20—4. Haughton, Journal of the Geological Society of Dublin, vi. p. 210. Ante, pp. 613—15.

[‡] Haughton, Journal of the Geol. Society of Dublin, vi. p. 208. Jukes, Du Noyer, & Willson, Explanations to accompany Sheet 184 of the Geol. Survey of Ireland, pp. 20—4. Ante, p. 613.

[§] Ante, pp. 612,-15.

^{||} Haughton, Journal of the Geol. Soc. of Dublin, vi. p, 213. Du Noyer & Willson, Explanations to accompany Sheet 184 of the Geol. Survey of Ireland, p. 37. Ante, pp. 518—19.

metalliferous deposit, however, intersects the slate in one part of its range, but separates the slate from the limestone in another. Where both sides (walls) are of slate shallow portions of the matrix consist of slaty clay mingled with earthy brown iron-ore, enclosing nodules of hematite, and angular masses of slate often encrusted with copper-pyrites, together with grains of purple, grey, and native copper; at greater depths siliceous slate—the principle ingredient—is frequently veined with quartz and speckled with copper-pyrites; moreover, where opposite sides of the deposit are bounded by rocks of different kinds, that portion of it which adjoins the slate-although rather richermaintains, in other respects, its normal character; at its contact with the limestone, on the contrary, it comprehends ill-defined beds of grey limestone and calcareous spar which embed considerable quantities of grey and purple copper, with smaller proportions of copper-pyrites.*

Kenmare is about thirty miles N.N.W. of Castletownsend, and about twenty-eight " E. by S. of Cahirciveen; but, inasmuch as it is less open to the ocean and more enclosed by mountains than they are, any difference between its mean temperature and theirs,† may, perhaps, be rather in defect than excess.

Whilst the Ardtully mines were deepened, observa-

^{*} Haughton, Journal of the Geol. Soc. of Dublin, vI. pp. 212—13. Du Noyer, & Willson, Explanations to accompany Sheet 184 of the Geological Survey of Ireland, p. 37. Ants, pp. 616—19, Table XXI.

[†] Lloyd, Trans. Royal Irish Academy, XXII. pp. 416,-23. Ante, p. 741.

tions were made, at various times, in different parts of them, with the undermentioned results.

1	Depth below the	Tempe	rature.
Localities.	surface.	1840. October.	1841. June.
North, Engine, or Ardtully, lode.			
Water, a small stream cosing out of slate in the N. side (wall) (a)	17.	50.6	•
,, ,, ,, lode W. (b)	,	51-25	
,, , a large stream out of slate (a)	20.		58.
,, , ,, ,, lode (b)	2 2·	51.25	
,, , ,, ,, (ð)	27·		55.
,, , pumped to the surface in 1840 from	22.	51.	
,, ,, ,,, 1841 ,,	27.		58.

ENGLAND.

CORNWALL.

THE CARADON DISTRICT,-

which rises from about 600 to 1,200* feet above the sea,—comprehends rocks of granite,† slate,‡ elvan,§

[•] Mac Lauchlan, (De la Beche's), Report on the Geology of Cornwall, Devon, and West Somerest, pp. 14, 18. Ante, p. 696.

[†] Boase, Cornwall Geol. Trans., IV. pp. 170, 209—10. De la Beche, Report on the Geology of Cornwall, &c., pp. 157,—9. Whitley, Reports of the Royal Institution of Cornwall, XXXII. p. 31. Thomas (Charles), Remarks on the Geology of Cornwall and Devon, p. 15. Webb & Geach, History and Progress of Mining in the Caradon and Liskeard District, p. 67. Holl, Quarterly Journal of the Geol. Society, XXIV. p. 440. Western Daily Mercury, No. 2,463 (28th May, 1868), p. 2. Ante, pp. 656—60,—62—66.

[‡] Rogers, Cornwall Geol. Trans. II. pp. 218—20. Boase, Ibid, IV. p. 208. De la Beche, Report on the Geology of Cornwall, &c., p. 79. Giles, Cornwall Geol. Trans, VII. pp. 155—6,—8. Webb & Geach, History and Progress of Mining in the Caradon and Liskeard District, p. 67. Holl, Quarterly Journal of the Geol. Society, XXIV. p. 444. Ante, pp. 656—60,—67—70.

[§] Boase, Cornwall Geol. Trans., IV. pp. 209-10. De la Beche, Report on the Geology of Cornwall, &c., pp. 159,-88,-85. Giles, Cornwall Geol. Trans., VII.

and greenstone. The lodes wrought at South Caradon, West Caradon, and Gonamena, on the S., traverse granite and elvan, abound in fluor, and yield only copper and copper-ore; † whilst those opened at Marke Valley and the Phænix mines, towards the N., intersect slate, granite, and elvan, contain no fluor, but afford the ores of both copper and tin, Crossveins occur in several parts of the district; § but neither of them has been traced throughout its entire breadth.

The mean temperature of Plymouth, some sixteen miles S.E., deduced from 43,824 horary observations, made at about 60 feet above the sea, during five years, was 52°081.

At different depths, in various parts of the Caradon district, the undermentioned temperatures were observed:—

Localities.	Rocks.	Depth below the surface. fms.	Relation to the sea-level, fms. ¶	Temp.
Cheesewring Hotel. Water, in a deep well, full to within 4 fms. of the surface (25th July, 1867)	Granite .	4.	180· A	50·9

pp. 158, 201. Webb & Geach, History and Progress of Mining, &c., pp. 33,—6. Holl, Quarterly Journal of the Geol. Society, xxiv. pp. 415,—41,—45. Ante, pp. 660,—1.

[•] Rogers, Cornwall Geol. Trans., 11. pp. 218—21. Boase, Ibid, IV. pp. 207—9. De la Beche, Report on the Geology of Cornwall, &c., p. 79. Giles, Cornwall Geol. Trans., VII. pp. 156,—8. Holl, Quarterly Journal of the Geol. Society, XXIV. pp. 421,—23,—44. Ante, pp. 655,—71.

[†] Webb & Geach, History and Progress of Mining in the Caradon and Liskeard District, pp. 31,-6, 51-3. Ante, pp. 678-80, Tables XXIII., XXIV.

[†] Webb & Geach, History and Progress of Mining in the Caradon and Liskeard District, pp. 24—31. Ante, pp. 676—80. Tables XXV.—XXVI.

Whitley, Geological Map of the Caradon Mining District. Webb & Geach, History and Progress of Mining in the Caradon and Liskeard District, pp. 25, 31, -3, -5, 52. Ante, pp. 681-5; Tables XXIII.-IV.,-VI.

Harris, Reports of the British Association, VII. pp. 24-5, Pl. X.
Approximate.

Localities.	Rocks	Depth below the surface. fms.	Relation to the sea-level, fms.	Temp.
Gonamena.				
Water in the adit, a very large stream out of the (Country) rock and lode; taken for household use in the neighbourhood, as it flows out at the surface		20-	130· A	5η4
South Caradon.		,		
Water, a very large stream out of the Little Cross-course ,, pumped to the adit (14 fms. below			96∙ ▲	51.4
the surface) from	,,	128.	12· A	61.5
The Phænix Mines.		ł I	1 1	
Water, cozing out of the (Country) rock and lode at the bottom of the mine.		146·	14· A	67.
", pumped to the adit (26 fms. below the surface) from		"	,,	52·6
Marke Valley.				Į
Water, a small stream out of Sarum lode (back of level)	Slate	106-	14· A	60-5
,, , ,, (bottom ,,)	"	"	,,	62.5
,, , , , (end ,, E.)	",	79	,,	70.
,, , a large stream out of Marke lode, W.	Granite.	,,	,,	68-3
,, , pumped to the adit (26 fms. below the surface from		,,	,,	69-8

THE MENHENIOT DISTRICT,

rather more than three hundred feet above the sea,* comprehends an extensive area of—more or less calcareous †—clay-slate,‡ which contains, at intervals,

^{• &}quot;The Menheniot station on the Cornwall Bailway is 261.5 feet above the sea-level."—J. D. Sheriff, Esq., C.E., Engineer of the Cornwall and West Cornwall Bailways, MS.

[†] Boase, Cornwall Geol. Trans., IV. p. 212.

[#] Henwood, Reports of the Royal Institution of Cornwall, xxxIII. (1851),

numerous small cavities filled with earthy ferruginous matter (? of organic origin *); in some places the slate encloses, but in other it is interlaid by rocks of felspar and hornblende,† occasionally of schistose, though usually of massive, structure.‡ The only lode yet discovered in the neighbourhood has afforded, and—at more than two hundred and fifty fathoms deep—still continues to afford, great quantities of argentiferous galena § and smaller proportions of other, less valuable, ores. Barren (flucans) veins of clay || (heave) displace the lode in several parts of its range.

During the years 1833—1837 the mean temperature at Plymouth, some eleven miles S.E., was 52°.081.¶

Streams derived from various parts of the slate series have—at different times—shown the temperatures hereafter mentioned.

p. 39. Sedgwick, Quarterly Journal of the Geol. Soc., VIII. pp. 5, 17, 146. Giles, Cornwall Geol. Trans., VII. p. 201. Webb & Geach, History and Progress of Mining in the Caradon and Liskeard District, p. 38. Salmon, Mining and Smelting Magazine, II. p. 211. Holl, Quarterly Journal of the Geol. Society, xxiv. p. 423. Ante, pp. 700—4.

[•] Ante, p. 700.

[†] Rogers, Cornwall Geol. Trans., 11. p. 221. Boase, Ibid, IV. p. 211. Henwood, Reports of the Royal Institution of Cornwall, XXXIII. p. 39. Giles, Cornwall Geol. Trans., VII. p. 201. Ante, pp. 701—2.

^{\$} Ante, pp. 701-2.

[§] Henwood, Reports of the Royal Institution of Cornwall, XXXII. p. 40. Giles, Cornwall Geol. Trans., VII. p. 203. Salmon, Mining and Smelting Magazine, II. p. 218. Webb & Geach, History and Progress of Mining in the Caradon and Liskeard District, pp. 26, 36. Ante, pp. 703—14; Tables XXVII.—VIII.

[§] Henwood Reports of the Royal Institution of Cornwall, XXXIII. pp. 40,—2. Webb & Geach, History and Progress of Mining in the Caradon and Liskeard District, p. 37.

Tharis, Reports of the British Association, VII. pp. 24-5, Pl. X. Ante, p. 745.

Localities	Depth below the surface, fms,	Relation to the sea-level, fms.*	Temp.
Liskeard.†			•
Water in a closed well at the London Inn } 1851, Sept 16th.	0.6	73· A	53 -2
" Dean's (closed) well 1867, July 30th.	1.	71· A	54.
Menheniot.			
Water in an open well, midway between the Church and Wheal Mary Ann 1867, July 29th.	Surface.	65∙ ▲	54· 7
South Wheal Trelawny.			
Water, a small stream out of veins	<i>5</i> 3·	about sea-	56.5
" , pumped to the adit (13 fms. deep) from	78.	20·B	56.5
Wheal Mary Ann.;			
Water, a large stream out of lode, bottom of the mine 1851, Sept. 9th.	98.	48· B	67-5
" pumped to the surface 1867, July 29th from	280.	280· B	64.5
Wheal Trelawny.			
Water, a moderate stream out of the lode N	68.	18· B	60.
,, a small stream	95.	45∙ B	6 5 ·
, a large stream out of the lode 8.	105.	55· B	65·
" pumped to the surface 1867, July 29th from	210	160·B	65.8

The Lanbeath and Saint Pinnock District rises from $150\,\parallel$ to, perhaps, 200 feet above the sea;

[•] Approximate.

^{† &}quot;The centre of the Parade at Liskeard is 425 feet above the sea."

ALLEN, History of Liskeard, p. 454.

[†] Within these sixteen years Wheal Mary Ann has been deepened 182 fms

^{§ ,, ,,} Wheal Trelawny ,, 105 ,, . | "Moorswater, the head of the Liskeard and Looe Canal, is 150 feet above the sea."—ALLEN, History of Liskeard, p. 454.

and consists of calcareous slates, which sometimes contain organic remains.* The lode at Herod's-foot—the only one yet wrought to advantage—has yielded, and still yields, an abundance of argentiferous galena, and bunches of copper-pyrites, beside smaller quantities of several other ores.† The lode is (heaved) displaced by a cross- (flucan) vein; which consists mostly of schistose matter, but is, at intervals, thinly sprinkled with ore.‡

In climate, *Herod's-foot* can scarcely differ much from Plymouth, Caradon, and Menheniot.§

The undermentioned temperatures have been observed in different parts of the district:—

Localities.	Depth below the surface. fms.	Relation to the sea-level, fms.	Temp.
Duloe.			•
Water in an open well at Benoke 1851, Sept. 18th	Surface	33· A	55 ·4
Saint Keyne.			
Water in an open well ¶ 1851, Sept. 18th	Surface	25· A	55.4
Herod's-foot.			
Water, a large stream out of the lode S. 1851, Sept 15th	187	110·B	67-5
,, pumped to the surface 1867, July 27th from	160-	135· B	61.

^{*}Giles, Cornevall Geol. Trans., vii. pp. 97—9, 171. Peach, Ibid, p. 104. Sedgwick, Quarterly Journal of the Geol. Society, viii. pp. 5, 17. Holl, Ibid, xxiv. p. 423. Ante, p. 700.

[†] Giles, Cornwall Geol. Trans., vII. pp. 201—3. Salmon, Mining and Smelting Magazine, II. pp. 211—17. Webb & Geach, History and Progress of Mining in the Caradon & Lisheard District, pp. 16—18. Ante, pp. 705—15; Table XXIX.

¹ Ante, pp. 715-18, Table XXIX.

[§] Harris, Reports of the British Association, VII. pp. 24-5. Ante, pp. 745,-7. Approximate.

T Carew, Survey of Cornwall (1602) f. 180. Norden, Speculi Britannia Pars. p. 86. Souther, Poetical Works. p. 666. Blight, Ancient Crosses and other Antiquities in the East of Cornwall, pp. 90—2.

SHROPSHIRE.

At Eardiston, some five miles S.E. of Oswestry, the New Red Sandstone and a band, varying in width from a few inches to perhaps five feet, by which it is intersected, both consist, in great measure, of granular quartz; but, whilst the former is tinged, more or less deeply, by various proportions of ferruginous matter, the latter contains, at intervals, great quantities of earthy brown iron-ore, sometimes largely mixed with the green carbonate of copper, and occasionally thinly sprinkled with grey copper-ore. At a depth of sixteen fathoms, however, the iron-ore is replaced by blue clay, when all trace of copper-ore suddenly disappears.

A narrow ferruginous cross-vein intersects the whole formation; but occasions no (heave) displacement.

At Whittington, about five miles N.N.W. of *Eardiston*, the temperature, — between March 1842 and February 1843,— ranged from 11° to 81°, and averaged about 49°.7.†

[†] At Whittington, within five miles of Eardiston, the extreme and mean temperatures from March 1842 to February 1843, were—

•	1	9 д.м		. :	В Р.М		1 9	9 г.м	_		Registe	
Months.				Max.								
1842. March	50.3	35.7	14.	°55.	40.7	48·1	49·	36.	41.	61·	28.	43·1
April			45.8			53.8			42.3		24	44.2
May	60.5		54.2			59.8		39.3	48.2	69	33.	52.1
June	69.3	55.	62·	79.2	59.7	68.	64.	47.	55.9	81.	34.	59.8
July	69.	54.2	61.4	74	55.3	66.1	61.	49.	54.9	75.	41.	59.1
Aug	70.5	55.2	64.4	80.5	55.5	66.4	63.	45.	58.6	81.	42.	62.3
Sept	68.	50.	57.3	70.2	51.7	61.4	63.	42.5	56.6	74.	34.	55.4
Oct	55.	32.	45.7	57.5	40.	49.7	49.	32.2	41.8	59.	24.	44.5
Nov	48.5	34.	40.7	50.5	38.	43.4	47.3	30.	37.8	52·	28.	41.
Dec	55.2	31.	44.2	<i>5</i> 6·2	37.5	47.3	55.3	31.5	44.9	59·	27.	45.2
1843. Jan	52.	22.5	38.5	<i>5</i> 3·	32.2	41.8	51.	27.	38.8	56.	15.	39.1
Feb	42.3	19.	34.4	47.5	28•	36.8	46.5	18.	33.7	51.	11.	34-4
Extremes	70.5	19-	-	80· <i>5</i>	28.		64.	18.		81.	11.	
Means			49.4			53.5		۱	46.2	l	١	48.34

^{49.7} The Reverend C. A. A. Lloyd, MS.

^{*} Murchison. Silurian System, pp. 39, 298. Ante, pp. 515-16.

At Eardiston on the 17th—18th of November, 1842—

It may, perhaps, be desirable, to place the facts, already described, in such various points of view, as may disclose their respective peculiarities.

The mean depths of the mines in each district;—

```
,, ratios of increase in temperature, expressed in fathoms of descent requisite to an elevation of one degree; • ... annual temperatures at the surface... }
```

are set forth in the following columns:—

	1]	Surface.		
Countries.	Provinces & Districts.	Mean depth. fms.	Mean temp.	Mean ratios. fms.	Mean annual temp.
Сиці	Chaffarcillo	155	69°2	21.6	64· †
BRAZIL	Minas Geräes	53	67.9	26.2	60-49 ‡
United States	Virginia	15	<i>5</i> 6·8	§ .	55· †

For the foregoing extracts, from a *Meteorological Register* which extends from March 1842 to July 1851, the writer is indebted to the Reverend Albany Rossendale Lloyd of Hengoed and George James Symons, Esq., F.M.S., Editor of "British Rainfall."

^{*} Henwood, Cornwall Geol. Trans., v. p. 404.

[†] Keith Johnston, Atlas of General Phonomena, Pl. XVIII. Ante, p. 724.

¹ Ante, pp. 726-8,-9,-32. Tables XXX.-XXXI.

[&]amp; One observation only.

ı	U	Surface.		
rovinces & Districts.	Mean depth. fms.	Mean temp.	Mean ratios. fms.	Mean annual temp.
ark & Herm	87	56̂∙5	10-	51.6 ◆
Vicklow, Waterford, Cork, & Kerry	} 57	58-4	14·1 {	50-3 † 52-3 †
Jornwall	92	62·5	5.8‡	52·08 §
hropshire	15	54.	ſ	49-7 T
	ark & Herm Ticklow, Waterford, Cork, & Kerry ornwall	rovinces & Districts. Mean depth. fms. ark & Herm 37 loklow, Waterford, Cork, & Kerry 357 ornwall 92	rovinces & Districts. Mean depth. fms. Mean temp.	depth. fms. Mean temp. ratios. fms. ark & Herm 37 56.5 10. Toklow, Waterford, Cork, & Kerry 57 53.4 14.1 ornwall 92 62.5 5.8

Other details appear in Table XXXII.

The mean—depths,—temperatures,—and ratios in which the temperatures increase with the depths, in

[‡] In the principal mining districts of Cornwall and Devon four hundred and fifteen observations afforded the undermentioned results:—

Moan depth, fms.	Mean temp,	Mean ratios.	Districts.	Mean depth, fms,	Mean temp.	Mean ratios.
95	57 [.] 84	14.3	Camborne, &c.	98	62 ⁹ 13	10-6
129	63· <i>5</i> 6	11.2	Redruth, &c	132	71.87	<i>5</i> ·8
76	63-87	7.7	Saint Agnes	99	65-91	8.4
101	68-4	7-4	Saint Austell	186	70-62	5.
134	66-66	8.8	Tavistock, &c	72	59:07	8-9
	95 129 76 101	95 57°84 129 63°56 76 63°87 101 63°4	depth. fras. Mean ratios. 95 57.84 14.3 129 63.56 11.2 76 63.87 7.7 101 63.4 7.4	depth fine. Mean ratios. Districts. 95 57.84 14.3 Camborne, &c. 129 63.66 11.2 Redruth, &c. . 76 63.87 7.7 Saint Agnes . 101 63.4 7.4 Saint Austell .	depth fine. Mean ratios. Districts. depth fins. 95 57.84 14.3 Camborne, &c. 98 129 63.56 11.2 Redruth, &c. 132 76 63.87 7.7 Saint Agnes 99 101 63.4 7.4 Saint Austell 136	depth fine. Mean ratios. Districts. depth fins. Mean temp. 95 57.84 14.3 Camborne, &c. 98 62.13 129 63.66 11.2 Redruth, &c. 132 71.37 76 63.87 7.7 Saint Agnes 99 65.91 101 63.4 7.4 Saint Austell 136 70.62

Mean depths..... 112 fms.;-

HENWOOD, Cornwall Geol. Trans., v. pp. 402,-6.

Ansted, Channel Islands, p. 140. Ante, p. 785.

[†] Lloyd, Trans. Royal Irish Academy, XXII. pp. 416,—22,—3,—4. Ante, pp. 788,—9,—41,—8. R. J. Greer, Esq., MS. Ante, p. 739.

^{,,} temperatures -... 66°88;--

^{,,} ratios 6-8 fms.

[§] Harris, Reports of the British Association, VII. pp. 24,—5. Ante, pp. 745—7.

¶ One observation only.

The Reverend C. A. A. Lloyd, of Whittington near Oswestry, MS. Axie, p. 750.

the different rocks of the several districts already mentioned, appear in the following pages:—

Rocks.	depth, fins,	Mean temp.	Mean ratios,
New Red Sandstone*	15	84.	One observation only.
• "New Red Sandstone [is] very much affected in saturated with moisture. Two blocks,—of which the second was the harder mentioned powers of conduction when in different conduction to the second was the harder mentioned powers of conduction when in different conduction when in different conductions were second to the second was the harder mentioned powers of conduction when in different conductions were second was the harder mentioned powers of conductions when it is not conducted to the second was the harder mentioned powers of conductions when it is not conducted to the second was the harder mentioned powers of conductions when it is not conducted to the second was the harder mentioned powers of conductions when it is not conducted to the second was the harder mentioned powers of conductions when it is not conducted to the second was the harder mentioned powers of conductions when it is not conducted to the second was the harder mentioned powers of conductions when it is not conducted to the second was the harder mentioned powers of conductions when it is not conducted to the second was the harder mentioned powers of conductions when it is not conducted to the second was	-,—poss	essed t	
Dry 25	49		
Saturated with moisture 60			
Hopkins, Phil. Trans., Oxl.		808,	18,—19.
At the mine of Monk-Wearmouth—which was sunk, 284 fms. beneath the surface,		•	•
249-5 ,, ,, sea, to the coal-seams which underlay the magnesian limes the mean temperature at the surface			ı—
", the depth of 264 fms. If, therefore, the depth of the invariable plane be taken at 16.6 ,, we			se of tem-
perature equal to one degree for 9.9 fms. of descent. PHILLIPS, London and Edinburgh Phil. Mag., v.	op. 446-	—51 (A	bridged).
The coal-mine of Torcy (Department of Saône et Lo		(
was opened at	ove the	•	
the total depth thus being 325.2 "			
The annual mean of the climate			
,, temperature maintained at 303 fms. deep, in an aban of the mi	doned 1	part ?	,, 81°.
The works of Mouillelonge, some two miles distant,	_		
were commenced at 1	175·8 fb	ıs. abov	e the sea ;
and—having pierced the New Red Sandstone— } entered the Coal-measures ; "	20.7	, belov	7 ,, ;
		, deep	
reaching eventually a depth of 207.7 fms. below the sea, or		su	rface,
where the temperature	• • • • •	W	as 100°.9.
Now the difference			
between the annual and the temperature at 308 fms. in the mean at Toroy,		WAS	39° ·5 (1);—
, 385 fins. at Torcy ,, · · , , , , , , , , , , , , , , , ,	usiongs		52°·4 (3); 19°·9 (3).

Rocks.	Mean depth, fms.	Mean temp.	Mean ratios.
Limestones alternating	155	69 ·2	21-6

The ratio in which the temperature increases with the depth was-

in the first case, one degree for 9.2 fms.;—

" second,, " 8.5 ";—

" third ", " 7.2 ".

(WALPERDIN, Comptes Rendus). SMYTH (Annual Address), Quarterly Journal of the Geol. Soc., XXIV. pp. lxxix.—lxxx. (Abridged).

The Duckinfield colliery afforded rare opportunity for observing, in two shafts, the gradual increase of temperature with depth. Of the observations made by F. D. Ashley, Esq., the proprietor, a synopsis is presented in the following columns:—

FIRST SHAFT.					SECOND SHAFT.					
Extre	mes.	Means.			Extre	mes.	Mea	1		
Depth.	Temp.	Depth fms.	Temp.	Ratios.	Depth. fms.	Temp.	Depth. fms.	Temp.	Ratios.	
2·8 115·5 117·3 146·5 150· 179· 186·5 216·5 223· 241·7 243·5 266·5 298·5	51. 57.7 58. 59.5 59.9 62.5 64. 66.5 67. 67.2 67.7 69.7 69.9 71.5 72.2	\$\\ \begin{align*} \text{fins.} \\ \choose \ \ \choose \	54°4 58°1 60°7 65°2 67°1 68°8 71°5 72°1	18·5 12·9 9·3 15·4 13·6 9·6	83·7 119·7 127· 147·5 154· 179· 191·2 218·2 227·7 233·5	58. 59. 58.2 60. 62. 63.5 65.5 66.5	fms. } 95-8 } 136-6 } 164- } 204-5 } 230-6	58°2 59°4 61°3 64°	34· 14·4 15·	
322·7 325·5 358·5	72·2 72·5 75·	338.3	74.	15.3						

Rocks,	Mean depth. fms.	Mean temp.	Mean ratios.
Felspathic and Hornblendic Rocks	87	55°5	10-

It appears therefore, that the rocks in these neighbouring shafts maintain different temperatures at corresponding depths; and that at various parts—of even the same shaft—the temperatures increase, with the depths, in widely different ratios.

At the first shaft, indeed,

of these differences, however, only one amounted to one degree and a quarter, whilst most of the others were much smaller.

Nevertheless, that the temperature increases at an average of one degree for a descent of

14.8 fathoms in the first shaft, and of

16.8 ", " second,,,

is indisputable.

FAIRBAIRN, Report of the British Association, 1861, Part II. pp. 53-6. (Paraphrased and abridged.)

The Rose Bridge collieries, at Ince near Wigan, have afforded opportunity for the undermentioned observations:—

From 80.5 to 403 fathoms therefore the increase of temperature averages one degree for 11.1 fathoms.

For this interesting record the writer is indebted to
JOHN ARTHUR PHILLIPS, ESQ.

In the Coal-mines of Virginia, which are believed to be of the Oolitio period (LYELL, Quarterly Journal of the Geol. Soc., 111. p. 261), Professor W. B. Rogers observed that

D'ARCHIAC, Histoire des Progrès de la Géologie, I. p. 71.

Rocks.	Mean depth. fms.	Mean temp.	Mean ratios.
Clay-slate* Jacotinga †	70	61°8	11.8
Jacotinga †	40	67:8	_
Talcose, micaceous, and chloritic slates	18	62-9	‡
Talcose, micaceous, and chloritic slates	79	59·5	<i>5</i> ·8
			1

The mean—depths,—temperatures,—and rates at which the temperatures increase with the depths—of the mines which yield different metals and ores (Table XXXIV.), are—

Metals and Ores.	Mean depth. fms.	Mean temp.	Mean ratios.
Gold	51.	6 7 ·5	23.
Silver	155•	69.2	21.5
Lead	72.	60.7	8.4
Copper §	43.	53· 7	15.9

[•] From one hundred and thirty-four observations, in the mines of Cornwall, between 1830 and 1837, it appeared that the slate was about 3°.9 warmer than the granite at the same depth. But four hundred and fifteen observations made in Cornwall and Devon from 1830 to 1843 showed—

that at a mean depth of 116 fms. the temperatures of the slate averaged 68°-89; , 94 ,, , , granite ,, 60°-35.

Henwood, Reports of the British Association, vl. (1837) Part II. p. 37; Cormoall Geol. Trans., v. p. 403.

,, 18 ,, 29, ,, 26·3, ,, — ,, 77°·8—92°·5 ,, 87°·15; an increase at the rate of one degree in (2·12) little more than two fathoms.

These temperatures so greatly exceed those observed, at corresponding depths, in any other mine, and the vertical range of observation is so small, that they have not been used in deducing the foregoing means.

[†] The mine of Agoa Quente—notwithstanding its depth hardly exceeded thirty fathoms—discharged more than three hundred cubic feet of water per minute. Of this enormous stream—

[‡] Two observations only.

^{§ &}quot;Fox observed [that] * * * tin veins usually shewed themselves colder

Motals and Ores.	Mean depth. fms.	Mean temp.	Mean ratios.
Copper and Tin*	114	65.7	— †

But whilst Morro Velho and Gongo Soco have been rich in gold, by far the larger part of the auriferous deposits have consisted of iron-pyrites in one, and of specular iron-ore in the other. Moreover, in the lodes which yield copper-ore,—whether mixed or unmixed with the oxide of tin,—iron-pyrites always abounds.

Between the temperatures and ratios observed, at various depths, in mines which have afforded similar metals and ores in rocks of different character, as well as between the temperatures and ratios noticed in such other mines as have yielded different metals and ores in rocks of like nature, the following comparisons have been made (Table XXXV.):—

The following are the mean depths at which observations were made, and the mean temperatures observed in the *lodes* affording different ores in the principal mining districts of Cornwall and Devon.

Lodes.	Mean depth, fms.	Mean temperature.
Copper	140	72°-89
Copper and Tin	74	61.45
Tin	92 .	60-67

HENWOOD, Cormoall Geol. Trans., v. p. 404.

than those which yielded copper."—Annales et de Chimie et de Physique, xvi. p. 80; Edinburgh New Phil. Journal, xxiv. p. 140.

[&]quot;The tin mines of the Souberg at Ehrenfriedersdorf also show a remarkably low temperature; indeed it is a prevailing opinion that stanniferous mountains are colder than others."—REICH, Beobachtungen ueber die Temperatur des Gesteine in verschiedenen Tiefen in den Gruben des Sachsischen Erzgebirges, pp. 87, 107. BISCHOFF, Edinburgh New Phil. Journal, XXIV. p. 140.

^{*} Ante, p. 756, Note §.

[†]Means of five observations, but all at the same horison.

•	METALS AND ORES.								
		Gold		L	LEAD		L	COPPE	R.
Rocks.	Mean depth funs.	Mean temp.	Mean ratios.	Mean depth fms.	Mean temp.	Mean ratios.	Mean depth fms.	Mean temp.	Mean ratios.
New Red Sandstone						••	15	<i>5</i> 4·	•
Felspathic & Horn- blendic rocks				4 8	56 ⋅8	13.5	30	54.7	9.4
Clay-slate	67	68·4	25.2	93	63.6	10.7	57	58·4	14-1
Jacotinga	42	67:3							
Talcose and Micaceous slate	18	62.9	t						
Granite		••			••	••	32	51-4	+
Means	51	67°-8	23.	72	60°-7	8.4	43	53°-7	15.9

Amongst the mines described in foregoing pages, those which have yielded silver and gold occupy high ranges of mountains, within the tropics; whilst such as have afforded the ores of other metals have been wrought in less elevated parts of temperate regions. Between the mean-depths,-temperatures,-and ratios in which the temperatures increase with the depths, in works thus differently situated, a comparison is offered —as well in Tables XXXII. and XXXVI. as—in the following columns:—

Metals and Ores.	Comparative elevation of surfaces, fms.	Mean depth. fms.	Mean temp.	Mean ratios.
Gold and Silver			67·7	30. ‡
Lead, Copper, and Tin	Less than 200	Ģ1	<i>5</i> 7·	8.9
Means		62	62.3	16.3

^{*} A single observation.

[†] The only observations have been made at the same horizon.

[‡] In the clay-slate of Morro Velho, temperature increases with depth much

As far as these observations extend, therefore, it appears that at considerable altitudes within the tropics, the temperature is higher than at corresponding levels below the surface at smaller elevations in temperate regions; but that the ratio at which it increases with the depth is much less rapid in the former than in the latter.

The mean—depths,—temperatures,—and ratios at which the temperatures increase with the depths, of the mines before mentioned, irrespective of their geographical positions, altitudes, rocks, metals and ores,—are the following:—

Extreme depth, fms.	Mean depth, fms.	Mean temp.	Mean ratios.*
Surface to 50	28	61:)
50 , 100	65	60.5	21.4
100 "150	122	65.4]
150 ,, 200	155	72 ·	5.
200 and beyond.	227	73.2	60.
Means	62	62.3	16.3

less rapidly than it has been found to increase in the similar rocks of Cornwall.

Hemwood, Proceedings of the Royal Geol. Soc. of Cornwall, 24th Oct. 1865.

At Morro Velho in Brazil the rate at which the temperature increases is but one degree for (33.3 fathoms) 200 feet.—SMYTH, Quarterly Journal of the Geol. Soc., xxiv. (1868) p. lxxxvi.; Ante, p. 727.

^{*}One hundred and thirty-four observations in the mines of Cornwall and Devon afforded—between 1830 and 1837—the following results:—

But at great altitudes in tropical regions—where the temperatures are above, whilst the ratio is below, the average,—and at smaller elevations in temperate climates,—where the ratios exceed, whilst the temperatures fall short of it,—observations have not been made

Rocks.						
	SL	LTB.		1	Granite.	•
Extreme depth. fms.	Mean depth, fms.	Mean temp.	Ratios. a	Mean depth, fms.	Mean temp.	Ratios. a
Surface to 50	35	57°	8.8	31	51°6	11•4
50 ,, 100	73	61.3	8.1	79	<i>55</i> ·8	5.6
100 "150	127	68·	4.3	183	65-5	6.6
150 " 200	170	78·	6.1	_	_	0.0
200 and beyond	221	85.6		237	81.3	
Means	•••••		6.5	•••••		6-9

Henwood, Thomson's Records of General Science, iv. (1836), p. 198; Reports of the British Association, vi. (1837), Part ii. p. 36.

"By burying the bulbs of different thermometers at various depths below the deepest excavations of mines" the undermentioned results were obtained:—

Mines.	Depth below surface. fms.	Temperatures.	Ratios from surface.
Levant	230	80.	7.6
Tresavean	262	82-	8.
Consolidated Mines	290	8 <i>5</i> ·3	8.3

Fox, Report of the British Association, VI. (1837, Part I.), pp. 134-7. (Abridged.)

"Upon the whole, I believe that * * the ratio in which the temperature augments in descending is greater in shallow than in deep mines."

Fox, London and Edinburgh Phil. Mag., XI. (1837), p. 523.

At a mean elevation of about 240 feet above the sea, the ground, at a depth of three feet, maintained, throughout the year, an average temperature of 49°-86. One hundred and seventy-seven observations, in different parts, but not the a These columns are now added.

in the same proportions at different depths. Thus,-

Extreme depth, fms.	Observations in elevated tropical regions.	Observations at moderate altitudes in temperate climates.
Surface to 50	24	17
50 ,, 100	5	12
100 " 150	4	10
150 ,, 200	2	
200 and beyond	2	
Totals	37	39

deepest, of many mines in Cornwall and Devon, exhibit increments of temperature equal to 10° each at intervals of about 47, 79, and 125 fathoms of descent.

Whilst fifty-three experiments in the deepest levels or accessible parts of mines show the rock, water, and air to preserve in round numbers,—
a temperature of 60° at 59 fms. below the

The following columns show the respective ratios of increase in temperature expressed in fathoms of descent requisite to produce an elevation of one degree; deduced from four hundred and fifteen observations in the mines of Cornwall and Devon:—

Depth.	Granite.	Slate.	Rocks,	Cross- courses.	Lodes.	Tin- lodes. fms.	Lodes yielding both tin and copper ores. fms.	Copper- lodes. fms.	Means.
Surface to 50	9.3	5.	5.8	8.2	6.	8.6	6.5	4.6	6.8
50 ,, 100	9.1	7·1	8.1	6.	8.3	7.3	6.4	8.5	7.6
100 ,, 1 <i>5</i> 0	8.3	8.3	6.7	11.	7.8	8.5	10.5	8.	8.7
150 ,, 200		4.4	3.7	4.9	6.3		8.	4.5	4.5
200 & beyond	7.5	6.5	9.5	3.9	5.2	5.1		6.5	6.4
Means	8.5	6.2	6.7	6.8	6.7	7.8	6.6	6.4	6.8

Henwood, Cornwall Geol. Trans., v. p. 406; (D'ARCHIAC), Histoire des Progrès de la Géologie, I. p. 69.

This preponderance of observation—
at less than 50 fathoms deep in elevated, tropical, regions;
from 50 to 150 " " in lower, temperate, countries;
and at all greater depths ... at great altitudes within the tropics,
accounts for the apparently higher temperature from
the surface to fifty—than from fifty to one hundred—
fathoms deep. Yet, whether unequal numbers of observations—in each of several countries so far apart,—
at altitudes so various,—in rocks so different,—and in
mines yielding so many metals and ores,—can afford
results accurately representing the mean temperatures

The temperatures observed in the rocks or lodes at the deepest levels, and ratios at which the temperatures increase with the depths, of mines in various parts of Cornwal are—

Mines.	Ores.	Rocks,	Date of ob- servations.	Depth.	Temp.	Ratios.
Botallack	Copper & Tin.	Slate	1837	188	79°·	6.5
		Slate	1853	255	87.	6-9
Levant	Copper & Tin.	Granite	,,	,,	74.	10.6
		Slate	1857	,,	85.	7.3
			1822	230	75.5	9-
Dolcoath	Copper & Tin .	Granite	1857	272	73·	11.8
	(another lode)	•••••	L ,,	,,	79.5	9.2
Tuesamean	Con	Granite	1837	262	82.5	8.1
Tresavean	Copper	Granite	1853	352	90-5	8.6
United Mines	Copper	Slate	1853	275	94.	6.3
Par Consols {	Tin	Slate	∫1837	128	74·	5.7
Tur Consols 3	Copper	Siace	1837	208	84.	6.1

In the United Mines, the temperature of the hot at 265 fms. deep, spring was 116° & the ratio of increase 1° in 3°9 fms.;

Fox, Reports of the British Assoc. for 1857 (Abridged and Paraphrased).

"The North or Hot-Lode of the Clifford Mines, formerly known as that of the

and ratios on any one vertical line, may, perhaps, be open to question.

The composition and structure peculiar to different strata afford greater or less facility for the ascent of water and vapour; which, co-operating with the conducting power proper, in various degrees, to all rocks,—aid as well in transferring towards the surface of the earth some portion of the heat maintained within it, as in determining to each formation its due distribution of temperature. Occasionally, however, this normal equilibrium is disturbed by the miner; * through

United Mines, is one of a group of east and west [copper-] veins which are encased in the clay-slate or killas, on the east of the granite hill of Carn Marth. * * The author found, in 1855, the chief spring welling upwards in a level 251.6 fathoms deep, with a temperature of 114°. * * In [1864, however,] these parts of the workings had been laid dry by the extension of deeper galleries, and the point of egress of the springs was along the rich lode, advanced [much] farther eastward. * * * [In this part of the works] the principal body of the upward-flowing water was to be seen rising * * on the north side of the magnificent lode of cellular, black-stained, cinder-like pyrites. The next level [at 270 fathoms below the surface] is advanced farther eastward by some 70 fathoms; the lode exhibited a good breadth of fina black-coated copper pyrites; and small feeders of water, issued mostly from the north, or hangingwall, almost scalded the fingers holding the thermometer, which marked 122°. * * At the bottom level, which is 275 fathoms deep; in its end the lode was narrow, and very impervious to water, but a little rill trickling from it showed a temperature of 121°. * *

Between my last two visits * * the point of issue of the hotter water had been deepened 30 fathoms, and the temperature was increased by 8°. This would give 1° for 3.75 fathoms. • • •

SMYTH, Mining and Smelting Magazine, vI. pp. 193-6; Reports of the British Association, for 1864, Part IL p. 70. (Abridged)

e"Numerous observations show that, whilst the conditions of the works on mines are unchanged, the temperatures at considerable depths are constant; but it seems not to have been ascertained whether the temperature of any spot—after other openings were extended beneath it—remained the same as it had been when it was the bottom of the mine. To invite enquiry on this subject, I venture to offer the following comparisons.

whose shafts and (levels) galleries, water and vapour circulate more freely than they had previously circulated through the cleavage-planes, joints, and crevices

East Wheal Crofty (a copper-mine, rocks:-	wrought	in felspati	hic and h	ornblendic	
rocks:—	18	38.	1840.		
	Depth, fms.	Temp.	Depth, fms.	Temp.	
Longclose, Engine-lode	85	63 [°] 5	85	60°	
. , , ,,	••	••	115	64.	
Trevenson, Reeve's Lode	115	69-	115	62.	
,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	••	••	135	70.75	
Wheal Vor (a tin-mine, opened in clay-slate).	18	38. I	18	59.a	
Main Lode, W.			210	74.5	
", "W	••		222	75·	
", ", B	230	78·			
", ", E	240	{80·5 81·	} 240	74.	
" ; W	••	••	251	80-	
", ,₩	••		h	686.	
", ", E	••		311	82.5	
,, E., bottom of the level	••	••	}	90.25	
, ,	••	••	321	91.	
Water discharged by pumps at the Adit, from	240	69.	321	75.	
Thus, at East Wheal Crofty,				•	
on the Longclose Engine-lode, the temper at the bottom was 63°.5 in 1838 v			. OF 6	J	
but it had fallen at the	anen fue ,	WOLKS WEL	e 50 ims	. aeep ;—	
ame spot to 60 ,, 1840		99	30 ,,	deeper;—	
yet at the bottom it had risen to 64 ,, ,,			115 "	deep:	
a These observations were made by the late Coments which had been used by the writer in 183	aptain Pran 8.	,, cis Francis,			

in the rocks and *lodes*; as therefore, each successive extension of deeper works intercepts, in its turn, the ascent of warm currents, the temperature of the original bottom gradually declines.

That, here and there, portions of various rocks and vein-stones are cooler than those above them, seems too well authenticated to admit of question. Such, infrequent interpositions, however, are seldom of great vertical range, and there is reason to believe they have usually but small horizontal extent; moreover, between

```
at Reeve's Lode, the temperature
at the bottom..... was 69.° in 1838 when the works were 115 fms, deep;-
but it had, at the same
      spot declined to .. 62.
                                                            20 .. deeper:-
yet at the bottom it had
         risen to ..... 70.75 ,, ,,
                                                           135 ,, deep.
  At Wheal Vor,
the temperature on the Main Lode
at the bottom..... was 80°.5 in 1838 when the mine was 240 fms. deep;—
but it had at the same spot
          fallen to .... 74 a , 1859
                                                           81 , deeper;-
nevertheless at the bottom
   it had advanced to .. 91. a ,,
    HENWOOD, Cornwall Geol. Trans., v. (1843), p. 395; Reports of the Royal
       Institution of Cormoall, XLI. (1859), pp. 21-3; Annales des Mines, 5me
       Série, XVI. pp. 571-3.
At Wheal Vor, on the 30th September 1858, the mid-day temperature
                                                at the surface was .... 67°;
but at the bottom, 311 fms. deep, the air and the water issuing from
                                             the rock were both at .... 80°.
           SMYTH (Annual Address), Quarterly Journal of the Geol. Society,
             xxiv. p. lxxxv. (Abridged.)
```

"On comparing the results obtained in Dolcoath in 1821—2 and 1827, it appears that the temperature was increased only 4° in one level with an increased depth of 42 fathoms, giving a ratio between the stations of 1° increase in 10.5 fathoms; and in another level the temperature was actually 2° to 2°.5 less than in 1822, although 42 fathoms deeper than the mine was then."

Fox, Report of the British Association, for 1857, Part 1. p. 100. a Ante, p. 764, Sub-note.

766 W. J. Henwood, on Subterranean Temperature.

their temperatures and the temperatures of the warmer rocks above and below them, the differences rarely exceed two degrees, and usually they are much smaller. Whether this state of things is a natural one,—or whether—in fact—it may have been brought about by the shafts, *levels*, and other openings in which it has been observed,—is beyond the scope of this enquiry.

W. J. HENWOOD.

3, Clarence Place, Penzance, 1870, February 3rd.

On the changes of temperature which take place—at the same, and at different, times,—on the surface and at depths of three, six, and nine feet in the Canga, at Agoa Quente in Brazil

The following observations were made with a view to ascertaining the rate at which solar heat penetrates the earth.

The high granitic ridge of the Caraça, situate in Long. 43° 10′ W., Lat 19° 50′ S., is, on the W.S.W.—separated by a deep and narrow glen from a parallel, but less elevated, range consisting of talco-micaceous slate, and schistose iron-glance interlaminated with quartz (Itabirite \$\ddot\$), in which—at least—one conformable bed of auriferous (Jacotinga \$\ddot\$) manganese, iron-glance, and talc has been extensively wrought. Considerable portions of the talco-micaceous slate, as well as of the Itabirite and Jacotinga, are overlaid by (Canga ||) breccia, containing sub-angular masses of the selfsame rocks and of quartz, usually cemented by compact brown iron-ore, but sometimes imbedded in

[•] Ante, pp. 174-6.

[†] Ibid, pp. 176, 220.

[‡] Ibid, pp. 214,-21,-44,-8,-98.

⁶ Ibid, pp. 173, 214,—16,—19,—23,—7,—36,—46,—51,—86, 303, 729.

[#] Ibid, pp. 216,—17,—36,—45,—99, 319,—24.

[&]quot;The ground rang under the hoof as if iron-plated; * * . The appearance of the mineral reminded me of the laterite in Malabar and Western India, but here it is the richest hæmatite."

BURTON, Explorations of the Highlands of the Brazil, 1. p. 815.

reddle. Particles of gold * and nests of native copper * occur in the Canga, but too rarely to need further remark.

Some four hundred fathoms N.W. of, and perhaps sixty fathoms above, the works at Agoa Quente,†—that is to say about three thousand six hundred feet above the sea—the surface is partially clothed with a stunted coppice of (Lychnophora) Candeia;‡ and at this spot holes §—of two inches in diameter and respectively of three, six, and nine feet in depth—were sunk in the Canga.

Thermometers—adjusted to the Standard of the British Association, by Pastorelli of London—were placed at the bottoms of the holes; which were then carefully closed with long wooden plugs wrapped in

^{*} Ante, p. 236.

⁺ Ibid, pp. 224-42, 729-31.

^{† &}quot;Sur plusieurs pentes couvertes de pierres, je trouvai en grande abondance une espèce à petites feuilles du genre Lychnophora Mart. (Vulg. candeis), genre qui, dans les montagnes, caractérise les côtes pierreuses."—Saint Hilairs, Voyage dans le district des Diamans et sur le litteral du Brésil, I. p. 81.

Gardner, Travels in the interior of Brazil, p. 473.

^{§ &}quot;Mes observations sont comprises entre le 11º degré de latitude boréale et le 5º dégre de latitude australe. • • • J'ai toujours observé dans un endroit abrité, un res-de chausée, une cabane d'Indien, un simple hangar. • • • Dans le village de Zupia, mon thermomètre était placé au rez-de-chausée, dans un trou de 8 pouces pratiqué dans le, sol; ce trou avait un demi-pouce de diamètre. Le maison était couverte de fuilles de palmier. • • • Lorsque le thermomètre était en expérience, on bouchait l'orifice du trou avec un morceau de carton sur lequel on appliquait une grosse pierre.

[&]quot;La température moyenne du village de Zupia avait été fixée à 21°.5 C. (70°.7 F.) par de nombreuses séries d'observations thermométriques faites en 1825, 1826 et 1829. Zupia ést élevé au-dessus de la mere de 1,225 mètres (4,019 feet).

[&]quot;Je rapporterai maintenant la march du thermomètre au-dessous du-sol, telle que je l'ai observée dans différentes localités.—

cloth; and-except for a minute or two at each read-

	Zupia.	
	8 pouces sous terre.	Dans l'air.
1830.		
Août le 8 à 9 h.m	21°4 C 70°5 F.	21.7 C 71° F.
10	21.4 70.5	22.2 72.
11	21.5 70.7	22.2 72.
1	21.5 70.7	23.8 74.8
3	21.5 70.7	22.8 73.
le 9 à 8 h. m	21.4 70.5	20 68.
midi,	21.4 70.5	23·3 74·
5	21.4 70.5	22.2 72.
le 10 à midi	21.4 70.5	23.8 74.
4	21.4 70.5	23.5 74.3
le 11 à midi	21.4 70.5	22.5 72.5
le 12 à 9.h. m	21.3 70.3	20.5 68.9
′ midi	21.3 70.3	21·1 70·
le 13 à 9 h. m	21.3 70.3	20.6 69.1
3	21.5 70.7	22.6 72.7
4	21.3 70.3	23.9 75.
le 15 à midi	21.3 70.3	22·8 73·
le 16 à midi	21.3 70.8	22·8 73·
3	21.8 70.8	22.3 72.1
le 18 à midi.	21.3 70.8	24.4 75.9
a boule du thermomètre a été p	placée à un pied au desso	us de la surface du s
le 18 à 3 h. soir	21°5 C 70°7 F.	23°4 C 74°1 F
4	21.5 70.7	22.3 72.1
6	21.5 70.7	21.7 71.
9	21.5 70.7	22.2 72.
le 19 à 9 h.m	21.5 70.7	21.1 70.
midi	21.5 70.7	21.7 71.
2	21.5 70.7	22.8 73.
3	21.6 70.9	22.2 72.
6	21.6 70.9	22.2 72.
le 20 à 11 h.m.	21.5 70.7	21.1 70.
midi	21.5 70.7	21.7 . 71.
3	21.5 70.7	22.2 . 72.
3		I
le 21 à 3 h.s.	21.6 70.9	
• •••••	1	į
le 21 à 3 h.s		

770 W. J. Henwood, on Temperatures at the

ing of the instruments—they were never reopened.

"Pendant les mois de septembre, octobre et novembre, le thermomètre a toujours indiqué 21°.5 C. (70°.7 F.)

Marmato.

"Le thermomètre a été placé à 1 pied dans la sol, dans une salle basse de la maison du surintendant des mines. La température moyenne de cette maison déduite d'une année d'observations est de 20°.5 °C. (68°.9 °F.). Elle est élevée au-dessus de l'Océan de 1,426 mètres (4,679 feet).

	•			
1830. Septembre le 9 à 11 h.:	m	20°5 C.	. 68 ⁹ F.	
	****************	l .	68.9	
3	****************		68-9	
le 10 à 8 h.:	m	20.3	68·5	
11	•••••	.,	68.5	
1	•••••••		68.7	
2	*****************		68.9	
. 3	•••••••	. 20.5	68.9	

Anserma Nuevo.

"Le thermomètre placé à 1 pied de profondeur dans le sol d'un rez-de-chausée.

		Thermomètre sous terres.
1830.		200 2 700 7
Decembre le 16 à 8 h.m.	******************	23.8 C 74.8 F.
le 19 8	•••••	23.7 74.6
le 22 9	••••••	23.7 74.6
le 22 11		23.7 74.6
9 h.s.		23.6 74.5
10	•• •• • • • • • • • • • • • • • • • • •	23.6 74.5

[&]quot;Pendant les mois de janvier et février 1831, le thermomètre a toujours indiqué de 23°·6 à 23°·7 C. (74°·5—74°·6 F.).

Puracé

"Dans la Troja del Cura, élevée de 2,651 mètres (8,698 feet) au-dessus de la mer, le thermomètre a été placé dans le sol à 1 pied de profondeur.

		1	Thermomètre sous terre.			
1831 Avril le 17 8	11 h.m.					
	midi.		18.1	55.6		
	2	•••••	13.1	55.6		
	4 h.s.	••••••	13.1	55.6		
le 18	8 h. m.	•••••	13.1	55·6		
	9	••••	13.1	55.6		

[&]quot;Des observations faites par Caldas, dans voisinage d'Anserma, donnent à cette partie de la vallée du Cauca,"—élevée de 1,050 mètres—(3,545 feet) " une température moyenne de 23°-8 C. (74°-8 F.).

The temperatures observed at 6 A.M., noon, and 6 P.M., from the 22nd of May to the 13th of July 1849, in each of three holes; * and at 3, 6, and 9

Quito.

[&]quot;Pendant mon séjour à Quito, j'engage ai M. Salaza à suivre la marche de son thermomètre mis à 1 pied au-dessous de la surface du sol. Les observations furent faites dans une salle Casse.

·	.	1			Therm	omėtre.			
Mois.	Dates	·à 7 1	h. m.	å 11	h, m.	1 48	h.s.	1 34	h. s.
1331. Septembre	26	15°6 €.	59°9 F.	1 <i>5</i> °5 C.	<i>5</i> 9.9 F .	1 <i>5</i> °5 €.	<i>5</i> 9∙9 F.	15°5℃.	59°9 ₽.
-	27	15.5	<i>5</i> 9·9	15.2	59-9	15.8	59.5	15.5	59-9
٠	28	15.3	59·5	15.5	<i>5</i> 9∙9	15.5	59.9	15.5	59-9
	29	15· 5 .	59-9	15.2	59.9	15.2	59 ·9	15.5	<i>5</i> 9·9
	30	15.5	59· 9	15.2	<i>5</i> 9·9	15.2	59.9	1 <i>5∙5</i>	59-9
Octobre	1	15.3	59·5	15·5	<i>5</i> 9·9	15.5	<i>5</i> 9·9	15.5	59 ·9
	2	15.2	59-9	15.5	59.9	15.5	59-9	15.5	56.9
	3	15.4	59.7	15.2	59 ·9	15.4	59.7	15.5	<i>5</i> 9·9
	4	15.5	<i>5</i> 9·9	15.5	59 ·9	15.2	<i>5</i> 9·9	15.5	59-9
	5	15.5	5 9·9	15.4	59.7	15.5	<i>5</i> 9·9	15.5	59.9
	6	15.5	<i>5</i> 9·9	15.5	59 ·9	15.5	59.9	15.5	59.9
	7	15.4	59.7	15.5	59-9	15.5	59-9	15.5	59·9·
	}	1		ł		Ι.		,	

[&]quot;Les observations que je viens de rapporter éstablissent, ce me semble, d'une manière certaine, que la température moyenne d'un lieu abrité situé entre les tropiques, est donnée par la température du sol prise à 1 pied de profondeur."

[&]quot;La température moyenne de Quito,—élevée de 2,914 mètres (9,660 feet) a été fixée par deux observateurs, MM. les colonels Hall et Salaza; leurs observations donnent une température moyenne de 15°.55 C. (60° F.).

Boussingault, Annales de Chimie et de Physique, LIII. pp. 228-35.

[&]quot;La temperature de Rio-Janeiro a été evaluée, par M. E. Chevalier (Voyage de la corvette la Bonite, p. 18), à 24°.2 C. (75°5 F.) d'apprès des observations faites à 1 pied de la surface du sol et à la profondeur de 3 mètres (9.8 feet), dans un puits."—D'ARCHIAO, Histoire des progrès de la Géologie, I. p. 88.

^{*}Trevandrum is situate in Long. 5° 7' 59" E., Lat. 8° 30' 32" N.; the Observatory hill, which exposes a grassy surface, rises to about 200 feet above the sea, is composed of the stone called *Laterite*, and in this thermometers were placed at the respective depths of 3, 6, and 12 French feet (3°2, 6°4, and 12°8 feet English measure).

772 W. J. Henwood, on Temperatures at the

A.M., noon, 3, 6, and 8 P.M., and midnight, from the 1st of January to the 13th of July, at the surface, are compared in *Table XXXVII*.; whilst the highest, lowest, and mean temperatures during intervals of ten days each, in the same periods as shown in *Pl. VI*. and in the following columns.

The following columns contain the monthly means of observations made daily (except on Sundays) at 6 A.M., noon, 6 P.M., and midnight, as well on each of these thermometers as on others at the surface, from the 1st of May 1842, to the 31st of December, 1845.

Months.	Surface.	8 French, 3°2 English, feet.	6 French, 6.4 English, feet.	12 French, 12°8 English feet.		
January	78 [°] 930	84 [.] 954	85°618	86°528		
February	80.386	86.888	86.625	85·78 4		
March	82.730	88.789	88-110	86.373		
April	83.370	89.614	88·527 a	86.916		
May	81.603	88-413	88-224 8	_		
June	79.023	85.012	86-883	86-878 5		
July	78-450	83-250	85.144	86.537		
August	78-990	83-566	84.736	85.894		
September .	79-978	84-575	85.133	85-633		
October	79.076	84.722	85-632	85.680		
November .	79.750	84-622	85.271	85.651		
December .	78-030	84-228	85.303	85-607		
Means	80.025	85.715	86-264	86.043		

The following conclusions are plainly discernible;-

The temperature of the ground at Trevandrum is from 5° to 6° higher than that of the air;—the principal maximum temperature of the air occurs about the beginning of April, and the extreme range is passed through in three months, the principal minimum occurring about the middle of July, the remaining fluctuations indicate a slight maximum about the middle of October The epochs of temperature are retarded with the depth below the surface, and, at the same time, the ranges diminish and casual fluctuations disappear.

CALDROOTT, Edin. Phil. Trans., XVI. pp. 379-93.

a For two years. b For one year only.

		Temperatures observed at											
the surface,						depths of							
	Periods.		_	ice,	th	ree fe	et.	1 8	ix fee	t.	nir	ne fe	et.
		Highest	Lowet	Keen.	Highest.	Lowest,	Kenn.	Highest.	Lowest.	Meen.	Highest	Lowest.	Mean.
Man	11 — 20 21 — 30 31 — Feb. 9. 5. 10 — 19 20 — Mar. 1. 7. 2 — 11 22 — 31 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 11 — 20 7. 12 — 30 7. 13 — 30 7. 14 — 30 7. 15 — 30	84·3 84·8 79·5 81·5 80·5 80·5 80·5 70· 71·8 72·7 69· 73·4 70·2 67·5	61· 62· 63· 66· 66· 64· 57· 62· 62· 44· 44· 44· 44· 44· 48· 50· 43·	72·3 73·7 73·5 70·4 69· 73· 71·5 68·9 70· 71·1 68·4 65·1 59·3 61·9 61·5 61·2 59·3 62·1 57·7 57·8	72·3 72·5 72· 71·9	71·4 71·7 71·1	71·9 72·1 71·5 71·5	71·6 71·8 71·2	7 0 ∙6 71∙2	71°3 71°4 71°1	71.8 71.6 71.2 71.1 70.8	71°1 71·3 71· 70·8 70·8	71.4 71.5 71.1 71.
1st to	Rxtremes {	42.8	_	67·3	73.7		 72·1	71.6	70·6	71·2	71.8	71· ·· 8	71-2
January 1st to	Ranges		~	66·3	73.7	71·1 ::_	:: 71·9	::	•••	••	71.8	70.7	71-1

At Brussels the temperatures hereafter mentioned are the means of ob-

774 W. J. Henwood, on Temperatures at the

From the 22nd of May to the 13th of July 1849, therefore, the extremes, the means, and the ranges of temperature were;—

servations at different depths on opposite sides of the Observatory; from 1834 to 1839.													
to 1839.	SOUTH OF THE OBSERVATORY.— UWSHADED.										OBSE ADED.	RVAT	et.—
			1	Oopths			1			Dep	the.		
Months.	Surface, Noon.	0-15 Metre (0-49 foot). 9 years.	0-40 Metre (1-31 foot). 1-25 year.	0.60 Métre (1.96 foot). 4 years.	0.80 Mètre (9.68 feet). 4 years.	1 Metre (8.28 feet). 2 years.	Burface. 6 years.	0·19 Metre (0·58 foot). 6 years.	0.45 Mètre (1.47 foot). 6 years.	0-75 Mêtre (2-46 feet). 6 years.	1Mêtre (3-28 feet). 5 years.	8-9 Metres (13-8 feet). 5 years.	7.8 Metres (25.6 feet). 5 years.
Jan	36·7	3 4 ·	34·2	3 7 ·1	38.2	38· <i>5</i>	37·	38.9	4 0∙6	42 ∙1	44·3	53·4	5 4 ·5 .
Feb	39-	34.6	34.3	36.8	37-8	39.	38.7	38·7	30-8	41.	43.7	51.6	54·
March	45.5	39.3	40-2	39.7	39-9	40.6	41.5	40-9	41.5	42.2	44.	<i>5</i> 0·2	53· 4
April	<i>5</i> 1·3	42.6	43-9	43-1	42-9	42·8	44.6	43-1	43.3	43-6	45·	49-6	52·8
May	65.7	53-4	54 ·6	51.3	50-2	49· 5	53.5	50· 4	50·	49-3	49.5	49.8	52·2
June	72-4	60.8	60-6	60-2	5 8·4	58-4	62·	58·3	57-4	56·	55.7	50-9	51.9
July	74.	68-1	66-3	64.6	63-2	64.8	64.8	60.8	60· <i>5</i>	59-7	59.5	53· 4	<i>5</i> 2·1
August	71.4	60.3	67.	68·4	62·3	64.9	63-1	60.3	60.5	60·5	60.9	<i>55</i> ·6	52·6
Sept	62·	<i>5</i> 7·8	58.6	59 ·	58-9	59-9	58-1	56· 4	57·5	58·	<i>5</i> 9·1	<i>5</i> 7·1	53·3
Oct	<i>55</i> ·1	52 ·6	53 ·6	54.3	54 ·9	55 ·4	51.3	51.9	<i>5</i> 3·6	54 ·8	56· 4	57-6	54 ·
Nov	44.	44.6	43.7	46.	47.2	46.8	42.7	44.5	46-4	48.4	50.7	57.	54·5
Dec	39-4	39-9	40.	42.	42.7	44-4	38-9	41.5	43·1	45.	47:3	55.4	55.
Means	54.7	48-6	49-8	49.8	49-7	50.4	49-7	48.8	49.5	<i>5</i> 0·	51.3	53 ·5	53·3

The following columns show the times of lowest, highest, and mean temperature;—at the surface and at various depths;—under both southern and northern aspects;—

Place of observation.	Extremes.	Means.	Ranges		
Surface	43· —73·4	59°1	30°4		
Three feet deep	71.1—73.7	72.1	2.6		
Six ,, ,,	70.6—71.6	71.2	1.		
Nine " "	71. —71.8	71.2	0.8		

	Southern aspect.— N Unshaded.				OTRHERN ASPECT.— SHADED.				
Depth.	Lowest.	Spring mean.	Highest.	Autumnal mean.	Lowest.	Spring mean.	Highest,	Autumnal mean.	
Surface	Jan. 13	Apr. 25	July 9	Oct.20	Jan. 17	May 3	July 18	Oct. 21	
Mêtres. Peet.	a		a			l			
0-15 0-49	,, 14	May 2	,, §	,, 80		ĺ			
0-19 0·58	١				,, 29	,, 9	,, 24	" 26	
0-40 1-31	Feb. 8		Aug. 2			1			
0.45 1.47					Feb. 5	,, 18	,, 30	Nov. 4	
0.60 1.96	Jan. 30	,, 9	July 23	Nov. 2	ŀ	1	l		
0-75 2-46	1	·			,, 17	,, 18	Aug. 6	,, 8	
0-80 2-62	,, 30	" 12	,, 25	,, 8		1			
1 3.28	l a	a	ĺ	a],, 27	,, 24	,, 9	,, 13	
3.9 12.8	1	l"		l " -	" -	I	1.	Jan. 10	

Between the times at which the lowest, highest, and mean, temperatures, respectively, occur at the surface and at different depths, the hereafter mentioned periods, therefore, intervene:—

Surface	1	1	1 1	1 1	1		
Mètres. Feet.	ا		ا م		- 1		
0-15 0-49	1 day 7 d	ays	10 days				
0·19 0·58	, .	,		12 days	6 days	6 days	5 days
0-40 1-31	25 days .	. 24 days	1 :				
0-45 1-47	1 1			7 ,,	4 "	6 "	9 ,,
0.60 1.96	.		3 ,,				
0-75 2-46	.		٠	12.,,	5 ,,	в "	4 ,,
0.80 2.62	10	,,				İ	
1 3.28	7	"a 4 "	2 ,,4	10 "	6 "	8 "	5 ,,
3-9 12-8	.	• ,.		<i>5</i> 2	53 "	66 ,,	58 ,,

QUETELET, Memoires de l'Académie Royale de Bruxelles, X. pp. 3-80, XIII. pp. 3-52. (Abridged.)

776 W. J. Henwood, on Temperatures at the

From the 22nd of May to the 13th of July, however, the extremes, means, and ranges were;—

At Greenwich observations have been made daily, for many years, on thermometers—at the surface and at depths of 1 inch, 3.2, 6.4, 12.8, and 25.6 English (3., 6., 12., and 24. French) feet. "The soil [is composed] of beds of sand; of flint-gravel with a large proportion of sand; and of flints with a small proportion of sand, cemented almost to the consistency of pudding-stone. * * * [Those parts of the tubes which project above the surface] are protected by a wooden case or box fixed to the ground; the sides of the box are perforated with numerous holes, and it has a double roof. In the North face of this box is a large plate of glass through which the thermometers are read." The extremes and means observed during the years 1865, 1866, and 1867 have been;—

•	Jan.	Peb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
			A	t the	surf	ice.						
Highest	53.6	55·2	67·8	75·8	81· 6	88.9	84·5	8 4 ·7	82·7	73·2	63·	o 54∙8
Lowest	20-	30.3				54.6				ı	37-9	28.7
Mean	40.5	44.1	43-	l57·8	61.9	69-1	69-8	68-2	67.	57-5	47.7	43.1
	·	•	At a	depti	s of a	me in	ch.		•	•		_
Highest	50.2	50.7	50.1	59-4	65.	69.5	77.1	70.6	70.5	63-2	56.8	51.9
Lowest	. 29.	31.2	35·1	43.8	45.2	57.5	56.8	55.7	52·1	46.6	39-	3 4 ·3
Mean	40.2	42.7	40.7	51.2	55.5	63.	64.6	62.8	61.7	54.2	46.6	43.3
•						3·2 fe		-			•	
Highest	. 46·	46.	45.3	51.8	57.2	61.6	63.8	63.7	64.4	61.5	53.4	 48 •9
Lowest	. 39-1	39-4	39-1	40.5	48.6	53.6	59.4	59.	57.	52-2	45.6	42-2
Mean	. 42.6	42.8	41.4	47.	52-1	58-1	61.5	61.1	60.8	55-9	49.7	45.8
				-	•	6·4 <i>fe</i>						•
Highest												
		45· 8										
Mean	. 46.7	46.78	45.40	47.2	150.7	54.8	58.4	59.3	59.7	56.5	53.3	49.7
				_	•	ر 12·8						
Highest			1									
Lowest	47.6	45.8	1								53-1	
Mean	. 49.7	47.9						55-1	56.1	56-1	54.5	52 ·2
				-	•	25·6 j		_		_		
Highest	1		4	1	1	1	i .			,	52.9	1
	. 51.	50.2										52.3
Mean	. 52	51.2					50.1	50.8	51.5	52.2	52.7	52.6
				a Two	years	only.						

Place of observation.	Extremes.	Means.	Ranges.
Surface	43· —73·4	60°·1	30°·4
Three feet deep	71·1—73·7	71.9	2.6
Nine " "	70·7—71·8	71:1	1.1

The intervals between the times of highest and of lowest temperatures at the surface and at different depths, are shown in the foregoing columns.

The highest, lowest, and mean temperatures, as well as the ranges at the surface and at various depths, were—

	Highest,	Lowest.	Means.	Ranges,
Surface	84 [.] 7	20°	<i>55</i> °.8	64°.7
l inch in depth .	77-1	29.	<i>5</i> 2∙2	48.1
3-2 feet "	64.4	39-1	51.6	25.3
6.4 ,, ,,	61.4	44.2	52-4	17:2
12.8 ,, ,,	57·2	44.8	51.4	12.4
25-6 ,, ,,	52·9	48.	50-9	4.9
1			l	1

Magnetical and Meteorological Observations at the Royal Observatory, Greenwich, 1865, pp. XLII.—III., CCLXXXIII.—VII.; 1866, pp. XLIII. —IV., CXCII.—VI.; 1867, pp. XLIV.—V., CCLXIII.—VII.

At and near Edinburgh observations were continued from 1837 to 1842 at depths of 3.2, 6.4, 12.8, and 25.6 (English) feet;—

on the Calton Hill, in porphyritic trap, at 350 feet above the sea; in the Experimental Garden, ,, sand ,,, 70 ,, ,, ; & at Craigleith, ,,, sandstone ,,,, 150 ,, ,, ; with the undermentioned results;—

	Calton Hill.				Experimental Garden.				Craigleith.			
Depth.	Highest.	Lowest	Means.	Ranges.	Highest.	Lowest.	Means.	Ranges.	Highest.	Lowest	Means.	Ranges.
3-2 feet	56 · 2	3 <i>5</i> .7	45·5	20°5	<i>5</i> 7∙2	35°.1	46°1	22°·1	58.9	35°4	45°9	20°5
6.4 ,,	52·3	39.7	45.8	12.6	54· 6	38 ·6	46-4	16.	<i>5</i> 3·8	38·1	45.9	15.7
12-8 "	49-4	43.6	46.3	5.8	<i>5</i> 0·6	42·8	46.7	7.8	51-1	40.7	45.9	10.4
25.6 ,,	47·8	46.	46 ·8	1.8	48-2	46·	47.1	2.	48 ·5	43.8	46.	4.7

FORBES, Edin. Phil. Trans., xvI. pp. 194, 204, -7; Proceedings of the Royal Society of Edinburgh, 1. pp. 223, 344*.

778 W. J. HENWOOD, on Temperatures at the

The depths at which the observations were made;—
the mean temperature of each spot at the commencement of the series (on the 22nd of May, 1849);—and
the nearest periods at which these were, respectively,
the means of the climate; are shown in the following
columns:—

Depth.	1849, 22nd May. Mean temperature,			Interval,	
	underground.	Date.	Mean temp., surface.	days.	
Three feet	7å·3	Mar. 3	74°3	80.	
Six ,, *	71.4*	Apr. 6	72.7	46•	
Nine "	71.6	Apr. 6	72.7	46.	

At Upsal thermometers sunk 1.07 foot, 2.14 and 3.20 feet in the ground, and observed daily at six in the morning, two in the afternoon, and nine at night, showed temperatures of which the monthly means are set forth in the following columns:—

	Times.	1.07 foot.	2·14 feet.	8:20 feet.
1833.	July	60°54	<i>5</i> 9.	56.96
	Aug.	<i>55</i> ·61	55.45	55·18
	Sept.	53.92	53.61	53.47
	Oct.	48.14	48.34	49.26
	Nov.	39.	40.31	42.20
- .	Dec.	33.45	35.18	37.
1834.	Jan.	29.28	81.24	32.72
	Feb.	31.31	31.96	32.43
	March	32.63	33.13	83 44
	April	38.04	37.43	36.93
	May	48.02	46.56	45.10
	June	<i>5</i> 6·67	54.50	52.32
Mean	3	43.88	43.89	43.92

RUDBBRG, Ann. der Chem. und Physik de Poggendorff, XXXIII. Memoires de l'Académie Royale de Bruxelles, X. p. 36. Edin. New Phil. Journal, XXIII., p. 345

^{*} At Trevandrum the mean annual temperature was higher at six, than at either three or twelve, feet.—Caldrott, Edin. Phil. Trans., XVI. p. 392.

The proportion of solar heat absorbed by the ground must, of course, depend, in some measure, on the nature of the surface.*

The following columns set forth the means of observations, at the surface, as well as at depths of three,

"Temperature of the air and of different kinds of soil, Alverton, near Truro.

	Air.	Clay.	Siliceous Sand.	Peat.	Garden Loam.	Grass.
Date.	Mean temp.	Mean temp.	Mean temp.	Mean temp.	Mean temp.	Mean temp.
1852. Sept	57°7	56°6 .	60°	57 [°] 6	59°6	60°1
Oct Nov	51·1 50·9	50·1 46·1	50· 49·1	51·6 50·2	51·1 50·2	54·2 52·2
Dec	49·3	46.7	46.2	47.8	48·1	49.6
1853. Jan Feb	44·9 37·	42·1 36·	42·1 34·6	43·7 37·4	43·2 36·8	46·7 40·8
March	44-4	41.1	40.4	42.2	41.3	45.2
April	50.5	49.9	<i>5</i> 0·	50.8	50.8	53∙
May June	53·4 <i>a</i> 57·6	64·	57·2 63·9	57·7 63·5	58·3 65·1	60·1 67·3
Jul y	60-5	63·1	63.	68.8	64.3	67.2
Aug	60-5	62.	62.5	62.8	63.9	68·1
Annual Means	51·5	51.2	51.6	52·4	52.7	55-4

WHITLEY, Bath and West of England Agricultural Journal, III. pp. 12-15. (Abridged.)

[&]quot;The observations from which the following table has been deduced were made at Alverton, near Truro, in 1852—3. * * * Four pits, about two feet deep and two feet wide, were dug in good healthy garden loam. [The first] was filled with yellow clay frem the clay-slate; [the second] with pure white sand, from the sand-bed at St. Agnes Beacon; [the third] with peat, almost pure vegetable matter, well worked before put into the pit; [the fourth] with garden loam. The bulb of the thermometers was placed 4 inches below the surface in the centre of each pit, and another thermometer was placed in the same manner under the short grass of the lawn. Each variety of soil was thus subject to the same drainage below and to the same influences above. The readings of the thermometers were made in the morning when the temperature of the soil was lowest, and again in the evening when it was highest * * *

a 1852.

780 On Temperatures at small depths in Brazil.

six, and nine feet; at 6 A.M., noon, and 6 P.M., during the same period; *---

i		6 A	M.	1	1	No	on.	1	ј 6 р.м.			1	
Place of observation.	May.	June.	July.	Means.	May.	June.	July.	Means.	May.	June.	July.	Meens,	General moans.
Surface	57 [°] 8	56 [°] 8	51°9	55°5	56°∙5	60°3	64 [°] .7	62 [°] 5	63°	63 [°] 2	61°5	62 [°] 6	60°∙2
3 feet deep	72.7	72.7	71.5	72.3	72.7	71.9	71.4	72.0	72.6	71.8	71.4	72.	72-1
6 " "	71. 2	71.2	-	71.2	71-2	71.1	—	71.2	71.2	71.	_	71.1	71.2
9 " "	71.3	71.4	70.8	71.2	71.4	71.2	70.8	71.1	71.4	71.1	70.7	71.1	71-1
Means, at 3, 6, } and 9 feet deep }	71.7	71.7	71.1	71.5	71.7	71.4	71.1	71.4	71.7	71.3	71.	71.4	71.5

It would seem, therefore, that at depths of three, six, and nine feet respectively, the mean temperatures slowly declined, both from May to July, and from morning to evening.† These observations, however, extend to neither of the annual extremes.

W. J. HENWOOD.

3, CLARENCE PLACE, PENZANCE. 1870, March 80th.

[†] At Trevandrum observations—at intervals of six hours—from the 1st of May 1842 to the 31st of December 1845, afforded the undermentioned results.

Depths,	Hours of observation.					
Dopius,	6 A.M.	Noon.	6 P.M.	Midnight.	Means.	
3 feet	82 [°] -50	82 [°] 66	82°55	82 [°] .50	82 [.] 55	
6 ,,	83.79	83-98	83.88	83-82	83.87	
12 "	83-90	83-99	83.95	83.90	83-93	
Means	.88-46	88.54	83.46	83:41	83-45	

CALDECOTT, Bulletins de l'Académie Royale de Bruxelles, IX. Partie I. pp. 303
—10; Proceedings of the Royal Society of Edinburgh, I. pp. 432—3;
Edinburgh Phil. Trans., XVI. p. 391. (Abridged.)

[•] Such observations only as were made simultaneously at the several stations, have been used in deducing these averages.

EXPLANATION OF THE PLATES.

Copies of Working-plans and Sections of mines are marked with asterisks (*).

In the Plans lodes are represented by single, and eross-veins by double, lines.

In Longitudinal Sections the darkest shades indicate the portions which have been removed

CHILI.

Plate I.

Plan of the Mining District of Chafarcillo.

Plate II.

Longitudinal section of the Colorada lode, Chanarcillo. †

BRAZIL.

Plate III.

Bird's-eye view of the gold-formation at different depths in *Morro Velho*.*

Plate IV.

Fig. 1*. Plan of Gongo Soco.

" 2*. Longitudinal section of the Gongo gold-formation.

,, 8⁴. ,, ... ,, ... Cumba

4. Transverse section of the strate.1

CORNWALL.

Plate V.

Fig. 1*. Plan of West Caradon.

" 2*. " South Caradon.

,, 3*. ,, Wheal Trelawny.

,, 4*. " Wheal Mary Ann.

[†] Drawn from survey by Edwin Price Waring, Esq., Superintendent of Colorada.

t ,, Captain John Luke of Gongo Soco.

Plate VI.

Projections of temperatures observed, at the surface and at depths of three, six, and nine feet, within the tropics.

WOODCUTS.

		NORT	H-WESTERN INDIA.
Fig	. 1.	The Danda mine;	Section. Beds of metalliferous quartz; in talcose and chloritic slates.
"	2.	" Dhunpoore ";	,, . Jointed structure common to limestone and slate.s
"	3.	" " " " " "	View of the joints, and of the (bunches) masses of copper-ore which occur at their intersection.
"	4.	""";1	Plan ,, ,, ,,
"	5.	District of Agur;	" beds of iron-ore, in quartzose-talc, and in clay-slates.
			CHILI.
,,	6.	" Chañarcillo ; l	Plan. Jointed structure of the first lime- stone.
"	7.	" " ; ^r	Transverse Section. Colorada, Waring, Descubridora, and Candelaria lodes; Manto de Ossa; first limestone.
"	· 8.	"";]	Plan. Dykes of felspathic and horn- blendic rocks, intersected by the Colorada and Candelaria lodes.
,,	9.	" " 5	" . Meridional positions of (bunches) masses of ore in different lodes.
"	10.	Quebrada Seca. T	ransverse Section of the hornblendic, quartz, and felspathic rocks, and of the metalliferous beds and veins.
,,	11.	San Josè; Plan. V	Veins of metalliferous quartz.

BRAZIL.

Quartz-rocks and quartzose slates.

- " 12. Catta Branca. Plan (Sketch). Auriferous quartz, in micaceous (and talcose?) slates
- " 13. Catta Preta; Transverse Section. Beds and floors of auriferous quartz in quartz-rocks.

Clay-slate.

- Fig. 14. Gongo Soco (Camara); Plan (Sketch). Isolated (bunches)
 masses of auriferous quartz, in variously
 coloured clay-slates.
- " 15. Morro Velho; Transverse Section (Sketch). Formation of auriferous iron-pyrites and quartz, in clay-slate.

Jacotinga.

- ,, 16. Pitangui; Transverse Section (Sketch). Bands of auriferous Jacotinga.
- " 17. Agoa Quente; Transverse Section (Sketch). Alternating bands of iron-ore and quartz (Itabirite), displaced (heaved) by minute veins of quartz.
- ,, 18. ,, Horizontal Section (Sketch). Schistose rocks of iron-glance and quartz (Itabirite) interfoliated with auriferous Jacotinga.
- " 19. Catta Preta; Transverse Section. Isolated masses of auriferous quartz imbedded in iron-glance, manganese, and talc.
- " 20. Cocaës (Alto da Cruz); Longitudinal Section. Layers of talc in iron-glance and quartz.
- " 21. Gongo Soco; Horizontal Section. A conformable layer of rich auriferous Jacotinga interlying a (horse) mass of iron-glance and quartz (Itabirite), surrounded by Jacotinga thinly sprinkled with particles of gold.
- " 22. " View of flexures in schistose iron-glance interfoliated with quartz (Itabirite).
- ,, 23. ,, ; Transverse Section. Contorted layers of tale and auriferous Jacotinga.
- " 25. " ; Longitudinal Section. Alternating layers of iron-glance and slightly auriferous Jacotinga displaced (thrown) by thin crossveins of quartz.

UNITED STATES.

" 26. North American Mine. Massive and cellular quartz sprinkled with native copper.

- Fig. 27. South Cliff Mine. Minute grains of native copper and trap in a vein of prehnite traversing trap-rock.
- ,, 28. Douglas Houghton (Henwood) Mine. Longitudinal Section of the Works.
- " 29. " . Grooved surface of vein-stone and native copper.

CHANNEL ISLANDS.

- ,, 30. Sark's Hope Mine. Longitudinal Section of the works.
- "81. Knockmahon Mine; Stage lode. Longitudinal Section of the works.

CORNWALL.

" 32. Wheal Trelawny. Section of a concretion imbedded in clay-slate.


INDEX.

Agoa Quente, subterranean temperature at, 729-31,-56.
, climate of, 729; Table XXXI.
, rain at, 730.
, temperatures at depths of three, six, and nine, feet, at,
768-80; Table XXXVII.
, growth of fish in warm water at, 355, 731.
Agordo, treatment of inferior copper-ores at, 579.
Agua, Manto de, a calcareous bed impervious to water, at Chafarcillo, 77.
Agur, the rocks, iron-ores, and smelting-works of, 21-2, 57.
Agur, Al, the rocks and copper-ores of, 11.
Air, issue of from joints in the Itabirite, at Gongo Soco, 250.
-, chalk at Belvedere, 250.
Airy, G. B., Esq., subsidence of rocks, and veins at Dolcoath, 666.
Akysa, sandstones and iron-ores of, 67.
Alabama, auriferous rocks of, 371.
Alderley Edge, ores of copper in sandstone at, 77.
Ale and Cakes, early use of plunger-pole at, 570.
Alfred, Wheal, silver ores obtained at, 112.
, ancient mode of operation, long continued at, 146.
, profits realized at, 445; Table XIV.
Alfred Consolidated Mines, profits realized at, 445; Table XIV.
Algæ, growth of in water impregnated with mineral salts, 585,—8
Alger, - Esq. and Dr. C. T. Jackson, copper-ores of the Nova-
Scotian coal-measures, 507.
Algonquin mine, rocks and ores of, 479-80.
Alison, R. E., Esq., qualities of Chilian copper-ores, 166-7.
Allen, John, Esq., altitude of Liskeard and its neighbourhood, 748.
, damage to fertile land by water from mines, 353.
, increased value of land consequent on mining operations
in it, 694.
, mining in the Caradon district, 694.
Alloy, nature and proportion of in Brazilian gold, 175-6,-9-80,
205-6,-36-7,-86,-99,311,-
334,-7,-8,-40; Tables VII.,
IX., X.
Canadian gold, 384.

Alloy, nature and proportion of in New	Granadian gold, 360.
Sibo	
, Vir	
, We	lsh gold, 642.
Almora, composition and structure of	
, frequency of earthquake-shoo	•
Altitudes, temperatures of mines at var XXXV., XXXVI.	•
Aluknunda, auriferous sands of the, 3,	46—7.
Amalgamation, 97, 877; Tables VII.,	
America, North, auriferous rocks of, 3	
•	68—870.
Amygdaloidal rocks, south of Lake Su	
Analyses, of mineral-waters, 573,—86	-
, " rocks, 75,—8, 80, 240,—	
and assays of different meta	
197-8, 204-6, -37, -86, 334-5, -8, -5	
523,-86,-45-8,-50,-66,-8,-71,-81-	
Tables VII., XXII.	, , , , , , , , , , , , , , , , , , , ,
Andrada, Senr. Gomes Freire de, his s	ttempts to preserve the forest
of Brazil, 348.	• •
Angles, included between lodes and cr	oss-veins, 127-9, 225, 527,-
_	558-60,-98, 608,-22,-88-85
	17; Table III.
, str	•
Ann, Wheal, silver and silver-ore of, 1	
Anna, Santa, auriferous deposit of, 21	
Ansermo Nuevo, temperature of, 770.	
Ansted, Professor, on the auriferous se	eries of Virginia, 379—81.
, ' detrital gold	_
, proportion	of gold extracted at the
	am mines, 377—9.
, geology of	•
	Herm, 736.
, climate of (Juernsey, 735.

- Ansted, Professor, on the wells of Herm, 737. Antimony, ores of, 118-19,-21,-79, 334,-74, 526,-35-6,-45,-8, 566, 617, 707. Antiquities, discovered in copper-mines near Lake Superior, 415-19. --, imbedded in detrital matter, 344-5. Antonia Pereira, disappearance of rivulet beneath rubbish, in the dry season, 353, -, ores, metals, mines, and smelting-works at, 213, 303,-59. rocks of, 303. - rubbish from mines carried into river Gualaxo during the rains, 303. Ants, damages done by, 850-1. -, discovery of gold in a nest of, 248. Apire (labourer), the weight of ore extracted by, in Chili, 147. Apjohn, Professor, analyses of lead, zinc, iron, and auriferous ores b_{y} , 545—50. Aplin, P. H., Esq., proportion of tin-ore in the vein-stone of the Saint Ives Consolidated Mines, 472. Arabia, sounds emitted by moving sand in, 154. Archiac, Viscount D', on the temperature of coal-mines in Virginia, 755. Ardtully Mine, rocks and metalliferous deposits of, 615-19, 742; Table XXI. -, subterranean temperature of, 744. Argall, Mr. William, on the proportion of tin-ore in the vein-stone of Wheal Vor, 472. -, profits and losses at , 446. Arrastres, quantities of ore-ground, and of gold extracted, by, 203; Table VII. Arsenic, compounds of, 118,-94, 213, 834,-74, 526, 617,-79. Ashburner, W., Esq., on the produce and quality of gold, the cost of extracting it, and the loss of mercury during its extraction, in
- Ashley, T. D., Esq., on subterranean temperature in the *Duckinfield* Colliery, 754—5.

California, Table XXII.



Ballymurtagh, precipitation of copper from mine-water at, 573-88.
Bamboo, growth of in water occasionally frozen, 45.
Banat, proportion of silver in auriferous iron-pyrites, of the, 100.
Banka (Banca), the rocks, tin-ore, and gold of, 631.
Bankart, Frederick, Esq., on the copper-ores of, Eardiston, 515. ——————————————————————————————————
590.
Bankart, Howard, Esq., on the quality of precipitate from water of
the Gwennap adit, 587.
Baranquilla, on recent sandstone at, 155.
Barley, cost of at Chafarcillo, Table V.
Barnett, Capt., R.N., on Magnetic declination in Jamaica, 511. Baroolee, the iron-ores of, 33.
Barrel-work, proportion and quality of, in the mines near Lake
Superior, 434.
Barreteros (miners), wages of, in Chili, 151; Table V.
Barrow, comparative cost of work by it, and by tram-way, underground, 144.
Barrul Cajoor, iron-ores and sandstones of, 67.
Barytes, the sulphate of, its modes of occurrence, 513,-46, 707.
Basset, family of, their heir-looms of plate manufactured from the
silver of Dolcoath, 113.
Basset & Grylls, proportion of tin-ore in the veinstone of, 472.
Basset, West Wheal, produce and profits of, 449-50; Table XIV.
Basset, Wheal, silver-ores of, 113.
Batēa, use of in Brazil; Table IX.
, when no longer serviceable, burnt for sake of the gold im-
bedded in fibres of the wood, 641.
Bathurst, on the rocks near, 490-500.
, lignite of, 510.
Batten, J. Hallet, Esq., on the cultivation of the Bhabur (Turace), 40.
, — disappearance of mountain-streams,
beneath gravel in the low-lands, 44.
, his extensive works for irrigating the Upper Bhabur, 43.
on the iron mines and miners of Agua 91

INDEX.

Batten, J. Hallet, Esq., on the iron-mines and miners of Dhunneea
Kote, 34.
Munglalekh,
26.
, on the ravages of wild beasts in the Hima-
laya, 17.
Bauerman, Hilary, Esq., on the drift of Lake Superior, 485.
Bayfield, Capt., R.N., on the conglomerates of Lake Superior, 403.
,sandstones of ,, , 392.
, magnetic declination in New Brunswick,
492.
Beans, prices of, in Chasarcillo, Table V.
Bearhaven mines, cross-course of, 608.
, lodes of the, 603—8, 741; Table XX.
, one of the, inclines at a higher angle
than the cleavage-planes of the rocks, 564, 604.
, rocks of the, 602—3, 740.
, the workmen of sometimes cook their food under-
ground, 610.
, water, quantities of, at different seasons, 609.
, a steam-engine and forges underground, 610.
, subterranean temperature, 742.
Besumont, de, M. L. Elie, on the structure of gneiss, 519.
Beche, Sir H. T. De la, on the rocks intersected by the Agua Alta,
511.
, —————————, elevation of the Caradon district, 696.
, —— granite of Brownwilly, 662.
, —— hornblendic rocks of Saint Cleer and
Menheniot, 672.
, — joints in the granitic and schistose
rocks of Cornwall, 673.
,, rocks N. of Hessenford, 701.
,
646.
, —— silver-ore of Herland, 111.
, — the alleged waste of copper near
Tavistock, 693.
Reckett. Mr. Denuty Collector on iron-smalting in the Himeleye 57


```
Beds, of pebbles and gravel in the Himalaya, 45.
 ---, ,, shingle, gravel, and sand in Chafarcillo, 142.
   -, ,, shingle, gravel, sand, and shells near Caldera and Copiapo,
           139,—57.
 ----, metalliferous, 4, 5, 7-16, 18-33,-5-41, 67, 82, 161,-79-83,
                      87-207,-14-16,-19-32, 237-41,-2-6,-8-86,
                      298-804,-11-40,-72,-5,-7,-80, 496, 509,
                      513, -43 - 69, -74 - 8, 616, -20, -8, 726, -8, -9,
                      732,-3,-8,-44.
Beef, price of at Chafarcillo, Table V.
Beeralston, proportions of silver in the lead-ores of, 109-10.
Bejapoore, the ferruginous conglomerate of, 39.
Bella Fama, on the employment of fire-flies to light the miners at
                209.
Belden, Dr. J. H., on the use of fire-flies to give light to miners at
                     their work in Brazil, 209.
Bellar, the copper-ores of, 4.
Belt, Thomas, Esq., on the proportions of gold at various depths in
  the quartz formations of Australia, 179.
Belvedere, issue of air from joints in the chalk at, 250.
Bengal, on certain iron-ores and sandstones in, 64-8.
Bento Rodriguez, on the (Cascalho) detritus of, 343.
Beresite-rock, description of, 177.
Berezovsk, on the auriferous deposits of, 177.
         -, quality of the gold obtained from, 335.
Berger, Dr., on the stanniferous granite of Devon, 175.
         -, - silver-ores of Cornwall, 110,-11.
Berggieshübel, on the ores of iron, zinc, and lead at, 546.
Berthier, M., his analysis of argentiferous lead-ores, 102,—3.
Betsy, Wheal, proportion of silver in the lead-ore of, 110.
Bhabur, bodies of detritus at the mouth of every glen in the, 42.
, unwholesome climate of the, during the warm season, 40.
      —, irrigation of the, 43.
      -, iron-ores and furnaces of the, 40.
  -----, rivers disappear beneath detritus on entering, and reappear
  on leaving the, 43-4.
Bhamouree, the limestones and sandstones of, 38.
Birch Torr, proportion of tin-ore in the vein-stone of, 472.
Bischoff, Professor, on the temperatures of mines, 757.
Bismuth, compounds of, 111,-79, 334,-5,-6,-7,-74, 546, 640.
Black workpeople employed in the mines of Brazil, and the United
  States, 289-96, 382-3; Tables VI.ª VII.
```


Blamey, Capt. Thomas, on the auriferous deposit of	f Agoa Quente, 283.
,	- Gongo Soco,
250,-4,-7,-60,-3,-6,-8,-9	
, on the manufacture of iron in	Brazil, 262.
precipitation of copper 589.	•
, ——assistance afforded to Table VIII.	the writer by,
Blamey, John, Esq., notice of diamonds discovered at São Miguel, 242.	the Corrego de
Blas San, produce of, 124; Tables IV. XIV.	
Blast, means of obtaining, 54, 219-20, 528.	
Blasting, extent of at Morro Velho, Table VII.	
dexterity of the Cornish miner in, 151-2.	
, quantities of gunpowder used under various	circumstances
in, 201.	
, of ore extracted by, at Morro Ve	lho, Table VII.
, in other min	es, 200.
, at different d	
Velho, 199	
Blavier, M., on the argentiferous lead-ores of Aveyro	
Blende, occurrence of, 86, 112, 545-6, 621,-40, 707	
Blight, Mr. J. T., on the occurrence of stream tin-ore district, 695.	
Boase, Dr. H. S., on concretionary structure, 702.	
, ,, the directions of valleys, 718.	
, ,, relations between the structure lodes, 187.	of rocks and
, ,, curvatures in beds of schistose r	ocks, 259.
, " granite of Caradon, 662,-72.	
, ,, hornblendic rocks of Saint Cleer: 662,-72, 702.	and Menheniot,
, ,, junctions of granite and slate in	Cornwall, 667.
Boa Vista (near Ouro Preto), on the topaz formation	
Bocona, extent and produce of works at, 124; Table	es IV. XIV.
Bohemia, proportion of silver in lead-ore of, 101.	
Boiling Well, proportions of silver in lead-ores of, 12	
Bolaco lode, extent and produce of works on, 124; T	ables IV. XIV.
Bolivis, quality of detrital gold in, 360.	


Bond, Thomas, Esq., scenery of the Caradon district, 697. Bonina, Padre, water-wheels first erected by, in the gold-districts of Brazil, 361. Bonnard, M., on the proportion of silver in the lead-ore of Bohemia, Bonegarh, on the iron-ore of, Table II. Borer, the, differently used in different districts, 151-2. Borgnet, Professor, on the influence of forests on the sources of the Meuse, 138. Borlase, Dr. William, on the native-silver of Saint Just, 110. -----, ----- argentiferous lead-ores of Cornwall, 118,-19. tin-mines of Polberrou and Polgooth, 453,-4. ---, ---- detrital tin-ore of Cornwall, 486. ------, on early copper-mining in Cornwall, 687-8, 691,-2. , on the Wringcheese & Kell-mar'r rocks, 696. ·Borriga, on the copper-ore's and sandstones of, 514. Borringdon Consolidated Mines, proportion of silver in the lead-ore Borrow, Henry, Esq., on profits obtained at Levant, 442; Table XIV. Boscean, proportion of metal in the tin-ore of, 472. Botallack, extent of submarine works at, 599. _____, fresh-water below the sea-level at, 539. ____, eels in the mine-water at, 717. ____, on the granite and slate of, 659. ____, profits made at, 442-3; Table XIV. _____, proportion of tin-ore in the vein-stone of, 472. Bournon, M. de, on the proportions of silver in the ores of Chalanches, 103. Boussingault, M., his analysis of mine and detrital gold, 360. _____, on the mines of Marmato, 137. _____, ___ diminished flow of water from springs when ground is stripped of wood, 138. ————, mean temperatures within the tropics, 768-71. Brady, Fort, annual fall of rain at, 477. Branches, enrich the lodes with which they unite, 83, 91, 607; Table III. Bray, Capt. Walter, on the auriferous deposits of Gongo Soco, 255

257,-60,-6,-71,-2,-3,-4.

	-, the forests of, 170, 296.
	, destruction of, occasioned a scarcity of water, 344.
	, laws for the preservation of, 296, 343-4.
	, manufacture of charcoal in, 219, 226-32.
	-, pastures of, 168-9, 296.
	-, rocks of, 171-3,-6-8,-80,-6, 209-12,-16,
	221,-42-4,-7-50,-97-307.
	-, discovery of gold in, 360.
	-, gold-mines of, 175,-7-84,-7-207,-9-16,
	221-41,-5-7,-9-88,-98-304.
	-,, people employed by British
	Mining Companies in the, Tables VII. IX. —, detrital gold of, 342-59.
	•
	-, proportions of gold obtained from different
	formations, 369.
	-, alloys of gold ,, ,, , 206, 286, 333-40.
	-, iron smelting-works of, 219,-42,-7,-61.
	-, mining laws of, 361-5.
	-, water-wheels first used in the mines of, 361.
	-, machinery of British manufacture now in use at the mines of, 288.
•	-, British capital invested in the mines of,
	246,–89, Tables IX. XIV.
	-, returns and profits made by British Mining
	Companies in, 208,-89; Tables VII, IX. XIV.
	-, quantities of gold obtained in, 366-9.
	—, duties paid by miners to the government in,
	208,-89, 361-7; Tables VII. IX. XIV.
	-, supercession of mining by agriculture in, 370.
	- Slavery at Gongo Soco, 289-96.
	 Slavery at Gongo Soco, 289–96. Slaves of native masters, permitted to search
	-, Slaves of native masters, permitted to search
	—, Slaves of native masters, permitted to search for gold on their own account
	-, Slaves of native masters, permitted to search










Caradon, the mining district of, 655—99.
,, granitic, slaty, elvan, and hornblendic rocks of, 655—
664,-6-74.
, " stanniferous granite of, 664—6.
, ,, directions, dips, widths, and compositions of lodes in,
674—81.
, ,, tin, copper, and iron ores of, 676-80,-95.
, ,, directions, dips, widths, and compositions of the cross-
courses in, 681—5.
, " intersections and (heaves) displacements of the lodes
by the cross-courses, 688-5.
, " precipitation of copper in, 585, 686.
, " statistics of mining in, 457, 694—5,—8; Table XIV.
, ,, injury to meadows, by sand, mud, and foul water from
the mines, 853.
, " subterranean temperature in, 744—6.
Carboniferous limestones, 611,—18,—20,—2.
slates, 614, (?) 616.
Carclaze, area of the open-work at, 576, 665.
Carew, Richard, Esq., on resemblances between lodes and the rocks
they traverse, 23.
, , partition of tin-ores amongst the partners
in stream-works, 94.
,, early copper-mining in Cornwall, 686,-90,
692.
Cargoll, proportion of silver in the lead-ore of, 120.
Carn Brea mines, proportion of metal in the tin-ore of the, 472.
, produce and profits of the, 448,—59; Table XIV.
, winding-engines at the, 143.
Carne, Joseph, Esq., on the veins of Cornwall, 341.
, , produce of parallel lodes on the same
meridian, 258.
, ,, native-silver and silver-ores of Cornwall,
· 110,—14,—16.
, " argentiferous galena " ,
118—20.
, , economy of steam winding- , ,
engines 142.
, ,, substitution of tram-ways for wheel-
barrows in Cornwall, 144.


Carne, Joseph, Esq., on opening mines by means of levels and winzes, 146.
, on the submarine mines of West Cornwall, 600.
Carnon-Stream, on an issue of inflammable gas at, 453.
Carnyorth, on proportion of tin-ore in the vein-stone of, 472.
Carter, Edwin, Esq., on the argentiferous lead-ore of Irebisken-
green, 120—1.
, proportions of tin-ores in the vein-
stones of Lanivet and Mulberry Hill, 472.
Carriage, difficulties and expenses attendant on, in Brazil, 288.
Carvallo, mine of, extent and produce of works at, 124; Table XIV.
Carvosira, formation of, at Catta Preta, 287.
Cascalho (detrital) gold-formations in Brazil, 342-5.
Cascajo, (,,) ,, Chili, 345.
Catalan iron-furnaces, used in Brazil, 219.
Cattas Altas, ancient gold-mines near, 221.
, force employed and gold obtained in the vicinity of,
· 800.
topaz in the neighbourhood of, 800.
Catta Branca, quartzose talc-slate of, 178.
, auriferous quartz of, 179.
quantities and proportions of gold obtained from,
179, 827,—67.
substances allowed with
179, 386.
Catta Preta, auriferous deposits of, 180-2, 287-41.
quantities and qualities of gold obtained at, 237—40.
Caunter lode (Chasarcillo), direction, dip, and produce of the, 81,
92; Table III.
Cavities in the carboniferous limestone, 625.
,, metalliferous deposits, 75, 89, 126, 812,-77, 481, 625,
681, 707.
Cazotte, M., statistics of copper-mining in Chili, 166.
Cellular structure in metalliferous deposits, 75, 89, 126, 812,—77,
481, 625,-81, 707.
Central mine, metalliferous deposit of the, 483.
-, expenditure and profits of, 459; Table XIV.
Chacewater mine, precipitation of copper at, 585.
Chain, quantities of, used for the winding-engines at Carn Brea, 148.
Chalanches, on the rocks of, 517—19.







Cliff mine, rocks of the, 422-8,-87.
, relation between rocks and vein-stones at the, 437-8
Table XII.
, direction, dip, width, and composition of the lode at 423-39.
, relations of copper to the structure of the vein-stone
482.
, proportion of (native) copper in the lode of, 434—6. , ,, fine copper in the crude metal of, at differ ent depths, 436.
, crystals of copper in, 431.
, endlong dip (shoot) of copper in the lods, at, 438.
, native copper sometimes encrusted with native silver, at 436.
, slide, and displacement of the lode by it, 439.
, man-engine at, 439.
, produce and profit at, 440,-59; Table XIV.
, subterranean temperature at, 784.
Cliff mine, South, rocks of the, 460.
, relations between the rock and lode at, 460-2, masses of copper obtained at the, 462.
, proportions of crude metal contained in the vein
stones, 462.
, drift overlying the surface, 460.
Clifford Amalgamated mines, analysis of water from hot spring at the 586.
, subterranean temperature at the, 762.
Climate, of Brazil (Minas Geraes), 349, 726-33,-71-80; Table
XXX. XXXI. XXXVII.
, " Chili (Chanarcillo), 724-5.
, ,, India (Turace of Kumaon), 44.
——, ", Norway, 479.
, ,, (the United Kingdom), Channel Islands, 734-7.
, ,,, England, 744-51.
, ",, Ireland, 787-44.
, ,, (the United States), Michigan, 465,-77,-8,-9, 788-4.
, ,,, Virginia, 782-8.
Cloak, Mr. N. S., on the produce and loss at Carnon Stream, 458
Table XIV.
Closer rocks of 685.7



Collins, Capt. William, on the auriferous jacotinga collected from the
stream at Taboleiro, 252.
Colorada, the rocks of, 70—80.
, the lodes of; their directions, dips, widths, compositions, and structures, 81-92; Table III.
Table III.
, sometimes divide into veins on passing from
one rock to another, 84.
, occasionally enclose (horses) masses resem-
bling the adjoining rocks, 84.
, of cavernous structure, 77, 89.
, unite, 92.
, are enriched at their union with veins, 88, 91.
, richest when most highly inclined, 82.
, kinds, quantities, and qualities of the silver-
ores they afford in the several limestones, 95–121; Table IV.
, exceptionally rich portions of, 90-8.
deflected on the line of their dip by a bed
of limestone which is rich in their vicinity,
82, 114—19.
, one lode in, intersects a cross-vein, 129.
, cross-veins of, their directions, dips, widths, compositions, and structures, 71, 126; Table III.
, intersect, and frequently (heave) displace the lodes, 71, 127—35; Table III.
, displace the rocks in one instance only, 71,
126.
hornblendic dykes at, 80.
subterranean temperature at, 724.
, amounts paid for labour, food, and materials at, Table V.
Combes, M., on the quantities of gunpowder needed for blasting, 201.
, , comparative values of hemp and of wire ropes,
148.
wheel-barrows and tram-
waggons underground, 144.
, , railways in the mines of Cornwall and Devon,
669.
Combe Martin, mines and silver and lead at, 109.
Composition of deposits containing in gneiss, 526,
the ores of antimony.

-	of deposits containing in talcose, micaceous, and chlo the ores of antimony, ritic rocks 179, 884,-7, 617.
	in clay-slate, 335,-7, 545,-8
	566, 707.
	in calcareous, alternating with
	hornblendic, rocks, 118, 617
	in hornblendic and felspathi
	rocks, 121, 535,-6.
	of deposits containing) in talcose, micaceous, and chlo
	the ores of bismuth, ritic rocks, 179, 884,-6,-7
	640.
	, in clay-slate, 111, 334,-5,-6,-7
	374, 546.
	of deposits containing ore of Chromium, 647-9.
	ores of Cobalt, 111,-12,-16,-21
	523—6; Table XVI.
	of deposits containing) in granite, 511, 675-80; Table
	copper and its ores, XXIII.—XXVI.
	- , in gneiss, 526.
	- in micaceous, talcose, and chlo
	ritic rocks, 4, 6-8, 11, 15, 18
	873—8, 495, 596-7, 605—6
	638—41,—53.
	, in felspathic and hornblendi
	rocks, 162—3,—5, 411—12
	420 – 38, – 60 – 5, –8–73, –5, –9
	497, 585–6,–8, 785,–7.
	, in clay-slates, 194—9, 545–50
	555-7,-64-9,-74-8, 676-7.
	, in porphyry (elvan), 441, 677
	Tables XXIII. XXIV.
	, in carboniferous slates, 605—8
	616—18.
	, in carboniferous limestones,
	618—19,—24—6.
	, in the coal-measures, 503—5
	506—10.
	, " New Red Sandstone,
	513—16.
	, ,, Canga, 236, 316.
	— of deposits containing in granite, 175, 311,-20,-32
	gold, { 384,—7.


Composition of denosits contain	ing in talcose, micaceous, and chlo
	old, ritic rocks, 177-80, 210,—47
6	298, 800, -2, -11-12, -15
	820-1,-82,-4,-5,-7,-72-4
	883, 638- 41 .
	—— in quartz-rocks, 181, 321, -32, -4,
	837.
	in clay-slates, 182-4,-94-9, 206,
	207-10,-98-9, 812-18,-21-2,
	828-9,-32,-4-5,-6; Table
	VI.
	in calcareo-siliceous rocks, 245-9,
	815,-23.
	— in Itabirite and Jacotinga, 212-
	216,-21-2,-4,-6-42,-6-7,
	254 – 84, - 6 - 98, 803 - 4,
	813-14,-25-8,-80,-2,-5,-7;
	Table VIII.
	, in Carvoeira, 239-41.
of deposits contain	ning) :ita 175
ores of i	ron, } in granite, 175.
	, in micaceous, talcose, and chlo-
	ritic rocks, 180,-4, 211-86,
	298,-99-300,-3,-13,-16,-18,
	8 21-3,-4-5,-78.
	, in clay-slates, 184,-94-9, 812-
	818,-17,-21-8, 548-9,-58-5,
	568-9.
	in Canga, 216,-36,-47,-8,-99,
	816, 767-8.
of denosits contain	
ores of 1	$\begin{pmatrix} 1116 \\ 122 \end{pmatrix}$ in gneiss, 98, 526.
	—, in felspathic and hornblendic
	rocks, 533-7.
	, in clay-slate, 110-12,-16-21,
	705-14.
	——, in quartz-rocks, 643-4.
	——, in quartz-rocks, 040-4. ——, in carboniferous limestone, 619-
	·
-C 3	621.
of deposits afford	ing in hornblendic slate, 527.

Composition	of deposits affording ore of manganese, in talcose slate, 495.
	, in Jacotinga, 215,-19,-23,-63, 264,-5,-9, 803,-14. in Carvoeira, 289-40.
	of deposits containing the ore of molybdenum, in micaceous slate and porphyry, 654.
	the ores of nickel, in gneiss, 521-6. in hornblendic rocks, 121.
	, in the coal-measures, 503.
	palladium, in granite, 175, 311,-34,-7.
	of deposits containing in Jacotinga, 248,-86, 335,-7, platina, 338,-40.
	of deposits containing silver and its ores, } in gneiss, 103, 521-6.
	in clay-slate, 105,-9-21, 545, 549,-66,-8, 708-11.
	rocks, 533-8.
	, in hornblendic, alternating with calcareous, rocks, 89-94, 98-121.
	of deposits containing in talcose, micaceous, and chlotellurium and its ores, in talcose, micaceous, and chloritic rocks, 180, 299, 315, 334,-6,-7,-76,-82.
	of deposits containing) in granite, 77, 175, 359, 472-3,
	the ores of tin, \\ 486, 631,-64-5,-76,-8,-9.
	, in clay-slate, 77, 202, 359, 472-3,-86, 630,-76,-80.
	, in porphyry, 665.
	of deposits containing the ores of zinc,
	ritic rocks, 617,-40.
	, in clay-slate, 112, 545-7, 707.
	, in hornblendic and felspathic,
	alternating with calcareous, rocks, 86.

Composition of rocks containing the ores of zinc, in carboniferous limestone, 621
of cross-veins, 125, 232,-86,-8, 330,-40-2, 439,-7 526,-58,-98, 608,-82, 716.
Conceicao, rich bunches of gold at, 216.
Concretionary structure, of hornblendic rocks, 702.
, ,, quartzose clay-slate, 702-3.
,,, copper-ore, in granite, 512.
-, ,, osppor ore, in grantes, orz.
, ,, ,, New Red Sandstone,
, ,, lead-ore in clay-slate, 702-3. [513]
, ,, iron-ore in the coal-measures, 66.
Conducting powers of rocks for heat, 753.
Condurrow, proportion of tin-ore in the vein-stone of, 472.
, tin-ore of, separated from its matrix by the natural
action of running-water, 354.
Conglomerate, 30,-9, 394-6, 403-4,-39,-74,-80, 501-3, 601.
Configuration of vein-stones and of the rocks which bound them,
188, 207,–23–4,–8,–30,–46,–59.
Connorree, the rocks of, 542,-63.
deposits of; sitions and structures, 545,—8,—9, 553,-63-8, 738.
, , , ; one of them more highly inclined than the cleavage-planes of the rocks, 564; Table XVIII.
, , , ; argentiferous and auriferous pyrites in,
, precipitation of copper at, 569-71,-87. [568.
, subterranean temperature at, 738.
Consolidated mines, produce, costs, Royalties (Dues), and profits of, 452; Table XIV.
Constancia, extent and produce of works at, 124; Table XIV.
Cook's-kitchen, on the alternations of granite and slate at, 658.
proportion of tin-ore in the vein-stone of, 472.
Coolies (labourers), payment of, in the Himalayan iron-works, 58-9.
Copiapò, consequences of lessening stream in the river of, 139.
Copper and its ores, 4-11,-18, 20, 62, 77,-9, 112,-17-18,-43,-54,-62,
163,-5,-6,-95, 236-7,-86, 334-8,-40,-78, 413,-21-2,-5-86, 461-5,-8-73,-5-8,-80-4,-7-9,-94,-5,-7, 500,-8,-6,-18-16,
525-6,-31,-6,-45-9,-53-7,-61,-4-8,-74-8,-97,-9, 606-7,
617-19-24-6395377-816-948. 707.


Copper, ancient implements of, at Lake Superior, 412-19.
, native, structure of, 431.
, specific gravity of, 429.
, proportion of, in the <i>lodes</i> and vein-stones of Lake Superior, 421,-6-7,-34-6,-40-1,-70-1,-8,-82,-8-9.
, electro-deposited, structure of, 431.
,, specific gravity of, 429–30, after compression, 430.
,, ,, after compression, 430.
, smelted, structure of, 430-1.
, specific gravity of, 429-30.
, precipitation of, 562,-9-92; Table XVIII.a
XXXIV. XXXV.
, ,, in Cornwall, 686-94.
Copper Falls, construction and operation of Stamping-machinery at 426.
, costs of extracting (native) copper at, 426.
, expenditure and profits at, 459.
, subterranean temperature at, 733.
Cornwall Silver-mines, East, proportions of silver in the ores of, 115
Cornwall, on the stanniferous granite of, 175, 664.
, ,, directions of lodes and cross-veins, in, 309, 674 681, 704,-15.
silver eres of 110 19 91 710 11
diminuous anto (Louise) of Jodes in 199
, ,, displacements (neaves) or todes in, 128. , ,, subterranean temperature in, 752, Tables XXXIII XXXIV. XXXV.
, the miners of, frequently employed as Superintendents o
mines in North and South America, 152.
Corri Charmaig, on the chrome-ore of, 643.
Costs of obtaining pyrites (Sulphur-ore) in Wicklow, 562.
native copper at Lake Superior, 426.
gold, by English Mining Companies, in Brazil
827.
,, in California, Table XXII.
Cotta, Prof. von, on the pyrites (Sulphur-ore) of Rammelsberg, 544 546,—52.
Crennis, on the native silver and silver-ores of, 118.
, produce and profits of, 456; Table XIV.


Crennis, East, produce, expenditure, and profits of, 459.
Crenver, ventilation of, 220.
Crickitt, J. K. A., Esq., on a rich bunch of gold at Gongo Soco, 272.
Crofty, Wheal, tin-ore of, separated from its matrix by the natural
action of running-water, 354.
Crofty, East Wheal, subterranean temperature at, 764-5.
Cromlechs in Upper India, 19, 28.
Cronebane, the rocks of, 541.
, ,, metalliferous, their directions, dips, widths, compo-
deposits of; sitions, and structures, 543-57.
, , , masses (Horses) of slate conformably
enclosed in, 551,-3.
, " , mixed with auriferous silver, 549.
,, cross-veins in, their directions, dips, widths, and
composition, 558.
, , displace (heave) the metalliferous
(Sulphur-course) deposit, 558.
, dimensions of holes, weight of powder used, and quantities of stone broken in blasting at, 200.
, precipitation of copper from mine-water at, 572-8,-87.
introduced from Commell 505
, introduced from Cornwall, 585. , subterranean temperature at, 738,-52; Tables XXXIII.
XXXV.
Cross-veins, in granite, (?) 176, 341, 681-5.
, " gneiss, 526–7.
, ,, clay-slate, 558-9,-98, 608, 715-17.
, ,, Itabirite and Jacotinga, 225-6,-32,-86-7, 341-2.
,, hornblendic (trap) rocks, 439,-74.
, " and felspathic, alternating with calca-
reous, rocks. 71, 124-85.
, ,, carboniferous slates, 608,-22.
, ,, limestones, 622.
, directions of, 124, 232, 526,-58,-98, 608,-22,-81, 715.
, approximate the directions of one series
of joints in the rocks, 124, 559, 681.
————, dips of, 124, 282, 474, 526,-58-9,-98, 608,-22,-82, 715.
, at higher angles than those of the lodes, 527, 682.
,, towards bodies of granite, 125, 682.
———, widths of, 125, 226,-32,-86, 341, 439,-74, 527,-58-9,
598, 608,-22,-82, 715.



٠,

Crystalline action, determined in saturated solutions, by the intro
duction of solid substances, 713.
Cubert United mines, on the argentiferous lead-ore of, 120.
Cuddra, proportion of tin-ore in the vein-stone of, 472.
Cumba, proportion of gold in the Jacotinga of, 258.
Cumberland, proportions of silver in the lead-ores of, 108-9.
Cuming, — Esq., on the preparation of lime from shells in Chili, 155.
Damsel, Wheal, profits realized at, Table XIV.
Danda, on the copper-bearing beds of, 7.
Daniel, John, Esq., on the method and cost of stamping vein-stones
near Lake Superior, 425-6.
of extracting large masses of re-
tive copper from the mines, 427.
, " precautions adopted to prevent the effects
of cold in mines, 478, 734.
, , man-engine at the Cliff mine, 439.
Darlington, John, Esq., and Phillips, J. Arthur, Esq., on the produce,
costs, and profits of mines, 459.
Darlington, West Wheal, on the silver-ores of, 116-17.
Darwin, Charles, Esq., on the valley of Copiapo, 140.
, ,, sounds occasioned by moving-sand,
140.
, , beds of shells at considerable elevations
in Chili, 155,-9,-60:
, methods by which ores are brought to
the surface in Chili. 147.
Daubreé, M. A., on the rocks and metallic)
Daubreé, M. A., on the rocks and metallic minerals of Norway, 543-6.
, , silver and silver-ores ,, , 100.
, , (supposed) prevalence of fluor in tin-lodes,
680.
auriferous sands of the Rhine, 348,-51-2,
856,-8.
Daubuisson, M., on the ventilation of mines, 220.
, ,, proportions of silver of Saxony, 101-2.
, " " " France, 104.
Davey, Richard, Esq., on the produce and profits of Wheal Buller,
Wheal Unity, and Poldice, 451-2.
Davey, Capt. Simon, on the ventilation of mines, 220.





```
M. P. van, on the detrital tin-ore of Banca, 631.
Lesch, F. Esq., on the use of sodium amalgam in Brazilian
 reduction-works: Table XXII.
b, of metalliferous deposits, 4, 5, 7, 9, 11, 19, 20,-2,-5,-6,-8, 30,
                                33,-5-7,-9, 41, 71, 80-1, 161,-4,
                                 810,-72,-75-7,-9, 408-9, 520-1,
                                532, -42, -75, -96, 604, -16, -18,
                                 620-3,-37,-47,-52,-74, 704,-37.
                            , usually greatest when rich, 189, 270.
               "
                            , greater when they have a N.—S. than
               "
                                an E.-W. direction, 405-9.
-, of auriferous deposits, generally, conformable to structure of
                                         the rocks adjoining them,
                                         310, -72, -5-7, -9; Table
                                         X.
                                     , parallel to the contour of the
                ,,
                                         nearest mountain ranges,
                                         310; Table X.
-, of the greater number of lodes, towards the nearest granite,
       675,-83.
-, alteration of dip on passing from Manto into limestone at
       Chafarcillo, 82.
 -, of cross-veins, 124,-5, 439, 526,-58,-75,-98, 608,-22,-81,-5,
                      715.
                  , higher than the dip of lodes, 526, 682—3.
                  , towards the granite, 125, 682.
  -, strata different on opposite sides of a cross-vein, 71, 82.
 Ding Dong, proportion of tin-ore in the vein-stone of, 472.
 Directions of metalliferous deposits, 4, 5, 7, 9, 11, 19, 20,-2,-5,-6,
             28, 80,-3,-5,-7,-9, 41, 71, 80-1, 161,-4, 809,-72,-5,-7,
             379, 408-10, 520-1,-32,-42,-75,-96, 603,-16,-18,
             620-3,-37,-47,-52,-74, 707,-37; Tables I. II. X.
           of auriferous deposits usually conform to the strike of the
             rocks above and below them, and to the directions of
             the neighbouring mountain-chains, 310,-72,-5-7,-9;
             Table X.
         - of cross-veins, 124-5, 439, 526,-58,-75,-98, 608,-22,
             681,-5, 715.
 Displacements, 127-35, 225-6,-32,-86-8, 341-2, 439,-74, 503,-27.
                 558-60,-75-6,-98, 600,-22,-81-5, 715-17, (See
```


Heaves. Intersections. Leaps. Throws.)

Displacements, h	orizoi	ntal, of	rocks by n	netallifero (?) 657—	us (veins) deposits 60.
	,,	,			71, 126,-80-1,-3, 225-7,-86-7.
,	"	,	"	"	(partial), 225-6, 286-7.
 ,	"	, of	metallifero depos	us by jo its 18	ints in the rocks, 3, 841, 559.
,	"	,	"	by <i>cr</i> eent	oss-veins of differ- widths. 127,—9, 3, 685.
 ,	"	,	,,	by th	e same <i>cross-vein</i> different depths, 3, 685, 717.
,	,,	,	"	at so	ome, but simple ersections only at er, depths, by the
,	"	,	,,	598 by di the 598	3, 717. fferent branches of same cross-vein, 3; Tables XIX.
,	"			by cr	XIII. XXIV. oss-veins of small al range, 225,-32,
 ,	"	, of o			s deposits (lodes) n, Tables XXIII.
	"	, of 1			alliferous deposits
 ,	"		metallifero		cobalt, 527.
-,	"	,		,,	copper, 558-60, 598, 683-4.
 ,	"	,		,	lead, 716-17.
,	"	,		,,	silver, 127—8, 527,
,	"	,	:	,,	tin, 684; Table XXVI.
 ,	"	, of r	netalliferor	as deposits	iron-pyrites,

iplacements,	horizo	ntal, of	metallifero	us deposits) gold, 183, 225, containing 232,-87, 341.
,	"	, of	metallifero depos	
 ,	"	,	"	in gneiss, 527.
 ,	,,	,	"	in clay-slate, 183, 493,
				558-60,-98, 716-17.
,	"	,	"	in elvan (porphyry), 684 — 5; Tables XXIII. XXIV.
 ,	"	,	"	in felspathic and horn- blendic rocks, 439, 474.
,	, 'n	.,	"	in <i>Itabirite</i> and <i>Jaco-tinga</i> , 225 - 6,-86 - 7, 341.
	, ,,	,	37	in calcareous, alter- nating with felspathic & hornblendic, rocks, 125—9.
	, ,,	,	"	in the coal-measures, 503.
	, ,,	, e	xtent of, 12 685, 715-	7-8, 439, 558,-75,-98, 684-
	, ,;	, , t	owards the g	reater angle, 127-8,-83, 841, 527,-58,-60,-75,-98, 684,
	'e e'	, ,	,, sı	717. maller angle, 127—8, 439, 558, 684, 717.
	' 9 9:	, ,		ght-hand, 127-8,-83, 341, 527,-58,-98, 683-4, 715-17.
	· 2:	, ,		oft-hand, 127—8, 341, 439, 474, 558—60,—75, 684,
Displacement	s, verti	cal, of		715—17. (See Heaves. Intersections.) etalliferous deposits (?), 657—60.
	· ,,	,	" by cro	088-veins, 71, 126,-30,-1,-3, 225,-86-7.
	· "	, horizo	ntal range	,, of small vertical and (partial), 225-6,-86-7, 342.


Displacements, vertical, in silver-mines, 71, 126,-30,-8.
, ,, in gold-mines, 225-6,-86-7, 342.
, ,, towards the greater angle, 126,-30,-1,-3.
, ", " smaller angle, 225-6,-86-7, 341.
(See Leaps. Throws. Intersections.)
Dissemination, of metals and ores through rocks:-
gold, 175,-7,-86,-96, 255,-81,-99, 300,-2,-3,
318,-16,-20,-2,-73 .
silver, 114,-19; Table IV.
copper, 403,-67,-75,-95, 500,-3,-9.
tin, 175, 664,-6.
Divining (Dowsing)-rod, the, still used in Dauphiny, 527.
Debereiner, M., on the platina alloyed with gold in the sands of the
Rhine, 360.
Dol, on the graphite of, 40.
Dolcoath, on the rocks of, 447.
mbiles of the make (Country) and lades at GGG
proportions of tip are in the wain stone of 479
, " separation of tin-ore from its matrix by the natural
action of running-water at, 354.
, ,, ancient works at, 146, 447.
,, improved system of mining at, 448, 645—7,
, conditions under which blasting takes place at, 200.
,,, ventilation at, 220.
,, plans and sections of, 448.
, ,, produce, expenditure, and profit at, 447; Table IV.
Dolcoath, North, on the silver-ore of, 113.
Dolomite, the, of Brazil, 178.
Domeyko, M. I., on the rocks of Quebrada Seca, 161.
, ,, calcareous débris of Chafarcillo, 75.
, " hydraulic lime of " , 75.
, " gold-mines of Chili, 167.
, " detrital gold of Chili, 345,-60.
, " statistics of mines, and miners of Chili,
148-9.
, ,, the climate of Chafarcillo, 724.
Don Pedro North d'el Rey, on the proportion of gold in the Jacotinga
of, Table XXII.
Douglas Houghton (Henwood) mines, on the rocks of, 466-7.
, ,, (copper-bearing),
conglomerate, of, 474.

Duglas Houghton (Henwood)	mines, on the lode of, 468-74.
	copper extracted from, and the cost of ex-
	traction, 471—3. (heaves) displacements
	of the lode by a cross- vein and a joint, 474.
	, ,, formation of ice under-
	ground at, 478, 734.
Downhill, altered value of lan been opened in it, 694.	d at, in consequence of mines having
	s under which blasting takes place at,
	n of tin-ore in the vein-stone of, 472.
Dressing (washing and cleaning	g) auriferous pyrites, 198-9, 208, 857; Tables VI. ^a VII.
	— —— Jacotinga, Table IX.
·	quartz, cost of, Table XXII.
Drift, deposits of, 42, 460-84,	622.
, ,, , containing n	nasses of native copper, 483,-7.
 , ,, ,,	,, gold, 842,-8,-58-9,-83.
	rystals of gold, 356,-9.
Duchanoy, and Rivot, MM., or Hungarian ores, 100-1.	n the proportions of gold and silver in
Ducky, Wheal, on the silver-or	res of, 114—15.
Duck, gold discovered in the c	rop of a, 356.
Duckinfield-colliery, subterran	ean temperature at, 754-5.
Dufrenoy, M., on the detrital g	
Du Noyer, G. V., Esq., on the	e rocks, lodes, and submarine works at Knockmahon, 593-7,-9-601.
,, , , ,,	rocks and <i>lodes</i> of Kenmare, 611, 616,-18,-21-2.
 ,, ,,	limestone of Brownstown, 622.
Dupee & Co., Messrs., on the 1488—9.	production of copper at Lake Superior,
Durham, proportions of silver	in the lead-ores of, 108.
Duras, Capt. Paulo Rodriguez	
	native copper and gold obtained by, 286.



hannes to Chillen miner along her lacked mater 01					
hances to Chilian mines closed by locked gates, 91.					
by, J. S., Esq., on the jointed structure of granite, 75, 672.					
hidote, occurrence of, 80, 398,					
khwege, Baron von, on the elastic sandstone of Villa Rica (Ouro Preto, 172, 210.					
, ,, Itacolumite of Brazil, 172.					
, ,, clay-slate of ,, , 173.					
alternations of Itacolumite and clay- slate, 211, 312.					
, " Itabirite of Brazil, 172, 211,					
298. —————, " Carvoeira of " , 237—8.					
, ,, Carvoera of ,, , 257—8. , ,, gold-mines of, ,, , 184, 244, , 300—1.					
the Canga of, } ,, 217.					
obtained in 366.					
, iron-ores and fur- naces of, 236-7.					
, ,, topaz-mines of ,, , 306.					
Eperanza, extent and produce of the works at 124; Table XIV.					
, quantities and qualities of the ores obtained; Table IV.					
Etheridge, Robert, Esq., on shells in the raised beaches near Caldera, 156.					
, ,, plants in the coal-measures of New					
Brunswick, 504-5.					
Euclase, matrix of the, in Brazil, 308.					
Europeans employed in the mines of Chili and Brazil, 152; Tables V. VI. ^a IX.					
Evans, T. F., Esq., on the different qualities of water at Mona, 584.					
per and production of copper and production of ochre					
, " rain " , 582.					
Evaporation, different amounts of, affected the speed of the water- wheels at Gongo Soco, 351.					
Exmouth and Adams, Wheal, proportions of silver in the lead-ores					
of, 110.					
Expectativa, on the New Red Sandstone, and the copper-ores obtained					
from it, at, 514.					

Expectativa, on petroleum of, 516.



Food, cost of, in Chafarcillo, 150; Table V.
, of slaves in Brazil and Virginia, 292, 382.
-, of serfs in Siberia, 348.
Forbes, David, Esq., on the auriferous deposits of Merioneth, 636-9.
, " quality of the gold of ", 642.
detrital gold of Bolivia, 360.
Forbes, Principal J. D., on the climate of the Alps, 528.
-, , temperature at the surface and at
various depths near Edinburgh, 777
Forbes, Sir John, on a spring of fresh-water beneath the sea at
Botallack, 539.
Forests, 21, 55, 61, 296, 890, 416.
, succession of different growths in, 416.
———, destruction of, 21, 55-6, 843,-90.
, consequences of, 344.
Forest mine, ancient implements discovered in the, 415.
Forges underground at Bearhaven and Wheal Vor, 610
Forster, Westgarth, Esq., on pipe-veins of lead-ore, 620.
" Merioneth ", , 636.
", Кетту ", 613.
Foster, C. Le Neve, Esq., on the Molybdenite of Mount Sorrel, 654.
Foster (J. W., Esq.) & on the rock-formations of Lake Superior,
Whitney (J. D., Esq.), 385-404,-39.
, iron-ores of Lake Superior, 388-9.
, copper-lodes of , , 405—
412,-20-5,-7-8,-81-4, 436,-9,
466-7,-75,-7.
, ,, native silver of Lake Superior,
419,-36-7.
, ,, cross-vein (Slide) at the Cliff mine,
439.
, " drift of Lake Superior, 483-5.
, ,, ancient mining-works near Lake
Superior, and on the weapons,
tools, and implements found in
them, 412-17,-19,-75.
, ,, temperature at Fort Wilkins, 477.

Whitney (J. D., Esq.), Superior, 477.
, ,, subterranean temperature, 733-4.
, " amount of rain at Fort Brady, 477.
Fowey Consolidated mines, on the native silver of, 118.
, ,, produce and profits of, 456; Table
XIV.
Fox, Robert Were, Esq., on the structure of cross-courses, 461.
, " subterranean temperature, 723,-60-1,
Fraga, on the auriferous talc-slate of, 301,-20, 732. [762,-5.
, ,, subterranean temperature at, 732.
Francis, Capt. Francis, on subterranean temperature at Wheal Vor, 764.
Francis, Mr. Henry, on the coincident directions of lodes and glens
in West Cornwall, 718.
Francis, Mr, William, on the gradual filling of Carnon valley with
sand and mud from the mines, 353.
Francisco Nuevo, San, extent and produce of the works at, 124;
Table XIV.
, quantities and qualities of silver-ore obtained
from, Table IV.
Francisco Viejo, San, extent and produce of the works at, 124;
Table XIV.
, quantities and qualities of silver-ore obtained
from, Table IV.
Francisquito, San, extent and produce of the works at, 124; Table XIV
, quantities and qualities of allver-ore obtained
from, Table IV.
Franklin mine, on the costs and profits of the, 459; Table XIV.
Frank Mills, proportion of silver in the lead-ore of, 110.
Freiberg, establishment of the Mining Academy at, 691.
Friendship, Wheal, on the inclined tramways of, 144.
produce and profits of, 459; Table XIV
Friendship, North Wheal, proportions of silver in the lead-ores of 110.
Fuel, charcoal as, manufacture and cost of, 55-6, 219,-61-2, 390
,, , from hard woods better than from soft, 55-6, 219
, coal cheaper than wood as, even in the forests of Lak Superior, 478.

Fuller, Dr. Thomas, on the silver of North Devon, 109.
Furnaces, Himalayan, blast supplied by air-bags of skin, worked by
hand, 54-5.
, Brazilian, blast supplied by fall of water into a pneu-
matic trough, 219,-60-1.
Furze Hill Wood, proportion of tin-ore in the vein-stone of, 472.
•
Gad and pick, skill of Cornish miners in using, 152.
Galena, 100-10,-16-21, 376-7, 545,-8-9, 615,-20,-40, 708,-8-11.
, argentiferous, 100-10,-16-21, 545,-8-9, 615-20,-(?) 40,
708,-8-11.
,, , enclosing threads of native silver and masses
of vitreous ore, 120-1.
, auriferous, 100, 878, 549, (?) 640.
Gallà, iron-mine of, 22; Table I.
Gama, Col. Ignacio da, on the gold-mine of Catta Preta, 287.
Gambling, propensity of Chilian miners towards, 91.
Garby, John, Esq., on the native silver and silver-ores of Cornwall,
Gardner, Dr. Daniel, on the gold-mine of Catta Preta, 237. [118.
Gardner, George, Esq., on the gold-mines of Brazil, 170,-85, 245,
803, 500-2.
, ,, detrital gold of ,, , 348.
, " fishes of " , 354—5.
, " scenery of " , 169, 297.
Garnets in the copper-bearing rocks of Chili, 161.
" auriferous deposits of Virginia, 381.
Garnett and Moseley mines, on the rocks of the, 379-81, 732.
auriferous deposit of the, 381-3.
, " proportions of gold at different
depths in the, 383.
, " quality of the gold, 381.
, , detrital gold of the neighbour-
hood, 384.
, ,, subterranean temperature at the
788,
, , food of the slaves at the, 382.
Garras (Guarneck), proportions of silver in the lead-ores of, 118.
Garras, South, proportions of silver in the lead-ores of, 120.
Gas, issue of, from joints in the Itabirite at Gongo Soco, 250.
-, inflammable, from crevices in the auriferous formation at Morre
Velho, 195—6.

Gas, inflammable, from vegetable matter overlying stream-tin-ore near Perran Wharf, 453.
——, coal, used for lighting the underground works at Balleswidden, 669.
Geach and Webb, Messrs., on the Caradon, Menheniot, Lanreath, and Saint Pinnock districts, 659,-61,-4-5,-76-7,-95, 704,-15-16. Geikie (Archibald, Esq.) and Murchison (Sir Roderick Impey), on the gneissose rocks of Breadalbane, 646. Georgia, on the auriferous rocks of, 371.
, ,, an ancient Indian village entombed in the auriferous drift of, 384.
Gesner, Dr. A., on copper-mining in New Brunswick, 506. Geyserberg, on the stanniferous porphyry of, 665.
Gilbert, C. S., Esq., on the silver-ores of Cornwall, 111,—14. Gilbert, Davies, Esq., on the argentiferous lead-ore of Wheal Rose, 119.
Giles, Mr. John, on the slates of the Caradon district, 667. ——————————————————————————————————
afford, 708. Gillman, Robert, Esq., on the produce of gold at Clogau, 641;
Globe, the, specific gravity of, 186. [Table XXII. Gneiss, the, of Dauphiny, 517.
, ,, of the Garnet and Moseley mines, 379. , ,, of Breadalbane, 646. Goats used as beasts of burden in Upper India, 47. Godoi, Juan, his discovery of silver in Chañarcillo, 69. Godolphin mine, profits of, 445; Table XIV.
Gola, beds of gravel and sand above the bed of the river at, 45. Gold, of Australia, 175,-9, 343,-56,-9; Table XXII. ——, ,, Banca, 631. ——, ,, Bolivia, 360.
——, "Brazil, 163-870; Tables X. XXII. ——, " California, 338-9,-43; Table XXII. ——, " Chili, 166, 345,-60,-78. India 3 46-7

Gold, of Ireland, 549-50, 627-84; Table XXII.					
, ,, New Granada, 860.					
, ,, the	e Rhine, 348,-	56,-8,-60 ; Table XXII.			
, ", Sil	peria, 175,-7, 8	85-6,-40,-2,-3,-5,-6,-8,-60; Table XXII.			
—, " Vi	rginia, 371-84	; Table XXII.			
, ,, W	ales, 685-42;	Table XXII.			
		h granite, 3, 175, 320.			
,	"	talcose, micaceous, and chloritic rocks,			
•		177, 800-3,-20.			
—,	"	quartz-rocks, 316.			
—,	"	clay-slate, 186,-96, 210.			
 ,	77	calcareo-siliceous rocks, 245,-9, 304,			
		315,-19,-23-4.			
 ,	27	Itabirite (Jacotinga), 215,-19,-21,-42.			
,	99	Carvoeira, 240.			
—-,	"	Canga, 217, 315-16,-19.			
, limite	ed to certain p	ortions } 175, 311,-20,-34.			
	or beds of g	granite, \ \ 110, 011, 20, 012.			
~ _,	"	talcose, micaceous, and chloritic			
		rocks, 176-7,-80, 298, 302,			
		311 - 12, - 15, -19, -20, -34, -72,			
		636,-42.			
 ,	,,	quartz-rocks, 178-9,-80-2, 312,			
		334.			
 ,	"	clay-slate, 182,-3,-4, 312-13,-17,			
		318,-20,-35,-49-50,-76.			
 .	,,	calcareo-siliceous rocks, 636-42.			
 ,	,,	Itabirite (Jacotinga), 214,—16,			
		221,-7,-35,-42,-6,-62-82, 313-			
		314,-18-19,-25,-35,-75.			
 ,	"	Carvoeira, 240.			
 ,	,,	which conform to the cleavage of			
		the rocks, 178,-80,-2,-3,-4,-7,			
		188, 207,-14,-16,-21,-3,-7,			
		243,-5,-9,-98, 300,-1,-2,-9-23,			
		372, -5,-7,-80.			
 ,	"	which conform to the joints, 181,			
		187, -8, -91, 207, -21, -3, -99,			
		323.			
-, limited to certain portions) which conform neither to cleavage-					
•		f rock, planes nor joints, 637.			
		· - ·			

Gold,	matrix	of, its c	omposition,	in g	ranite, 46, 175, 811,-82,-4,-7.
	. ,,				alcose, micaceous, and chloritic
•	•	·		•	rocks, 177-80, 209-11,-47,
					298-803,-11-12,-15,-20-1,
					332,-4-7,-72-83, 639-42.
				αt	artz-rocks, 181, 321,-82,-4,-7.
,	, ,,	,			ay-slates, 182-4,-94-9, 206-7,
—-,	79	,	" ,	,, CE	210,-98-9, 312-13,-21-2,-8-
					329,-32,-4,-5,-6,-7; Table
					VI.
,	"	,	"	,, са	lcareo-siliceous rocks, 245-9,
					815,-28-4.
	77	,	" ,	" It	abirite (Jacotinga), 212—16,
					221-4,-6-35,-42,-5-7,-54-84,
					286,-98, 808-4,-13-14,-24-
					330,-2,-4-7; Table VIII.
 ,	"	,			rvoeira, 289–41.
 ,	"	, dip o	f (Shoots)	maa	ses coincident with indulations
					16,-34,-59,-64,-9, 319,-23,-6.
 ,	most pl	entiful i	n most high	hly ir	clined portions of formations,
		270,-81,		•	•
				e) of	pyritic quartz in clay-slate,
,			f its matrix		184,—97.
,		"			Jacotinga (Itabirite), 259,-70,
•		••			824,-5,-8,-9.
	relation	between	it and the) in	clay-slate, 199-201, 328;
•			its matrix		Tables VII. XXII.
,		"		in	Jacotinga (Itabirite), 232,
•		•••			329.
—,	differen	t distrib	ution of, in	diffe	rent ores of iron, 328.
,	crystals	of, in	shallower p	parts	of the rocks and in detritus,
-		•			215,-36,-41, 803,-59.
 ,	proport	ion of, di	isseminated) tal	cose, micaceous, and chloritic
•	• •	•	through		rocks, 177.
 ,	2)	,	"	qu	artz-rocks, 182, 821.
	"	,	33		y-slates, 186,-96, 321.
	99	,	,,		careo-siliceous rocks, 249,
•	•	•	••		321.
—,	,,		"	Ja	cotinga (Itabirite), 255, 325.
,	"	, cc			talcose, micaceous, & chloritic
,	"		ertain beds		rocks, 177,-9, 320,-7,-73,
				•	883, 688,-41.

Gold,	proportio		ntained in artain beds		rocks, 182, 321.
 ,	"	,	"	of clay-sla	te, 184,-96-8, 201-6, 827,-79.
 ,	"	,	"		te (Jacotinga), 290-2, 79-81, 326,—7.
 ,	**	, ob	tained from different	•	e same formation, 383.
 ,	"	,	**	depths in	the clay-slate series, 201-5, 331; Tables X. XXII.
 ,	"	,	")	Itabirite (Jacotinga), 280-2,-5,-79-81, 881; Tables X. XXII.
—,	77	,	"	22	Carvoeira, 241.
 ,	29	, (n	neggets) ma		ust in <i>Itabirite (Jaco-</i> <i>tinga)</i> , 274-7, 330.
—,	analysis	of, 334,	-5,-60,-74.		
			mite, 382.		
<u> </u>			•	oons and ch	loritic rocks, 832,-74,
,	"		-	·	383, 642; Table X.
 ,	29			5-6, 832,- 88	
,	"	, "Iu	ıbirite (Jac	otinga), 236	5,-85-6, 832,-8; Table X.
,	"	, at diff	ferent depti		ne formations, 205-6, 75; Tables VII. X.
		diff	- int-1	-	t forms, 835.
,	"			or ameren	t tornes, ooo.
,	nature a	of its	alloy \ " E	granite, 175,	
 ,		"	,, t		ceous, and chloritic ,-80, 299, 384,-7,-74,
,		"	" 9	uartz-rocks,	334,-7.
 ,		"	" c	lay-slate, 20 377–8.	06, 884—5,—7,—40,
—,		,,	C		eous rocks, 642.
					cotinga), 236, 885,-7.
,		"		840.	
		"	,, a		depths in the same -86,338,-40; Tables
	5 x			VII. Х	, , , , , , , , , , , , , , , , , , , ,

```
Gold, alloyed with antimony, 179,-80, 334-7,-74.
                  arsenic, 835,-7,-74.
           11
                  bismuth, 174, 334-7,-74.
           "
                  copper, 286, 335-7,-40,-59,-74,-5, 642.
           "
                  iron, 835-6,-74, 632,-42.
           "
                  lead, 335-7,-40,-74,-7.
           ,,
                  palladium, 175, 215,-86, 335-7,-40,-59.
           "
                  platina, 286, 335-7,-40,-60.
           ,,
                  silver, 236,-86, 334-5,-7,-40,-59,-60,-74,-7,-8,
           "
                            632-42.
                  tellurium, 180, 299, 334,-6-7,-82.
  -, molten, of higher specific gravity than native, gold, 335.
    --mines, temperature of, 725—33,—56—8; Tables XXXII.—
                                                          XXXVI.
----, detrital, ancient deposits of, 342-51,-83-4, 627-33,-42;
                                       Table XXII.
                                   , 46, 283, 348,-52-9,-84.
             . recent
         ••
             , contains the same alloy as the gold of the neighbouring
         99
                 mines, 359.
             , always of superior quality to mine-gold, 859-60, 632.
         "
             , crystals of gold not uncommon, 356,-9, 632.
             , shaped implements sometimes mixed with, 344-5.
             , proportions of, extracted from sand and gravel, 346,
             , occasionally swallowed by water-fowl, 356.
                           extracted from mud in the streets, 217.
    -, cost of extracting, in various countries, 327,—46,—8,—58;
                                                     Table XXII.
   -, quantity of, obtained in Australia, Table XXII.
                                Brazil, 866-70; Table XXII.
                        "
                                 California, Table XXII.
           ,,
                        "
                                 Ireland, 633.
                        "
           "
                                Nova Scotia, Table XXII.
                                 Siberia, 346; Table XXII.
           ,,
                                 Wales, 641; Table XXII.
                        "
    -, Royalties (Dues) and Duties in Brazil, 208,-89,361-8;
                                                 Tables VII. IX.
                   paid by miners, \( \)
                                  , "Siberia, 327.
                         "
Golden, Wheal, proportion of silver in the lead-ore of, 120.
Gold-Hill (California), auriferous deposit of, 839.
                    -, alloy of gold from, 339.
Gonamena, on the stanniferous granite of, 664-5.
```

INDEX.

		erranean temperature at, 746.
Gongo Soco, the	a rocks o	of, their direction and dip, 248—9.
 ,	"	, joints in, 250.
 ,	"	slate (auriferous at Camara), 183, 247,
		827,-41.
 ,	"	calcareo-siliceous matter mixed with par-
		ticles of gold, 249, 804,-19.
 ,	. ,,	Itabirite, 249,-97, 818.
 ,	"	", ", mass (Horse) of, imbedded in
		(yet containing lines of)
		auriferous Jacotinga, 251,
		824; Table VIII.
,	"	", displaced by small cross-veins,
		286—8, 341.
 ,	"	Jacotinga, the Gongo formation, 250,
		252-4.
 ,	"	", ", "Cumba ", 252-4,
		256- 8,
	>>	", ", ", ", both
		conform to the direction, dip, and
		cleavage of the rocks above and below
		them, 248, 310,—25.
 ,	"	Jacotinga, the Cumba formation, at times
		thinly sprinkled with gold,
		255.
 ,	"	", productive, mostly, on certain
		parallel lines (veins) in both,
		255,-7,-62,-4,-5,-8 2 .
,	"	", productive, mostly, on certain
		thin lines (veins) of limited
		horizontal & vertical range,
		257,-65-7,-70,-1,-82.
 ,	"	", productive on the same meridian
		269,-70.
,	,,	", productive masses (shoots) of,
		all dip (endlong, with the
		groovings of the rocks), to-
		wards the E., 259,-68,-4,-6,
		270,-82.
	"	", productive portions the most
		highly inclined, 281, 310.

Fongo	Soco.	the	rocks of	Jacotinga	, F	roduc	tive, c	composition of, 268-
	,			•	•	,	-	269, 314,-25-7.
			,,	11		"	, 1	nardness of, 283-4.
		,	"	"	,	"	, (disposition of masses
		•	•	,,			(nu	ggets) and particles
							of g	gold in, 271—80.
		,	,,	,,	,	"	,1	proportions of gold
								in, 274-7,-9-81;
								XXII
		٠,	,,	"	,	produ	ctive,	quality of gold con-
								at different depths,
						289,	, 555 - 6	; Table X.
		,	"	"	,	nature	oc pi	coportion of alloy of ained in, at different
						gora	COLU	86, 838; Table X.
						debr	ativo	lines take, at times,
	-	٠,	"	"	,	prou u	diron	tions of the joints,
						264		atoms of all jems,
		_			_			influence of the
		•	"	"	,			252,-67,-70.
		•.						ve portions of, 270.
		••	"	"				unches) of iron-ore
		,	"	"	•			260-1.
		-,	"	magnesiar	ı-l			
		-, ⊽€		n of works	b	y air	issuin	g from joints in the
			Itabirit	e, 250.				
		-, al	imate of	E, 349,-52,	7	26; T	able :	XXX.
		-, BC	ıbterran	ean tempe	rai	ture at	, 727	-9,-51,-6,-8; Tables
								XXXII.—XXXVII.
				t, 290—6.				
								Table XXII.
		-, q	uantitie	of gold o	bti	ained	by E	uropean miners, 284.
		-, a	mount o	i Duties pe	ud - 1	to the	Braz	ilian Government at,
			289, 80	87; Table	8 1	And D	. V • 	. 920 S67 · Tables
		-, p	roauce,	exbenmen exbenmen	₽, 	ани р	a ve (at, 289, 367; Tables 290—7. [IX. XIV.
Good	oin T	-, n	names, b	n the mo	u i	ation (of go	ld in iron-pyrites at
			184, 82		r		- 5º	
Goori	khaa 4	atter	ntion to	the draina	œ	of mi	nes b	v the, 50.
Gord	on. J.	N.	Esa. or	the rain-	o∨ [a]	l at M	orro	Velho, 349.
	,	~~·,	, v			nean		
			,	temper				, , 726–7.

Gorland, Wheal, on the profit made at, 451; Table XIV.
Goron, on the copper-bearing talc-slate of, 5.
Gossan, a matrix of copper-ore, 162,-5,-7, 545,-64,-95, 605,-16,
——, ,, lead-ore, 706-7. [624,-39,-76,-8.
, ,, silver-ore, 525,-34.
, " gold, 167,-83, 299, 301,-2,-12,-73,-5-7,-81, 549, 639.
Graham (Callcott), Mrs., on the elevation of the coasts of the Pacific
by earthquakes, 157.
Granada, New, on the different qualities of mine and detrital gold in, 360.
Granite, composition of, 2, 29, 170,-4-6, 247, 311,-85-7, 662-4,
———, auriferous, 3, 175-6, 311,-16,-20. [666-7,-72.
, quality of gold in, 332.
, alloy ,, , 334.
——, stanniferous, 175, 664-6.
——, cupriferous, 512.
, structure of, 3, 316,-85, 512, 673.
——, cavern in, 27.
, position of, in opposite sides (walls) of lodes, 657-60,
, interlying slates, 658-9. [669-70.
, veins of, 3, 27, 386-7,-91, 490-2, 512.
, relation of its contour to the cleavage of contiguous slates, 259,-310.
, relative position to the dips (shoots) of ore in lodes, 32,
, rate of blasting in, 200. [122, 259.
, quantity of it shipped from Cornwall, 666-7.
, subterranean temperature in, 756,-8,-60-1; Tables XXXIII.
Graphite in the clay-slate of Upper India, 41. [XXXV.
Grasty (Chancellorsville), auriferous rocks of, 372-5.
Gravel, beds of, at great altitudes, 45, 66, 70, 156.
Gravity, specific, of native copper, 429.
, ,, electro-deposited copper, 429-30.
, ,, ,, ,, ,, ,after compression, 430
, ", " smelted copper, 429–30.
, ,, native gold, 835, 431, 632.
, " , molten " , 385, 481.
Great Work, proportion of tin-ore in vein-stone of, 472.
Greenstone, 79, 398, 496-7,-9, 650-1.
money ave, van

Greenstone, dykes of, 80.
Greenwich, extreme and mean temperatures at the surface and at
various depths, 776-7.
Greer, R. J., Esq., on the mean temperature of Waterford, 739.
Gregg (R. H., Esq.) & Lettsom (W. G., Esq.), on the silver-ores of
Cornwall, 116,-18.
Gregory, Capt. Thomas, on the rate of blasting at Drake Walls, 200.
, " proportions of tin-ore in the vein-stone of , , 472.
Grylls and Basset, proportion of tin-ore in the vein-stone of, 472.
Gualaxo, character of gold obtained from the, 359.
Guernsey, on the climate of, 735.
Gueymard, M. E., on the rocks of Chalanches, 518-19.
———, " metalliferous deposits of " , 519-26.
, ,, climate of the Alps of Dauphiny, 528.
Guias de Carvallo, veins (branches) of, 81.
, horses worked underground at, 76.
Gungolee, on the copper-mines of, 4, 5.
Gunnai, auriferous sands of the Ramgunga near, 5.
Gunnis Lake, produce and profits of, 457; Table XIV.
Gurhwal, on the iron-furnaces of, 57.
, Government Royalties of, 62-3.
Guy, Capt. Benj., proportions of gold from the calcareo-siliceous
formation at Gongo Soco, 249.
Gwinear, subterranean temperature in the district of, 752.
Haidinger, H.W. von, on changes in the characters of copper-ores,
Hall, Capt. Basil, on the valley of Copiapo, 139. [679.
, mode of bringing ores out of the Chilian mines, 147.
Hall, James, Esq., on fossils from the shores of Lake Superior, 393.
Hals, William, Esq., on early copper-mining in Cornwall, 686,-90,-2.
Hambly, Capt. W., observations on Gongo Soco, 257, 350.
Hammer (Mallet), use of by miners in different districts, 151.
Hansteen, Prof. Chr., on Magnetic declination, 520.
Hardness, a favourable indication in the auriferous 328; Table deposit of Morro Velho, XXII.
, at different depths ,, 199, 206; Table XXII.
Harris, Capt. N., on the auriferous deposit of Gongo Soco, 250,-5,

Harris, Capt. N., on the manufacture of iron at Gongo Soco, 2 Harris, Sir W. Snow, on the climate of Plymouth, 745,-9.	62.
Hartz, proportions of silver in the lead-ores of the, 101.	
Haughton, The Rev. Dr., on the deposits of sulphur ores in Wick	low.
547,-58,-5	
make of the Kenmare district f	
• • • • • • • • • • • • • • • • • • • •	614.
mandallifamana)	514-
	621.
, ,, cross-vein at Ardtully, 622.	
, , , drift at Kenmare, 622.	
, " precipitation of copper at Be	ally-
murtagh, 574.	
, analysis of the hematite of Ballymurt	tagh, 47.
, " limestone of Kerry, 6	13.
grey copper-ore of Ardtully,	817.
Hawkins, John, Esq., on the throws of lodes in Cornwall, 226.	
, ,, copper-ores of the Parys moun	tain,
574,	_
, ,, stanniferous porphyry of Geyers	berg, 65.
Heaves, 127-85, 225-6,-32,-86-8, 841-2, 439,-74, 508,-27,-58	-60,
575-6,-98, 600,-22,-81-5, 715-	
(See Displacements. Intersections. Leaps. Throws.)	
Heber, Bishop, on the frequency of earthquakes in Kumaon, 4	9.
Helmreichen, H. Virgil von, on displacements of beds of aurife	erous
granite at Candonga, 176.	
, on the geology of Gongo Soco,	170,
250,-68,-6,-8, 815	,-54.
, ,, manufacture of iron at G	longo
Soco, 262.	
, ,, calcareous rocks of Ante	onio
, ,, flint arrow-heads from diam	ond-
washings in Brazil, 344.	
, ,, Magnetic declination in M Geraës, 177.	(lin a s
Helston, subterranean temperature in the district of, 752.	
Hemp, rope of, inferior to iron-wire rope, 143.	


Hender, Wheal, proportion of silver in the lead-ore of, 120.
Hennezel, M. de, proportion of silver in the lead-ore of Bohemia, 101.
Henry, Dr. William, on the argentiferous lead-ores of Crone Bawn, 745.
, " precipitation , 572.
of copper at) ", ", ", ", ", ", ", ", ", ", ", ", ",
Henwood, contact of granite and slate at, 655,—7.
Henwood, Mr. John, a shareholder in the first Cornish silver-mine,
Henwood mines, 465-79. (See Douglass Houghton.) [111.
Herbert, Capt., on the auriferous granite of Kedarnath, 3.
, ,, sands of the Aluknunda, 46.
, , metalliferous rocks of the Himalays, 12, 21.
works of } 41,55.
, ,, drainage of the Dehra Doon, 42.
, ,, disappearance and reappearance of rivers in
the Terrai, 44.
Herland, on the silver and silver-ores of, 111—12.
Herm, on the metalliferous deposits of, 786—7.
, " subterranean temperature at, 737.
Herod's-foot, on the rocks of, 700; Table XXII.
, " organic remains they contain, 700.
, ,, lode of, 704—15.
, proportions of silver contained in the lead-ores of, 120, 709.
, ,, cross-vein of, and the heave it occasions, 715-17.
, , machinery in use, population employed, produce,
and profits at, 719—20.
Heron, Robert, Esq., on the produce and profits of Knockmahon, 442;
Table XIV.
Herring, Charles, Esq., on mass (shoot) of ore at Morro Velho, 207.
, single-handed boring at , , 199.
, on the specific gravity of the ore at VII.
, " loss of mercury) ", Table
during amalgamation at \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \
Hewas, proportion of silver in the lead-ore of, 120.
Hickson, Mr. Richard, on a rich mass of gold at Gongo Soco, 272.
Hides, their use in separating gold from stamped ore, 203; Tables
Hiel, nature and proportion of alloy in the gold of, 335. [VII. IX.
Higgs, Samuel, Esq., on the produce and profits of the Providence
mines, 444; Table XIV.

Higgs, Samuel, Jr., Esq., on copper-ore in the New Red Sandstone of Alderley, 516.
Hilaire, M. A. de Saint, on the gold-mines of Brazil, 185, 237,-43-4.
,, implements used in them, 218.
inon formone of Ging 918
details and of Desail 849
, , scenery and , , 168, 209-10, vegetation 236.
Hill, — Esq., on the direction, dip, width, and composition of the Cliff lode, 423.
, , occurrence of native in , , 486.
Hill, W. G., Esq., on the proportion of tin-ore in the vein-stone at
Wendron Consols, 472.
Himalaya, Mining in the, 4-42.
, inefficient tools used by miners in the, 51—3.
, iron-smelting in the, 53-62.
, varied occupations of the inhabitants of the, 56.
, earnings, , , 60-1.
Hind, H. Y., Esq., on the granite of New Brunswick, 490,-1,-8.
missesser alots of 400
andstones and)
conglomerates of , , 501,-2,-3.
———, " cupriferous shales of " , 503,-6.
, " copper-ores of " , 500.
, ,, detrital gold of ,, , 498, 500.
Hitchens, Capt. Joel, on the suriferous deposit of Agoa Quents, 233.
Hitchins, The Rev. M., on the silver and silver-ore of Herland, 111.
Hoare, J. S., Esq., on the detrital gold of Wicklow, 688.
Hocheder, J. C., Esq., on the auriferous deposit of Gongo Soco, 178,
251,-5,-6,-60,-2,-5,-9,-71,-3,-4; Table XXII.
Hockin, John, Esq., on the qualities of gold from different)
depths in <i>Morro</i> 205.
Velho)
, , nature of the , , 206.
analysis of 994 74
note of blooding 900
, " rain-fall at Morro Velho, 849.
, ,, climate of ,, , 726.
, subterranean , 727.
temperature at) ", '21.

Hodge, John, Esq., on the (sulphur-ore) of Wicklow, 545,-7,-8,-66.
, " copper-ores " , 547–8.
, ,, auriferous iron-ores of Ballymurtagh, 550.
, " unfavourable influence) Sulphur-course
of cross-veins on the of Ovoca, 560.
, ,, displacements (heaves) of the ,, , 558.
, " detrital gold of Wicklow, 629. [559,-60.
, " precipitation of copper at Ballymurtagh,
574-8.
Hoffman, Prof., on the dissemination of gold through granite and
slate, 175.
Holdsworth, — Esq., on the rocks of Bonmahon, 597.
Holes, experiments on boring and blasting, 199, 200; Tables VII.
XXII.
Holl, Dr. H. B., on the hornblendic rocks of the Caradon district, 662.
, " fossiliferous " " Liskeard " ,701.
Hollingshed, R., Esq., on the proportion of silver in the lead-ores of
Devon, 109.
Hollow, Capt. W., on the proportion of tin-ore in the vein-stone of
Providence Mines, 472.
Holman, Capt. James, on the ventilation of mines, 220.
Holmbush, on the proportions of silver in the lead-ores of, 120.
Hooker, Dr. J. D., on the Coal of Fitcooree, 65.
, " structure of Himalayan sandstone, 66.
Hopkins, Evan, Esq., his experiments on the auriferous iron-ores of
Ballymurtagh, 550.
Hopkins, William, Esq., his observations on the conducting powers of rocks, 758.
Hore, Mr. James, on ancient works at Knockmahon, 753.
Hornblendic granite, 2, 511,-17.
gneiss, 517.
and felspathic rocks, 72,-3,-9,-80, 154,-62-5,-73, 386_
387,-8, 530,-1,-41, 650,-3,
701–2.
,, influence of on silver-lodes, 86,-8.
, subterranean temperature in, 754,
755,-8; Tables XXXIII. XXXV.
slates, 519, 646-7,-61-2,-71-2.
Horses (masses) of granite enclosed in lodes, 683; Table XXIII.
tologo and missessus slate in Jades 991
,, calcone and micaceous siste in todes, 301.

Herees, (masses) of clay-slate in lodes, 190,—4, 812,-17,-28, 550, 551,-68; Tables XXVII. XXVIII. XXIX.
felspathic and horn- blendic rocks, in lodes, 424; Table XVII.
,, Itabirite enveloped by Jacotinga, 250-1.
,, intersected by cross-veins, 287.
micaceous and oxydulated iron imbedded in Carvoeira, 240.
,, carboniferous slate ,, lodes, 605.
,, limestone ,, ,, 618,-20,-4.
penetrated by lines of the ores which envelope them, 251, 551,-96, 718.
, their cleavage-planes mostly coincide with those of neigh-
bouring slates, 28, 190, 251, 324,
551,-64,-96, 605, 718.
, ,, are not always coincident, 713.
Horses and mules, their liking for water holding iron-ores in sus-
pension, 354.
Houghton, Dr. Douglass, on the metalliferous rocks of Lake Superior,
487.
in the drift of , ,484.
Howe, T. M., Esq., on financial operations at the Cliff mine, 440;
Table XIV.
Huidobro, ores of copper in New Red Sandstone at, 77, 518-16.
, petroleum obtained from ,, , 516.
Huldwanee, works of irrigation at, 48.
Humboldt, M. A. de, on the disposition of silver-ore in veins, 327.
, ,, proportions of silver in the ores of
Mexico, Peru, and Chili, 97-9.
, ,, comparative values of wheel-barrows &
tram-waggons in mines, 145.
, ,, loads brought to the surface by Mexican
labourers, 147.
Humming-birds, their nests and note, 181.
Hungary, proportions of gold, silver, and lead in the ores of, 100-1.
Hunt, Robert, Esq., on the proportions of silver in the lead-ores of
the British Isles, 105,-6,-7,-8,-9,-10,
120.

Hunt, Robert, E		ores of	d proportions of silver in less East Cornwall, 708-9,-20.	
	, ,, pi	oduce of	copper-ores in East Cornwe	all, 98.
	, ,,	" W	silver-ore at \ , 7. heal Ludcott, \ , , , 7.	11.
Hurchinolee, on	the iron-ore	s of, 33,-	-5.	
Ice, formation of	, in grounds	where the	he Bamboo flourishes, 45.	
 , ,,	, in mines,	165,-78-	9, 784.	
Inclination of pa Chilian mines		h ores are	e taken out of the Mexican	and
Indiana mine, o	n the rocks	and lodes	of the, 464.	
Inficionado, auri				
Intersections, un		•		
intersections, th	mocompanie.	atalliforor	18) 81,-122,-87, 818, 420,	_3.
	rocks by m	oraniusus teorrak	a, \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	-80.
		dehopri	515,-20,-31,-2,-64,-	-96.
			603, -25, -57, -60, -8	- 9 .
			674, 704.	, ,
			, which are elsewhere the	em-
	"	"	selves intersected by	
			same rocks, 192, 318.	
			, which are deflected by the	nem
	"	,,	in the same mines, 82	
	" by c	ross-veins	, 125,-9, 287, 474, 515,-	-26,
			608,-22,-81, 71	
	"	"	, some of which displace neighbouring strata, 18	
	. 1110			
of	metallitero	ra ceboeri	by rocks, 176,-92-8, 818.	,
	99		by other metalliferous for tions, 92, 127, 538.	ma-
	"		by cross-veins, 608,-22,-	-84, 17.
			by cross-veins, which sin	
	"		traverse them at so	
			but displace then	
			other, depths, 717.	
-			by cross-veins, which sin	
	"		traverse them but	dis-
			place other similar	r de-
			posits, Table XXI	V.
			4	



Iron, ores	of, in Coal-me	asures, 64-8, 510.
		d Sandstone, 514—15.
	~ .	
 , ,,		217,-36,-45,-8,-98,-9, 315.
		alaya, 48—55.
		219-20,-60-2.
		nalaya, 53—62.
		Superior, 388—90.
		Jpper India, 43.
		ct of, 214—18.
 ,	"	, quantities of gold obtained in, 216-17
	,,	qualities of
 ,	"	, mode of working and description of
,	,,	tools used in, 217—18.
 , ·	,,	, gold extracted from mud in the street
•	,	at, 217.
, ma	nufacture of i	ron in the neighbourhood of, 218—20.
Itabira, mi	ne of, rocks an	d metalliferous deposits of, 214—18.
Itabirite, de	erivation of th	e name, 172, 214.
		211-16,-19,-21,-4,-7,-42,-4,-8,-9,-50,-1
•	•	287,-97,-8, 803,-13,-24,-88-9, 767.
, st	ructure of, 21	11,-14,-16,-21,-3,-5,-7,-42,-9,-50,-1,-98
, co	ontains little g	old, 219,-42,-5,-9. [824,-9,-89, 767]
, m	asses (Horses) of, imbedded in, and veined with, aurifer-
	ous Jacotinga,	251,-65,-87, 324.
		low cross-veins of quartz sometimes simply
	traverse, but s	cometimes displace, the rocks, 225,-87, 341.
		the name, 178.
	, composition a	and structure of, 170,-2,-8, 301.
Ives, Saint,	subterranean	temperature in the neighbourhood of, 752.
Ives, Saint,	Consolidated	Mines, proportions of tin-ore in the vein-
		stones of the, 472.
		, profits of the, 443; Table XIV.
		•
Jackson, D	r. C. T., on the	e drift copper of Lake Superior, 485.
		ehistoric mining near ,, , 412,-14.
	_	e lodes, ,, ,420,—8,
		465,-79.
	 , ,,	distribution of copper in, ,, , 433.

Jackson, Dr. C. T., on the purity of copper near Lake Superior, 418.
, ,, mode and cost of)
extracting copper from, ,426,-8.
, ,, native silver of ,, , 436.
, subterranean tem-
perature)
, & Alger, Esq., on copper-ores of the Nova Scotian coal-measures, 507.
Jacotinga, derivation of the name, 214.
place of in the <i>Itabirite</i> formation, 214,-16,-19,-22,-3,
224,-7,-8,-9,-80,-45,-7,-50-1,-7,-9,-62-7,-71,-4,-81,-2,
318,-14,-19,-25,-6.
, encloses Horses of Itabirite containing lines of auriferous
Jacotinga, 251,-65, 319. ———, limited vertical and horizontal range of rich beds, 224,-7,
230,-2-4,-62,-4-7,-9,-71,-82, 325.
, composition of, 214-15,-19,-21,-3,-7-9,-30,-3-4,-44,-5, 251,-4,-5,-6-7,-62,-4-75, 313-14,-24,-7.
, structure of, 214,-19,-21,-3,-7,-8,-80,-46,-9,-56,-9,-60,
264-9,-70, 318-19,-29-30.
distribution of gold in, 215,-16,-19,-23-7,-30,-8,-4,-42,
245,-7,-57,-62-4,-6,-71,-2,-4,-7, 319,-26,-30.
proportions of gold in, 230,-2,-5,-55,-79,-81, 325,-7,-31.
, nature, proportion, and distribu-
tion of alloy in, 215,-36,-86,
835,-7,-8.
, displacements of, by cross-veins, 226,-8,-86-7, 341.
, temperature of mines in, 728-31,-51,-6,-7,-8,-61; Tables
XXXII.—XXXVI.
Jamaica, on the copper-bearing granite of, 511.
James, Captain Richard, on submarine works at Wheal Margery, 599.
James, S. H., Esq., on submarine works at Botallack, 599.
, on the proportion of tin in the vein-stone of, , , 472.
, " proceeds and profits of " ,443; Table XIV.
-, , existence of eels in mine- water at } ,, ,717.
Jars, M. G., on the proportions of silver in Norwegian ores, 100.
English load area 107
, ,, English less ores, 107.

Jars, M. G., on the ores of Fahlun, 548.
Jeffery, Captain William, on the auriferous deposit of <i>Gongo Soco</i> , 255,-66,-70-3.
Jenkin, Alfred, Esq., on the proportions of metal in the silver-ores
of Herland, 111—12.
Jenkin, Wheal, on the tin-ore of, 677.
Jennings, Captain William, on the auriferous deposit of Gongo Soco,
255,—66, 271—4.
, , of Antonio Ps-
reira, 804.
, on a land-slip at Gongo Soco, 350.
Jewel, Wheal (Calstock), on the silver-ore of, 116.
Jewel, Wheal (Gwennap), alleged waste of earthy black copper-ore
, profits of, 451; Table XIV. [at, 694.
Jham, on the ferruginous conglomerate of, 89.
Jherria, on the ferruginous sandstone of, 64.
Johnson, P. N., Esq., on the proportions lead-ore of Devonshire,
of silver in the \
——————————————————————————————————————
, analysis of Carvoeira from Catta Preta, 240.
Jacotinga , Gongo Soco, 267.
auriferous native copper from
Duraö, 237.
, ,, zinc, lead, and iron ores from
Wicklow and Anglesea, 546.
, " silver-ores from Sark, 536.
, , gold from Morro Velho, 334,-74.
, " from Gongo Soco, 286.
, on the lode of Sark's Hope, 531.
, , lead-ore of ,, , 536.
, ,, proportion of gold in the vein-stone of
Grasty, 373.
, ,, ,, of palladium (Candonga, in the gold of) 175, 834.
, " of alloy " Gongo Soco,
286, 885,
887,-8.
, " of tellurium " Catta Branca,
886.

Johnson,	P. N., I	Esq., on the qualities of gold in different rocks, 236, 332—6.
		-, his present of palladium to the Geological
		Society of London, 336.
Johnstone	, R., E	sq., M.D., on the elevation of land at Caldera by
	•	by an earthquake, 157-8.
Johnston.	J. W.	F., Esq., on the Geology of New Brunswick, 490.
		, ,, sandstones of ,, , 501.
		, " slate of the Tattagouche, 495.
Joints, di	rections	s of, in granite, 3, 841,-85, 492, 511, 678-4,-80-1.
	,,	, ,, gneiss, 520,—6.
		, ,, talcose, micaceous, and chloritic slates, 7, 12-
,	"	13, 15, 18, 26, 211,-98, 301,-2, 603,-37,
		651-2,-74.
		, ,, quartz-rocks, 80, 180.
	"	, ,, clay-slate, 26, 182-3,-6,-7,-8,-90,-1,-3,-5,
,	"	207, 317,-21,-3, 500-1,-59,-75, 674, 703;
		Tables II. XXIV. XXVII. XXVIII.
		, " porphyry, 651—2.
	"	, ,, hornblendic and felspathic rocks, 161,-4, 405,
,	"	648,-71.
		& limestance
,	"	which alternate with them, 74, 84,-8, 124.
		, " carboniferous slate, 603,—14; Tables XX.
	"	limentana C14 [VVI
	"	, ,, the Coal-measures, 503,—6.
-	"	, ,, New Red Sandstone, 515.
 ,	"	, ,, recent sandstone, 155.
	"	, ,, rocks generally, their influence on metallifer-
,	"	ous deposits, 474, 560, 603,-49.
		, ,, metalliferous deposits, 76, 84,-8, 181,-6,-7,
,	"	188,-91,-3, 207,-23,-5,-46,-50,-64, 421
		433,-67,-8,-80, 548,-51-8,-69,-97, 608,
		605,-80-1, 712; Tables III. VI. VIII.
		X. XVII. XVIII. XX. XXI. XXIII. XXIV.
		, ,, metalliferous de-) native copper, 433,-69,-80;
	,,	posits containing Table XIII.
	22	mold 1810 998
,	77	7, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,
		VIII. X.
 ,	••	, ,, ,, silver, 76; Table
,	"	, ,, , , , , , , , , , , , , , , , , ,
	K -2	 ·

Joints, dir	ection		alliferous its contair		f copper, 13, 187,-8, 207, 321,-8; Table X.
,	"	, ,,	,,	"	iron, 181, 225,—47, 264, 469.
 ,	"	, ,,	"	"	iron (auriferous), 187,—8, 207, 321, 323; Tables VI. X.
 , ·	"	, ,,	,,	29	molybdenum, 654.
 ,	"	, ,,	"	"	silver, 76, 88.
,	"	, ,,	"	fragmen	tary matter, 75, 225, 247.
 ,	,,	, ,,	"	clay, 18	, 182, 559, 648,–83.
 ,	,,	, " cro	ss-veins,	124, 683.	
 ,	"	, groov	es (striæ), on their	opposite sides, 433,
				469, 552, 6	48,–83; Table VI.
 ,	"	, displa	cement o	f metallifer	ous deposits by, 183,
——,	"	, air ei	nitted by	, 2 50.	[301, 474, 559.
——,	,,	, inflar	nmable g	as emitted b	y, 195-6; Table VI.
José, San	(Cop	iapò), copp	er-mining	g at, 164-5.	
José, San	(Cha	narcillo),	extent ar	nd produce	of works at, 124; Tables IV. XIV.
Juan, Sa	n, on	the ores of	copper i	n sandstone	at, 514.
		Isq., on the		er- \ Cork a	nd Kerry, 602-3,-11- 12.
		•	e carbonif imestones	- 5	,, 612–14.
			e carbonii ous shale		,, , 613.
		— , on the	e lodes of	Ardtully,	ľable XXI.
					it Morro Velho, 199.
				the Caradon cts, on the, 6	, Camborne, Helston, 556-60.
					on-wire ropes at, 143.
		the native			• •
				ure in, 752.	
Tala Das			_	-£ 01	

Kala Bun, on the calcareous rocks of, 31.

Kaleedoongee, on the detrital deposit of, 45.

Katharinenburg, on alloys in the gold of, 335.

Kempston, Nicholas, Esq., on the black copper-ore of *Connorree*, 565.

Kempston, Nicholas, Es	sq., on the use of a beechen
	pranger as j
	-, ,, wooden linings for ,, 570.
	ene bumbs at)
	profits made in Wheal Vor, 446.
Kenmare, mining distr	ict near, on the rocks of the, 611-15.
, "	, " copper and lead lodes of the, 616-21.
, "	, ,, cross-vein of the, 622.
 , ,,	, subterranean temperature observed
	in the, 742-4.
Keweenaw Point, on th	e configuration of, 394.
 , "	rocks of, 394-404.
 , ,,	lodes of, 405-89.
, "	native copper of, $421-2, -5-36, -62-3$; Tables XII. XIII.
 , ,,	native silver of, 436.
	cross-vein (Slide) at, 439.
, subte	erranean temperature observed at, 738-4.
, on th	e erection of a Man-engine at the Cliff mine,
	439-40.
 , "	produce and profits of mines at, 440,-59;
	Table XIV.
 , ,,	antiquities discovered at, 412-19.
Keyne, Saint, temperat	ure of water in the well at, 749.
Keyserling, Count A. v	on, on the gold-mine of Berezovsk, 177.
	-, ,, detrital gold and pla- 175, 340; tina of the Ural, Table XXII.
	-, ,, ,, parent rock of, 843.
	-, ,, copper-ore of the Coal-measures in the Ural, 508.
Khetsaree, on the iron-	mines, furnaces, and miners of, 21,-9,-57;
Khuloagarh, on the iro	n-ores of, 33,—5. [Table II.
Khurna, on the iron-or	es of, 33,—6.
Killarney, magnetic dec	lination at, 616.
Kinahan, G. H., Esq., o	on the chloritic minerals of Cork, 602.
Klaproth, M. M. H., o	n the argentiferous lead-ores of Cornwall,
Knockmahon, on the ro	cks of, 593, 601, 738. [118,-19,
	ganic remains at, 594—5.
	des of, 594-8, 739; Table XIX.
	088-veins of, 598.

Knockmahon, on the submarine works at, 599-600.
, ,, subterranean temperature at, 740.
, " produce, costs, and profit at, 442, 601; Table XIV.
Knott, Capt. William, on the proportions of metal in the silver-ores
of the East Cornwall Silver Mines, Wheal
Duchy, Wheal Brothers, Wheal Sisters,
Wheal Langford, and Wheal Mexico
(Calstock), 115—16.
Kokcharoff, M., on a mass of detrital gold discovered at Zarevo-
Alexandrofsky, Table XXII.
Kotelar, on the iron-ores and furnaces of, 29, 57.
Kumson, on the rocks of, 1-4.
, ,, copper-mines of, 4—18.
, ,, iron- ,, ,18-42.
———, " mining-works of, 48—53.
, " smelting- " ", 53—62.
, ,, Royalties paid in, 62-8.
———, ,, detrital deposits of, 42—6.
, ,, ,, streams disappear in, and reappear
from beneath, them, 44.
———, on the forests of, 21.
Kurn Pryag, gold in the Aluknunda at, 4.
Kurrye, on the rocks and copper-ores of, 4.
Kyroolee, on the iron-ores of, 33,—7.
Lac la Belle, subterranean temperature at, 733.
Lake Superior, temperature of, 734.
, navigation of, 434.
, subterranean temperature in the neighbourhood of,
, on ancient copper-mining near, 412-19. [738-4.
Lamborn, — Esq., on the manufacture of iron near Lake Superior, 390.
Lamellar structure common to rocks and lodes, 85,-8; Table III.
Lamps used by labourers (apires) in Chili, 147.
Samuel managed by growledge in the Himpleys 10
Lan. M. —, on the proportion of silver)
Lan, M. —, on the proportion of silver afforded by Spanish lead-ores, 103.
, " French " , 104.
Land, price of, near Lake Superior, 390.
,, in Cornwall, enhanced by the discovery of copper-ore

le

7.


s

ı

```
Landslips, frequent at Gongo Soco during the rainy season, 350.
Langford, Wheal, proportions of silver in the ores of, 115,-20.
Lanivet, proportion of tin-ore in the vein-stones of, 472.
Lenreath, on the Mining district of, 699-720.
       -, subterranean temperature of, 748-9.
Lapham, J. A., Esq., on the ancient copper-miners of Lake Superior,
Larraona, on the early discovery of copper-ore at, 154.
Laur, M. —, on the auriferous deposits of California, 838—9.
       <del>---</del>,
                     detrital gold ..... of
               22
                                                      , 343.
Laws, Mining, of Brazil, 861-5.
               of Chili, 148.
Lead, directions, inclinations, widths, and ingredients of deposits
        containing,-
   —, ores of, in gneiss, 517-27.
             , in quartz-rocks, 643-5.
             , in clay-slate, 699-720.
             , in hornblendic and felspathic rocks, 582-8, 701-2.
            , in carboniferous limestone, 610-21.
             , associated with gold, in Australia, 99.
                                   , in Brazil, 334,-75.
         "
                                   , in Virginia, 376,-7.
         22
                                   , in Wales, 640.
         ,,
             , associated with silver, in Australia, 99.
         ,,
                                    , in Bohemia, 101.
         "
                        "
                                    , in Chili, Table III.
         "
                        ,,
                                    , in Cornwall, 111-12,-15-21.
         "
                        27
                                                          707-14.
                                    , in Cumberland, 108-9.
                        ,,
                                    , in Derbyshire, 109.
                        99
                                    , in Devonshire, 109-10.
                        ,,
                                    , in Durham, 107-8.
                        "
                                    , in France, 103-5, 526,-37.
                        ,,
                                    , in Hanover, 101.
                                    , in Hungary, 100-1.
                                    , in Ireland, 105, 546,-8,-9, 621.
                                    , in the Isle of Man, 105.
                                    , in Northumberland, 107-8.
                                    , in Norway, 100.
                                    , in Sark, 121, 535-7.
                        ,,
                                    , in Scotland, 106, 643-5.
                                    , in Shropshire, 109.
                        ,,
```

Lead, ores of, associated with si	lver, in Sicily	and Calabria, 102,
 , ,, ,	, in Spain,	
		nited States, 99.
		, 106-7, 546,-76.
		by them at different
depths, 708-10		•
		d in them, 121, 710.
Lead-mines, subterranean temp		5,-8; Tables XXXIV.
T 35 (70)	.•	XXXV.
Lean, Messrs. Thomas and Joel,		-
Leaps (See DisplacementsVer		
Leitao, M., on the proportion of		
Lelivec, M., on the proportions ores, 103.	of silver affo	rded by French lead-
Le Play, M. F., on the detrital	gold of Siberi	a. Table XXII.
Lettsom, W. G., Esq. (& Gregg		
Levant, on the native silver of,		Cornwall, 116,-18.
, ,, extent of subma		
, ,, profits at, 442;		, 000
Levels and Winzes, when and		stituted for Stones in
	of Cornwall,	
Levy, Armand, Esq., on the iro		
		mines of Brazil, 369.
Liefchild, J. R., Esq., on the inc		
Lignite, in the copper-bearing		
 , ,, ,,		Siberia, 508–9.
 , ,, ,,	Coal-measures	of New Brunswick,
		506–10.
, , , , ,		of Nova Scotia, 507-8.
Lime, an ingredient of sea-sand	•	
, its use as a precipitant of		
Limestones of Brazil, 245,-9,-9		
,, Chili, 71–9, 82,-		•
	, 12-16, 18.	
	,-18-21,-2-6.	
, compositi	ion and struct	ure of, 71-9, 82,-5,-6, 89, 92, 304.
, siliceous,	"	, 5, 12—16, 18,
,,	,,	78, 245,—9,
		304,-13,-19.
, magnesian,	,,	, 85, 297-8, 304.

Limestones, carboniferous, composition & structu	
, sprinkled and veined with copper-ore	
, " silver and	silver-ore, 76 — 7,
, lodes in, 80-124, 616-21,-23-6.	[89-90.
	_
, subterranean temperature in, 724-5,	-42-4.
Lingula-flags, auriferous deposits in the, 635-42.	
Lisburne mines, on the returns, expenditure, and	profit of the, 459-
Liskeard, mining-districts near, 650-720,-44-9.	[460.
Little Bounds, extent of submarine works at, 600	-
Llama, unsuccessful attempts to naturalize the, in	Chili, 141.
Lloyd, The Rev. C. A. A., on the climate of Shre	
Lloyd, Colonel, on the mines of Chaffarcillo, 148.	
Lloyd, The Rev. Dr., on the climate of various	
Magnetic)	
declination } at ,,	, 541,- 75.
Lodes. (See Metalliferous deposits.)	
Logan, Sir W. E., on the sandstones of Lake Su	perior, 893.
, " of New Bru	ınswick, 502-3.
, ,, fossil flora of ,,	, 505.
Lonsdale, William, Esq., on the fossil flora of Ne	•
Lory, M. Ch., on the rocks of Chalanches, 519.	, , , , , , , , , , , , , , , , , , , ,
Loss of gold during the processes of reduction at A	forro Velho, 204-5.
Luc, J. A. De, Esq., on alternations of granite a	
kitchen, 658.	
Luçencia, on the precipitation of copper at, 590,	-2 .
Ludcott, Wheal, on the proportions of silver and	
of, 120, 710-11.	
madisa made and lass	at 719
Luke, Capt. John, on the auriferous deposits of	
	Fongo Soco, 278.
Lusghanee, on the iron-ores of, 22.	aongo soco, 210.
Luttea Garh, on the copper-ores of, 20.	
Lushington, G. T., Esq., on native-mining in the	Timelers 51
	umanaya, or.
————————, on the wages of native a	,, , 61.
, on experimental works	,,,, by a
in the)	Cornish Superin- tendent, 52.
on the use of pine-torches	in the Himalava, 52



Man, Isle of, proportion of silver in the lead-ore of, 105. Man-engines, notices of, 439, 609.
Manganese, ores of, in Bengal, 65.
, ,, France, 527.
, " the Himalaya, Table I.
, " New Brunswick, 496.
Manto de Agua, an impermeable bed of clayey limestone, 77.
Manto de Cachi, a thin bed of calcareous spar, 79.
Manto de Ossa, dimensions, composition, and structure of, 75, 82.
, displacement of by flucans, 125.
Marazion district, subtorranean temperature in the, 752.
Marchand & Scheerer, MM., on the specific gravity of copper after
compression, 430.
Margery, Wheal, extent of underground works at, 599.
Marianna, present poverty of the gold-mines near, 368.
Marke Valley, on the rocks of, 656,-9,-61,-7-8,-71,-3,-4.
in confronting (walls) sides of the lode
659-60.
, ,, lodes of, 676,-7,-9; Table XXVI.
, ,, the population employed, machinery in use at, and
produce of, 698.
, ,, subterranean temperature at, 746.
Marmato, diminution of springs consequent on clearing the surface at, 137.
, subterranean temperature at, 770.
Marrot, M., proportion of silver in the lead-ore of France, 104.
Martius, Dr. C. F. P. von, on the mountains of Minas 178.
& Spix, Dr. J. B. von,) Geraes,)
, ,, vegetation ,, , 169.
, rocks of , 174, 210,-11, 212, 304.
, , minerals of , , 174, 305-6.
, ,, gold-mines of ,, ,180,—4—5,
222-8,-88-9, 243, 808.
, , quality of } , , 213.
, iron smelt- works of , 212-13.

Martyn, Thomas, Es	q., on	the silver-ore of Wheal Basset, 113.
		clay-slates and greenstones of, 700-3.
,	"	direction, dip, width, and composition of the lode at, 704-14.
	"	brecciated and concretionary structure prevalent in portions of, 712-14.
,	"	(seemingly) isolated masses of vein-stone
		which represent it, 714.
 ,	"	proportions of silver extracted from the lead-ores of,
,	"	,, , obtained at different depths in, 709.
	,,	cross-veins (flucan) at, 715-16.
	"	", displacements of
,	"	the lode by, 716-17.
 ,	"	sulphate of lime contained in the water at, 717.
 ,	,,	subterranean temperature at, 748;
		Tables XXXIV. XXXV.
ф,	"	rate at which the works have been deepened, 748.
,	,,	population employed at, machinery in use
		at, produce and profits of, 718-20.
Mary, Wheal, propo	rtion	of tin-ore in the vein-stone of, 472.
Masses of native cop	per, 4	127,-8,-9,-34-5,-62,-70-1,-5-7,-81,-5; Tables XII., XIII.
	er, 91	,-2,-3,-123,-4, 436-7, 521; Tables III., IV.
,, gol	d, 21	5,-16,-33,-72,-3,-7,-9,-80,-3,-303;
	·	Tables VIII. XXII.
Mason, James, Esq.	(Baro	on de Pomeron), on the petroleum of Huidobro, 516.
Maton, W. G., Esq.,	on th	e profit made at Carnon-Stream, 452.
		produce and profits of West Wheal Seton, 449.
Matthews J Esq.	n the	proportions of tin-ore in the vein-stones
mucono, o, modi,	,	of different mines, 472.
	,,	produce, costs, and profits of mines near
,	"	Tavistock, 458,—9.
	,,	comparative values of barrows and rail-
,	,,	ways in mines, 144.


Matthews, J., Esq., on the	(alleged) waste of copper-ore near Tavistock, 693.
Watthew C. The service	•
	quality of Californian gold, Table XXII.
Matrix. (See Composition	
	paz and euclase of Brazil, 300,-5.
	ld-mines of ,, , 243, 351.
, ,, de	trital gold of ,, , 221, 301,-42.
, ,, ex	portation of gold by smugglers, 365.
Mayen, Dr., on the valley	of Copiapo, 140.
Meadow-mine, on the rock	
Mehelchowree, on the iron	
Mendips, on fossiliferous l	
	of, in use at Morro Velho, Table VII.
	lock of Walls TITT
 , , , ,,	, lost at ,, , 1 able v11.; 334,-74.
	, lost in California, Table XXII.
Merionethshire, on the au	
	pper turbaries of, 580.
	ntiful at small depths, 97, 121.2.
Wateling Color	
Metales frios, ,,	at great ,, , 97, 121-2.
	nation granite, 175, 811,-20, 511-12, rough 664.
, ,	talcose rocks, 177, 210,-99, 801, 302,-3,-20, 495.
·, ,	, porphyry, traversing mica-slate, 654.
 ,	, clay-slate, 175,-86,-96,-321.
	, calcareo-siliceous rocks, 249, 804,
	823.
, , , , , , , , , , , , , , , , , , ,	, Itabirite (Jacotinga), 219,—42, 249,-55.
, , ,	, calcareous, alternating with horn-
	blendic, rocks, 76-7, 89-90.
 , ,	anal-shalos 509-10
, , ,	conditions 519 15
	transpoor complements ATE
diggemin	nation of copper and the ores of copper,
, uioseilii	236, 475,-95, 503-10,-11-12.
, , ,	, gold, 175,-7,-86,-96, 210-19,-86, 249,-55,-6,-8,-99, 300,-1,-2, 303,-4,-11,-24,-75.

Metallic minerals, disse	mination of n	nolybdenite, 654.
 ,		ilver and the ores of silver, 76-
		77, 89-90.
,		in-ore, 175, 664.
		ions, inclinations, and widths—
;	in Brazil,	181,-2,-3,-4,-8,-9,-90,* 211,-14,
	216,-17,-1	9,-21,-3,-7,-9,-30-5,-6,-40,-2,-5,
	247,-9,-58	3,-9,-98-300,-1,-2,-3,-9,-10,-11,
	375; Tabl	les VI. VIII. X.
 ;	in the Chan	nel Islands, 531,—2,—3, 737;
	Table XV	и.
;	in Chili, 71	, 81,-3, 91-3, 133,-61-2,-4-5;
	7	Table III.
 ;	", div	ergent directions of, 81.
		(Cornwall, 410, 674-5, 704-5
;	in England;	Tables XXIII.—XXIX.
	* . 101	Shropshire, 515.
;		20—1; Table XVI.
 ;		livergent bearings of, 520,-6.
;		alaya, 4, 5, 7, 9, 11, 12, 14, 15,
	Tables I. I	4,-5,-6,-8, 30,-3,-5,-6,-7,-9, 41;
 ;		542-4,-63-4,-96, 603-4,-16, ables XVIII.—XXI.
	in Scotland,	
	in Spain, 51	
,	m opan, or	near Lake Superior,
		405,-6,-8,-9,-10,
		411, -23, -35, -60,
;	in the Unite	ed States, $\langle 463, -4, -5, -7-8, $
		479,—80; Tables XII. XIII. XV.
		All, Alll, AV.
		in Virginia, 872,-5, 376,-9,-80.
	in Wales, 57	
		ores of antimony, 520-1,-32,-42-
,	,	545,-63-4, 616; Tables
		X. XVII.
	,,	, of bismuth, 372, 542-4, 637-
,	,,	638 ; Table X.
) ,	of chromium, 647-9.
<u> </u>		,, of cobalt, 116, 520-1; Table
•	•	XVI.

Matallifarous deposits violdi	ng copper and its ores, 4,-5,-7,
mountainerous deposits, yieldi	9, 11, 12, 14, 15, 18, 161-
	162,-4-5,-88-90, 236,
	816, 405, -6, -8, -9, -10, -11,
	423, -85, -60, -8, -4, -5, -7,
	468,-79-80, 513-14,-15,
	542-4,-63-4,-75,-6,-96, 608-4,-16,-74-5; Tables
	VI, XII, XIII, XV. XXIII.
	XXIV. XXV. XXVI.
	gold, 181,-2,-3,-8-9,-90,
	211,-14,-16,-19,-21,-3,
	227,-9,-30-5,-6,-40,-2,
	245,-9,-54,-9,-64,-6,-7,
	298, 300,-1,-2,-3,-9,-10,
	811,-72,-5,-7,-9,-80, 687- 638; Tables VI. VIII. X.
	the ores of iron, 19, 20,-4,-5,-6,-8,
, ,,	30,-3,-5,-6,-7,-9, 41, 188-
	189, 211,-14,-16,-19,-21,
	223,-7,-9,-85,-6,-42,-5,
	247,-9,-53-4,-9,-64,-6,-7,
	298, 300,-9,-89-90, 542-
	544,-63-4, 647; Tables I. II.
	of load 500 1 99 690 AR
, ,,,	" of lead, 520-1,-82, 620,-48, 704-5; Tables XXVII
	XXIX.
	,, of molybdenum, 650,-4.
	" of nickel, 520-1.
, "	" of palladium, 214, - 16, - 49,
 , "	,, of platina, 249,-58-4,-9,-64,
	266.
· ,,	" of silver, 71, 81-8, 91, 188,
	188,-9,-90, 227,-9,-35,
	249,-53-4, 809-10,-72, 423, 520-1,-32,-42-4,
	616,-37-8, 704-5; Tables
	III. VI. VIII. X. XVI.
	XVII. XVIII. XXVII.
	XXVIII. XXIX.
, , ,,	" of tellurium, 376,-9-80.
······································	" of tin, 674-5; Tables XXV.
	XXVI.

Metalliferous deposits,	yielding the ores of zinc, 520-1,-63,-77, 620,
	637-8,-43,-704.
·	in granite, 674-5; Tables XXIII.—XXVI.
	in gneiss, 520–1.
	in talcose, micaceous, and chloritic rocks, 4,
•	5, 7, 9, 11, 19, 20,-2,-8,-6, 177, 211,-98,
	300,-1,-2,-8,-9,-10,-11,-72,-5,-7,-9-80,
	608-4,-52,-74-5; Tables I. X. XXV. XXVI.
	in quartz-rocks, 181; Table X.
	in clay-slate, 5, 25,-6,-8,-31, 182,-3,-8,-9,
,	190, 309,-10,-11,-72,-7, 542-4,-63-4,-75,
	596, 674-5; Tables II. X. XXV.—XXIX.
	in calcareous slate, 4, 12, 35,-6,-7.
	in Itabirite (Jacotinga), 214,-16,-19,-21,-8,
	227,-9-85,-42,-5,-9,-58-4,-9,-63,-4,-6,-7,
	298, 309-11; Tables VIII. X.
	in Hornblendic rocks, 405-10,-11,-20,-3,-60,
,	463,-4,-5,-7-8,-79,-80-1, 532, 647, 704-5;
	Tables XII. XIII. XV. XVII. XXVII.
	XXVIII.
	in Hornblendic rocks alternating with lime-
,	stones, 71, 81-3, 91, 183; Table III.
	in Carboniferous slates, 603-4,-16; Table
,	XX.
	in ,, limestone, 618, -19, -20, 623; Table XXI.
	in New Red Sandstone, 515.
,	bounded by different rocks on opposite (walls)
	sides, 616; Table XXI.
,	adjoining which granite occurs nearer the surface in the (hanging) upper, than in the (foot wall) lower, side, 658.
,	· (foot) lower, than
•	in the (hanging wall) upper, side,
	657-60; Tables XXV. XXVI.
	at the junctions of different rocks, sometimes
•	course awhile between them, before passing
•	from one into the other, 81-2, 616,-57-60,
•	701.

M + 11'6 1'41'-1 6	
Metalliferous deposits, which conform	
	8 9, 11, 19, 20, -1, -5, -6, -8, -30
	,-41,-82, 161,-77,-80,-2,-3,-4,
	14,-16,-19,-21,-3,-7,-9,-30,-6,
	59,-66,-7,-98, 300,-1,-2,-3,-10,
	463,-4,-5,-7,-79,-80, 542,-53,
554,-68,-75	, 602,-20,-37,-47,-52, 787;
Tables II. V	I. VIII. X. XV. XVIII. XXI.
, which conform	n to the cleavage of schistose
	rocks in direction, but are
	more highly inclined, 564,
	604; Tables XVIII. XX.
,, ,,	to the joints of rocks, 14, 15,
. "	22, 74, 81,-5, 164,-83,-7-
•	188, 223,-64,-99, 301,-2,
	815, 603,—18,—24,—74;
	Tables I. III. VI. VIII. X.
	XXIII.—XXVI.
	to the cleavage in some, but
 , "	with the joints in other,
	places, 187-8, 223,-64,-98-
	299, 301,—2; Tables VI.
	VIII. X.
, "	to neither cleavage nor joint,
	81, 405,-9,-10,-11,-22,-60,
	515,-20-1,-32,-54,-64,-96,
	602,-4, 704; Tables XII.
	XIII. XVI. XVII., XIX.
	XX. XXVIIIX.
· , ,,	to the joints, sometimes dip
	to the cleavage; - yet sometimes
dip towards	neighbouring bodies of granite;
674-5.	
, bodies (shoots) of metals, ores, and vein-stones
	ich conform, in endlong dip, to
the (ripples	s) flexures of slates, to the in-
clinations of	f different strata on the opposite
sides of joi	nts, and to the slope of neigh-
	nite, 26, 32, 122, 207,-15,-16,
),-63,-4,-6,-7,-71, 323,-6, 439,
	7,-78,-94,-7,-8, 608.
,,	, , , , , , , , , , , , , , , , , , , ,



Metalliferous deposits, often enclose (horses) masses of the contiguous rocks, 20, 84, 181,-90,-1,-4, 251,-65, 312,-17,-19,-73,-8,-81, 420,-4,-9,-61,-8, 480, 551,-5,-64,-8,-76,-96, 605,-24,-48, 712.
(Country) formations, and the included (horses) masses of rock, 190,-1,-3-4, 251, 265, 313,-18-19,-78,-81, 424-5,-61, 551, 564, 605-6,-24,-48, 712.
(See Composition.)
, bounded by joints, 85, 161,-4,-81,-7, 207,-28, 264,-99, 801,-2, 464,-7, 515, 604,-16,-18, 620,-49,-80, 714.
, ,, obliquely, by joints, 76, 181, 302, 468,-74,-80, 575.
, ,, transversely, by joints, 181,-7, 299, 480, 559,-78.
, , filled and clay, 13, 89, 182, 469, faced with 559.
metals and ores, 13, 39, 76, 181,-2, 223,-5,-8, 246,-64,-9,-80,-1,-3, 654.
vergently) scored, when faced with metals, ores, and clay, 13, 207, 378, 433,-69, 552, 648,-54.
-, laminated structure common to them and to the contiguous rocks, 85.

Metalliferous deponits	, of schisto		214,-23,-8,-9,-87,-46 254,-7,-9,-62,-5,-9 269,-98, 800,-1,-2, 811,-14,-18,-19, 551-2,-8,-64,-8-9, 575.
	9:	, , ,	often display (divergent- ly) striated faces of unctuous clay, 552.
	concretion	ary structure	e of, 182,-4, 318.
 ,			, 77, 89, 195, 225, 312,
			525,-81, 707,-18.
 ,	various	metals and o	them, sprinkled with res, 77, 90, 195, 225, 515, 625,-81.
,	relations b		in clay-slate, 199— 205, 829,-31;
,	• '	" " "	in Jacotinga, 232—
			4-7,-9,-31; Table X. hardness and their pro-
,			er), at different depths,
	in horn	olendic (trap) rocks, 326.
,	proportion	s of native	copper in, 421,—34—6, 470-1.
,	"	,,	, at different
			depths, 421,-36.
 ,	**	of copper-c	ore in, 693.
,	"	of gold in,	100,-1,-77,-9,-82,-4, 186,-96-8, 201-5,-8,
			230-2,-5,-7,-49,-55,
			258,-79-81, 312,-20,
			821 ,-2,-4,-5,-7,-81,
			834,-5,-78-4,-5,-7,
			878, — 9, — 88, 550, 568, 641; Tables
			VII. VIII. X. XXII.
	"	",	at different depths,
,	,,	,, ,	177,-9, 201-5,-84,
			831,-82;
			Tables VII. VIII.
			X. XX.
 ,	"	of lead-ore	in, 693.

Matallifornous density manualism of the sea 2 700 00 101 Kg
Metalliferous deposits, proportions of silver and \ 76, 90-121,-58, silver-ore in, \ 206,-36,-86, 334-835,-8,-9,-40,-73-874,-8, 437-8, 522-4,-35-6,-45, 546,-9,-66,-8, 617,-32,-42, 708-11; Tables VII. X. XXII.
, , at different depths, 206,–86, 338–9, 708—9; Tables VII. X. XXII.
, of tin-ore in, 472-3, 693.
-, intersections and) by other metalliferous dedisplacements of, posits, 92, 127,-8, 493, 538.
,, by cross-veins, 127,—9, 225,-87, 841, 489,-74, 503,-15,-27,-58,-75,-98- 599, 608,-22,-44,-83-5, 716-17.
, by rocks, 176,-92-4, 818.
——————————————————————————————————————
, subterranean temperature in and near, 728-84.
, issue of inflammable gas from crevices in, 195-6.
, " in Banca, 631.
, in Brazil, 283, 842-70.
, " in California, 343; Table
, ,, in Canada, 384. [XXII.
, " in Chili, 845,-60.
in India 2 4 46 7
, ,, in Ireland, 627-84.
in Now Cranada 260
, , in Nova Scotia, Table
XXII.
, in the Rhine, 348,-51-2, \$56,-8,-60; Table XXII.

Metalliferous deposits	(Detrital), o	f gold	in Siberia, 175, 342,-3,-6, 348,-60; Table XXII.
	99	,,	in the United States, 384.
,	"	"	, Ancient detritus, 283, 342-5,-6,-8.
 ,	,,	"	, proportions of metal ex- tracted from, 346,—8. 358; Table XXII.
	, ,,	"	, quality of metal extracted superior to that of mine- gold, 359-60.
	"	"	, crystals of metal extract- ed from, 856,-9, 632.
	, ,,	"	, traced to the parent formations, 175, 343,-59-60.
	, ,,	,,	, cost of extraction, 846,-8, 358; Table XXII.
	, (Detrital) o	f gold,	46, 346-8,-51-2,-9.
	Recent de	etritus,	} ±0, 0±0-0,-01 2, 0.
	, ,,		, crystals of gold in, 356,-8.
)))		, shaped implements and
			ornaments mixed with, 844-5.
	, ,,		, an ancient Indian village imbedded in, 384.
	, ,,		, grains of gold swallowed
			by water-fowl, 356.
	(Detrital)	of tin-	ore, in Australia, 286.
	· , , ,	,,	, in Banca, 631.
	, ,, ,	,,	, in Cornwall, 452-3, 695.
	, ,, ,	,,	, in Ireland, 629-31.
	· , ,	"	, mixed with gold, 286, 629-31.
	-, ,, ,	"	, associated with platina, 286.
	-, ,, ,	"	, superior to mine-tin-ore in quality, 359.
	-, ,, ,	"	, profit and loss on the extraction of, 452-3; Table XIV.
	-, ",	"	, issue of inflammable gas from ancient vegetable matter overlying it,
			453.

Meuse, the stream of the, affected by the condition of the woods at
its source, 188.
Mexico, proportion of metal in the silver-ores of, 98,-9.
———, weight of ore brought out of the mines in, by labourers, 147.
Mexico, Wheal (Calstock), proportions of metal in the silver-ores of, 115.
Mexico, Wheal (Cubert), on the native silver and silver-ore of, 110.
Mica, an ingredient of the rocks, 2, 8, 4, 29, 30, 64, 171,-2,-4,-7,
210,-11, 376,-9,-86,-7,-92, 491, 508,
511, 613,-43,-50,-62,-3,-7,-70.
, ,, metalliferous deposits, 511,-24, 675,
Michell, Edmund, Esq., on the profit made at Carnon-Stream, 452.
Michell, Edward, Esq., on the proportions of metal in the silver-ores
of Herland, 112.
" ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' '
Menheniot, 708-9.
, on the produce and profits of East Wheal
Rose, 454.
Michell, Mr. Francis (of Redruth), on the introduction of the plunger-pole at Ale and Cakes, 570.
Michell, Capt. H., his suggestion of the use of mine-water for pre-
venting dry-rot, 570.
Michell, James, Esq., on the proportions of silver in Spanish lead-
ores, 102.
Michell, John, Esq., on the native silver and silver-ores of Cornwall and Devon, 112,-16,-18.
, ,, argentiferous lead-ore of Wheal Rose, 119.
discovery of copper-ore in Cornwall, 689,-91.
وسواسته والمنس المعانونين وبالمواب
The state of the s
from Australia, 286.
Miguel, Corrego de Saő, on the metalliferous deposit of, 242.
, " iron-smelting works of, 242.
, ,, discovery of diamonds at, 242.
Miller, Major W., on the detrital gold of Virginia, 384.
Miller, Dr. W. A., on crystallization from saturated solutions, 713.
, analysis of mine-water from the Clifford mines,
Mine-gold, its inferiority to detrital gold, 359,-60. [586.

Mine-tin-ore, its inferiority to detrital tin-ore, 359, 472.
Miners, in Australia, earnings of, on quartz-formations, Table XXII.
, ,, , on detrital deposits, Table XXII.
, in Brazil, European (in mines worked by British Companies)
earnings of, Table
XXII.
, ", Native (", by Brazilian proprietors)
modes of labour
217-18,-21-2,-99.
, ,, ,, (in the beds of rivers in search of de-
trital gold), modes of labour and
earnings of, 356-8.
, ,, Slave, (in mines wrought by both native and
foreign proprietors), modes of
labour and earnings of, 217-18
221-2,-92,-3,-9, 301.
, ,, , (in mines wrought by both native and
foreign proprietors), hire of, 382
, in Chili, European (in mines owned by native proprietors)
Table V.
, , Native (in mines owned by native proprietors)
nature of labour, 146-7,-51-2.
, ", " (in mines owned by native proprietors)
amount of earnings, Table V.
, ,, ,, reckless character of, 91, 167.
, ,, ,, their dresses, and persons examined or
leaving the mines, 91.
, in France, Native (Orpailleurs), searchers for detrital gold
amount of earnings, 358
, in the Himalaya, Native (on their own account, but paying
a Royalty to the Government), nature of labour and
amount of earnings, 58-61.
, in Ireland, Native (County of Wicklow), amount of earnings
, in Nova Scotia, European, earnings of, Table XXII. [562]
, in Siberia, Native (searchers for detrital gold), amount of
earnings, 848.
, in the United States, Slave (Virginia), hire of, 382.
, , , European (Lake Superior), amount of
earnings, 428,-71.
, dangers to which they are sometimes exposed, 600,-66.
from Derbyshire and from Wales employed in the mines of
Devonshire, 109.

Mining, early system of	of, in Germany,	646.
, ,, mode of,	in Cornwall, 14	6, 448, 645-6.
	in Anglesea, 57	
	er in Cornwall, 6	
	in Chili, 154.	
	•	i- Commoll 146 645 7
		m in Cornwall, 146, 645-7.
, substitution of	r tram-waggons	for wheel-barrows in, 144.
, present modes		to the surface in, 142-4, 609,
, "	of ventilation in	n, 219-20. [668-9.
	of, and implement used by na workmen.	in Brazil, 217-18, 221- tive 222,-99.
, ,,,		, in Chili, 142-8.
		, in the Himalaya, 48-52.
, ,,	of prince fire for	or expediting operations in,
, ancient, in (th		ersetshire, 625. [51-2.
	Lake Superior, 4	
Mining-works, rate at	which they have been deepened,	in Brazil, Tables VII. IX.
 ,		in Cornwall, 448, 748,-64-5.
 ,		near Lake Superior, 439-40.
, submar		
		rom water entering, 562,-9-
, precipio	amon or copper r	592; Table XVIII.
	of fal in streams	
		issuing from, 854-5, 717,-31.
, vegetati	ion in	,, , 585,-8, 686.
, produce	and profits of, 9	05-6, 123-4,-79, 208,-16,-17,
2 85,-	47,-7 4-7,- 89, 38	3 , 44 0-59, 601,-41,-4,-94-5,
698, 7	720 ; Tables IV.	. VII. IX. XIV. XXII.
Mining-laws, 148, 861	-5.	
Minnesota mine, on the	e rocks of, 475.	
 , ,,	lodes of, 475-6	•
		conglomerate of, 476-7.
 ; ",		ofits of, 459; Table XIV.
		rous ore from Grasty, 874.
Mohs, H. Frederick, or		
tomsenet, m. D., on th	Cornwall an	different ores in the lodes of d Devon, 693.
, ,,	use of tram-wa	aggons in mines, 669.
	produce and p	rofits of the Lisburne mines,
, ,	- •	460.


Molybdenum, ore of, in Breadalbane, 650-4.
, ,, in Leicestershire, 654.
, ,, , in Norway, 546.
, ,, in Australia, 654.
Mona mine, on the rocks, vein-stones, and ores of the, 574-8.
, ,, treatment to which pyritic ores are subjected at the, 578-9.
,, precipitation of copper from mine-water at the 581-4; Table XVIII.4
, ,, quantities of ochre obtained ,, , 588-4
, ,, of rain at the, 582.
Monk Wearmouth, subterranean temperature at, 753.
Monlevade, M. A. J. de, on the iron-ores and smelting-works of Monlevade, 171, 219.
differences in the qualities of char- coal from different woods, 219
Monte Amargo, on the recent sandstone of, 155.
Montgomery, Dr. J. B., analysis of water in the river of Copiapò, 140
, on the identity of diatoms, in the present
strand, and in the raised beaches,
near Caldera, 160.
Moore, C., Esq., on a fossiliferous lead-lode at Charter House Warren, 625,-6.
Morcom, M., Esq., on the rate of blasting vein-stones at Polberro, 200.
, , proportions of tin-ore in , , 202, 472.
Morgan, John, Esq., on the auriferous deposits of Gongo Soco, 272. ————, " " of Tacquaril, 273.
, ,, quantities of gold coined at Rio de Janeiro 369.
, ,, ,, used by goldsmiths at Rio de Janeiro, 870.
Morro das Almas, on the auriferous deposit of, 224:
Morro Velho, on the rocks of, 185-6.
, which occasionally contain gold, 186, 321.
, their schistose and jointed structure, 186,-91,-2,-8.
, ,, , narrow highly-inclined mass of joint- ed rock between two bodies of vein- stone, 191.

Morro Velho, on	the rocks of,	transitions between them and metal- liferous deposits, 192,-8,-4, 207, 813.
, the		their directions, coincide sometimes with the cleavage, sometimes with the joints, but often with neither, 187-8, 90, 207, 310; Tables VI. X.
,	99	; their dip and width, 188-92, 811; Tables V. X.
,	"	; bodies (shoots) of the productive portions in, coincident in endlong dip with structural dispositions of rocks, 206-7, 328,-6.
 ,	**	; comparative hardness of, at different depths, 199; Tables VII. XXII.
,	"	; ingredients, 190,-4,-6, 202, 812, earthy, 813,-17,-21-2; Table VI.
,·	,,	; ,, , isolated masses of slate imbedded in, 190, 194, 312,-17,-28.
ş	"	; ,, , portions conforming to the joints or oblique to the cleavage, par- tially or entirely severed, for short distances, by certain beds, 192,-8-4, 318.
,	,	; ,, uniformity of cleavage in the (Country) walls, imbedded masses, and severing beds, 198-4, 818.
,))	; ingredients, 194-7, 312,-18,-21- metallic, 322; Table VII.
;	"	; ,, , proportions of gold contained in, 197,-8, 204-5; Tables VI. ^a VII.
;	"	; ,, proportions of gold extracted from, 197, 198, 201,-8,-4,-5, 312; Tables VI. ^a VII. X. XXII.

Morro	Velho, th	e metalliferous deposits of;	} i	metallic,)	proportions of gold extracted from ores obtained at different depths, 201,—3,—5, 831; Tables VII. X. XXII.
	,	"	;			proportions of gold extracted from ores of different hardness, 199, 201,—3,—6; Tables VII. XXII.
	,	,,	;	"	1	comparison between the proportions of gold contained in and extracted from the vein-stone, 198, 204, 205; Tables VI.• VII.
	 ,	"	;	99		analysis of gold, 834, 874.
	 ,	n	;			proportions of silver alloyed with gold, at different depths, 206, 882; Table X.
	 ,	"	;	"		quality of gold ob- tained at different depths, 205,-6, 875.
	,	"	;		feı	quality of gold in dif- ent vein-stones, 875.
	 ,	"		man per o	da; e,	th of hole bored per y at different depths 199-200; Table VII.
	 ,	,,	-	blasted, 19	99	e extracted per hole; Table VII.
	 ,	**	-	weight of	gr	of ore extracted to inpowder used, 200.
	,	n	·	stone rejection (of VII. XXII	cte gh dre L	d proportions of vein- ed as not containing to repay the cost of ex- essing), 202; Tables
	 ,	"		quantities a stone stam	an ipe	d proportions of vein- ed, 202; Tables VII. XXII.
	 ,	,,	;	numbers, viciency of VII. XXII		ight, speed, and effi- stamp-heads, Tables

Morro Velho		over which passed, 208	ed hides, and of baize, the stamped-ore is to; Table VII.
· · · · ·	, ,		f amalgamable ore col- lected by each series of <i>hides</i> and baize, Table VII.
	, ,,	; " of	ore escaping the <i>kides</i> and baize, but afterwards collected and ground in <i>arrastres</i> , 208, 855; Table VII.
	vein-stone, exten	t of granulation	on by stamping, Table VII.
,	quantity of merci	ry used, and t	ime required for amal-
•	•		ting the stamped ore,
			e VII.
	, , ,,	lost during	amalgamation, Tables VII. XXII.
•	costs of (dressing	cleaning the	tamped ore, Table VI.ª
	, ,, , total, of ex	tracting gold a	nd making it ready for the market, 327.
	icano of inflamma	hlo man from	•
	stones, 195 ; 7	lable VI.	crevices in the vein-
	, climate of, 849,-	52, 726.	
			6-7,-56,-8,-9,-60,-1,-2.
	, force employed a		
	, wages of Europe		
	, produce, expend	iture, and pro	ofit at, 208-9; Tables VII. XIV. XXII.
	, extent of water- ants, 350; Ta		njuries done them by
			ritic ore in suspension,
Morro, Vend	la do, iron-furnace	a near the 247	·
			e, and profits at the
,,			ley mines, 882-8.
			of slaves in Virginia,
Monacotia.	the flourishes in	water which	yields a precipitate of
	copper. 585		Arrem a brombrenc or


Mouillelonge, subterranean temperature in the coal-mine of, 753.
Mulberry Hill, proportion of tin-ore in the vein-stone of, 472.
Mules (and horses), their fondness for water holding iron-ore in sus-
pension, 854.
Mungla Lekh, on the iron-ores of, 26.
Murchison, Sir R. I., Bart., on the copper-bearing sandstones of Salop
514,-15,-16.
, (Verneuil, M. E. de, & Keyserling, Coun
A. von), on the auriferous Beresite o Siberia, 172.
, on the proportion of gold obtained a Berezovsk, 177-8.
, " gold and platinum of Siberia, 340
, ", detrital gold ", ,175
, ", a large mass (Nugget)
of, Table
XXII
copper-bearing coal-measures of Siberia, 508-9.
Murray, J. J., Esq., on the effects of an earthquake at Caldera, 158.
Music, Wheal, area of open mining-works at, 576, 665.
Nadhoolee, on the iron-furnaces at, 19.
Nanskêg (Nanskuke), on the argentiferous ore of, 119.
Napier, James, Esq., on the structure and specific-) of smelted copper,
gravity
, ,, of electro-deposit-
ed copper, 431.
absorption of oxygen by copper when in fusion, 430.
,, precipitation of copper at the Parys mine, 580-1.
National mine, on the native copper in conglomerate at the, 477.
, , proportion of , , 471.
, ,, produce and profits of the, 459; Table XIV.
wages of miners at the, 471.
Navigation of Lake Superior interrupted during winter, 434.
Nebal-bridge, detrital matter near, 45.



Oak, forests of, in the Himalaya, 55.
, rejected by charcoal-burners in the Himalaya, 55.
Oates, G., Esq., on the influence of cross-veins on the sulphur-course
of Wicklow, 558-9.
Ochre, lining crevices in the Itabirite at Agoa Quente, 225.
——, quantities of, deposited in the precipitation-works at Mona, 583-4.
Odernheimer, G., Esq., on the mica-slate of Breadalbane, 645,-50.
porphyries of , , 650.
, " lodes of " , 652.
Oil, mineral, of Borriga and Expectativa, 516.
Oldham, Dr. T., on subterranean temperature at Knockmahon, 740.
Oliveira, Col. M. F. de, gold-mines wrought by, in Brazil, 237.
Oliveira, Capt. M. J. F. de, on the gold of Catta Preta, 237.
Ollivant, S., Esq., on the gold produced in Minas Geraës, 869.
O'Niel, C., Esq., on the gold of Merionethshire, 641.
Oojowlee, on the iron-ores of, 83,-6.
Open mining-works, 222,-84,-7,-56, 576, 665.
Ores. (See Antimony, Arsenic, Bismuth, Chromium, Cobalt, Copper,
Gold, Iron, Lead, Manganese, Molybdenum, Nickel,
Palladium, Platina, Silver, Tellurium, Tin, Titanium,
Zinc.
Organic remains of the Carboniferous system, 618-14,-23.
in metalliferous deposits, 619 (?), 625-6.
in raised beaches, 156.
Ornaments, ancient, of native copper obtained from early mining- works near Lake Superior, 418-19.
Orpailleurs (gold-washers), quantities and values of gold obtained
by, from the Rhine, 348,-51-2,-8.
Ouro Fino, rocks and metalliferous deposits of, 184.
, proportion of gold in ,, , 184, 322.
Ouro Preto (Villa Rica), on the micaceous (? talcose) quartz-rocks (elastic sandstons) of, 209.
, " auriferous clay-slate of, 210,-11.
Ovalle, P. Alonso de, on the valley of Copiano, 139.
Ovoca, on the mines of iron-pyrites and copper-pyrites, in the vale of,
, produce of561.

UTOCA, OR THE	serunds or mine	rs in the vale of, 90%.
, "	precipitation of co from mine-	
 , ,, :	suriferous pyrite	s of Connorres and Ballymurtagh, 549-59.
	he proportions of ungary, 100.	silver and gold in the lead-ores of
•	ne association of a	antimony-glance and tellurium with
Pacific, shells		found also in raised-beaches near,
Pahlee (in Ram	gurh), on the spe	cular-iron of, 25.
		the brown iron-ore of, 26.
		us lead-ores of Sicily and Calabria, 102.
	" proportions	of silver in Spanish lead-ores, 102.
Pajonales, on th		nickel, and silver at, 121.
		old of Candonga, 175.
 ,	,,	Santa Anna and Itabira, 215.
	"	Gongo Soco, 286, 387,-8,-40.
	"	, in granite, 175.
	'n	, in Itabirite (Jacotinga), 215,-86,
,	"	, ,, , proportions of at different depths, 286, 888,-40,
	ome deposits which	ch yield silver-ore with others which
of	• •	r-ore in deposits having different
		the ranges of mountains in which
		21,-45,-8, 804,-10,-11; Table X.
		se near Ovoca to the cleavage of the adjoining rocks, 558.
,	, great body of	pyritous copper-} ,, 575.
,	, rich <i>bunches</i> (Kenm a re, (of copper-ore on different lodes near
 ,		some with the barren portions of
		s of the, contain chrome-ore and

Par Consolidated mines, proportions of tin-ore in the vein-stones of
Park mine, proportions of silver in the lead-ores of, 110. [472.
Parys mine, rocks of the, 574.
, metalliferous deposits of the, 575-8.
, cross-veins of, 575,
, analysis of the (blue-stone) ore of, 546,-77.
, treatment of pyritic copper-ore at, 578-9.
, drainage of the, 580.
, precipitation of copper from mine-water at the, 580-1.
, copper-turf of the, 580.
Pasco (Peru), proportions of silver in the ores of, 98.
Pascoe, Capt. H., on the lodes of Bearhaven, 608.
Patol, on the iron-ores of, 33,-7.
Pattinson, H. L., Esq., on the proportions of silver in the lead-ores
of England and Wales, 105,-7,-9,-19.
Paunchmaulee, on the clay iron-stone of, 66.
Peach, C. W., Esq., on fossiliferous rocks near Liskeard, 700.
Pebbles, beds of, far above existing streams, 45.
Pedro, San, gold imbedded in copper-glance at, 378.
Peever, Wheal, displacements of lodes at, 226.
Peixoto, Senr. J. de Souza, employment of slaves by, 301.
Pengilly, Capt. T., on the auriferous of <i>Gongo Soco</i> , 250,-7,-60,-3, deposits 266,-8,-72,-4.
, " of Agoa Quente, 288.
, " " unrecognized by workmen, 326.
, ,, treatment of auriferous sand, 854.
, ,, manufacture of charcoal, and the smelting
of iron-ore, in Brazil, 262.
Penhaldarva, proportion of silver in the lead-ore of, 120.
Pennance, proportion of silver in the lead-ore of, 120.
Pennant, T., Esq., on the treatment of at the Parys mine, 574,-6, copper-ore 577,-9.
, ,, copper-turf of ,, ,580.
Penrose, Wheal, on the argentiferous lead-ores of, 119.
Pentire Glaze, proportion of silver in the lead-ore of, 120.
Pepper, Prof., on the steam-engine placed underground at the Tamar
mines, 609.
Percy, Dr. J., on the structure and specific smelted copper, 429-30.
, ,, ,, ,, after pressure, 429,-30.



Phillips, W., Esq., on the granite and slate of Tincroft, 658.
, ,, alleged waste of copper-ore in Cornwall,
690-2.
, " silver-ores of Cornwall, 111,-12,-18,-16,
, " stream-tin-ore " , 859. [118.
birth and death of, 691.
Phænix mines, rocks of the, 656,-69.
, ,, on opposite sides of the <i>lode</i> at the, 657, 669.
, , metalliferous deposit of the, 676,-8,-9; Table XXX.
, subterranean temperature in the, 746.
, produce of, people employed at, and machinery at
work on the, 698.
Pick and gad, dexterity of Cornish miners in using the, 152.
Piedade, on the structure of the mountain, 298.
Pike, R. H., Esq., on the proportion of tin-ore in the vein-stones of
the Carn Brea mines, 472.
, , produce and profits of the , , 448,
Pindur, on the auriferous sands of the, 4, 46. [459; Table XIV.
Pinnock, Saint, on the rocks of, 99.
. 1110 1 1 0 0 000 100
, proportions of silver in the lead-ore of, 120,
, ,, produce of, people employed, and machinery
at work in the district of, 720.
Pipe-veins, of lead-ore, on, 620-1.
Pitangui, on the auriferous deposits of, 222-4.
,, conformity of some of them to the jointed structure
of the rocks, 223, 319,-23.
Plants, fossil, on, in the coal-shales of New Brunswick, 503-9.
, ,, Nova Scotia, 507-8.
S:h 508 0
Platina, associated with the gold of Brazil, 286, 335,-7,-8,-40.
of the Dhine 900
TT 1 00 T 40
, in Itabirite (Jacotinga), 286, 385, 887,-8,-40.
, , , , proportions
of, at different depths, 286, 388,-40.
, associated with stream-tin-ore in Australia, 286.
Plumbago in the clay-glate of the Himelern 41

Plunger-pole, notice of its early use in the mines of Gwennap, 570.	
, wooden, used at Connorres, in Wicklow, 569.	
Plymouth, on the temperature of, 745,-9.	
Poder, a name anciently given, in Cornwall, to certain ores of copper	
and (?) of iron, 689.	
Pokres, on the rocks and copper-ores of, 6.	
Polberro, on the produce of, 458.	
proportions of tin-ore in the vein-stones of, 202, 472.	
, ,, rate at which the vein-stones are blasted, 200.	
Poldice, proportions of tin-ore in the vein-stones of, 472.	
, loss during former operations at, 452.	
Polgooth, profit yielded by early works at, 454.	
Polkinghorne, W., Esq., on the proportions of tin-ore in the vein-	
stones of Cuddra & Par Consolidated	
mines, 472.	
Polwhele, The Rev. R., on the proportions of silver in Devonshire	
lead-ore, 109.	
, ,, silver-ores of Cornwall, 111,-14.	
Pontalba, Baron, on the proportions of pure copper in the native metal	
of Lake Superior, 471,-3.	
Pontes, Senr. M. J. P. da Silva, on the productive gold-mines of Itabira, 216.	
, on the \ , , 215,-16,	
quality of the gold afforded by \ 332.	
nonertions of polledium in \ , 215.	
proportions of panadium in)	
theft of gold by workmen at \ ", , 216.	
Pool, Wheal, on the argentiferous lead-ore of, 118.	
Pooley, Capt. E., on the proportion of tin-ore in the vein-stone of	
Trelyon, 472.	
Porphyry, felspathic, of New Brunswick, 491.	
, ,, of Breadalbane, 650-1.	
, , of Caradon, 660-1,-70-1.	
, ,, of Pra, 651.	
, stanniferous, of Geyserberg, 665-6.	
Portage Lake, on the sandstones of, 391-2.	
Potosi, on the proportion of silver in the ores of, 98.	
Potstone of Brazil, on the, 173.	
Powder, charges of, used for blasting in different mines, 200-1.	
, , , , , , , , , , , , , , , , , , , ,	

Pra, on the fe	lspathic porphyry ((Elvan) of, 651.
		iferous sand collected on, at Morro
` `		ho, 855.
Prideaux, Cap		rous deposits of Gongo Soco, 257,
	, on a landslip	
		l vein-stones of Sark, 532,-3,-4.
		, , 121, 586-7.
	-, " silver and	l silver-ores ,, , 121, 536-7.
Precipitation of	of copper from min	e-water in Anglesea, $580-2, -92$;
-		Table XVIII.a
	77	in Cornwall, 385-9,-92.
	"	in Cuba, 590-1,-2.
	"	in Devon, 584-5,-92.
	"	in Spain, 589-91,-2.
	"	in Wicklow, 562-87,-92.
·	"	, more rapid in warm, than
		in cold, weather,
		571.
	,,	, ,, in running, than
		in still, water, 571.
	,,	, quality of the precipitate
		affected by the nature of
		the precipitant, 588.
	"	, dues (Royalties) paid on,
	es. (See Mining.)	[588.
	of antimony, in gold	
	of arsenic, "	
	of bismuth, ,,	
		and silver-ores, 523.
, (of copper and copp	per-ore, in vein-stones, 426,-35,-6,
		462,-70-1,-5,-9,-82, 693.
 -,	"	, in gold, 286, 234,-8,-40,-74.
, (of gold, in vein-stor	nes, 177,-8,-9,-82,-4,-97,-8, 201,-8,
		280,-1,-2,-5,-49,-55,-8,-77,-9-
		281, 331,-4,-72-3,-4,-9,-83,
		550,-68, 641; Tables VI.ª
		VII, VIII. X.
,	" "	, at different depths, 201,-4,-5,
		284, 831; Table VII.
 ,	,, in lead and	l lead-ores, 99, 100,-1,-4.

Proportions of gold, in silver and silver-ores, 100,-1, 550.
, ,, in copper and copper-ores, 237, 378.
, ,, in different (Brazilian) rocks, 869.
, ,, in detrital matter, 845,-6,-8,-58.
, of lead and lead-ores, in vein-stones, 693.
, in gold, 334,-74,
, of palladium, in gold, 286, 338,-40; Table X.
, ,, at different depths, 286, 388,-40; Table X.
, of platina, in gold, 286, 338,-40; Table X.
, , , , at different depths, 286, 838,-40;
, ,, in detrital gold, 860. [Table X.
, of silver, in vein-stones, 91-8,-5-6,-7,-8-9, 100,-1,-2,
104,-12,-13,-14,-15,-16,-17,-18, 121, 487, 523-4,-36,-68, 711.
, ,, in copper-ore, 378.
, in lead and lead-ores, 99, 100,-1,-2,-8,-4,-5,
106-10,-18,-19,-20,
121, 586, 708-9, 11.
, , , , , at different depths, 708-9.
, , in mine-gold, , .206.
286, 335,-8,-40,-74; Table VII.
, ,, in detrital gold, 860.
, of tellurium, in gold, 336.
, of tin-ore, in vein-stones, 202, 472,-3, 693.
, , , in detrital matter, 680.
Prosper United mines, proportions of tin-ore in the vein-stones of,
Providence mines, rocks of the, 659. [472.
, proportions of tin-ore in the vein-stones of the,
, produce and profits of the, 444; Table XIV.
Providence, Wheal (Tremayne, Wheal), on the argentiferous lead-
Pryce, Dr. W., on the composition of lodes, 676,-8. [ore of, 118.
, ,, ,, when productive, 653.
, proportion of copper-ore in lodes, 693.
, on early copper-mining in Cornwall, 688-90,-1.
,, loss of copper-ore owing to careless treatment, 693.
,, argentiferous lead-ores of Cornwall, 118,-19.
, ,, stream-tin-ore ,, , 486.
, ,, precipitation of copper)
from mine-water ", , , , , , , , , , , , , , , , , , ,

Pryce, Dr	. W., on se	it water which	enters subr	narine mi	nes, dyy.
Pumps, in	n the mines	of Brazil, son	_	rked with	rag and
		C	hain, 345.		
,	"		n-work of, nufactured }		
 ,	"	,, , wood	l-work of, i	on the spo	ot, 288.
 ,	"	Cornwall, c	comparison ciency of, 5		ated, with
Punaar, or	n the graph	itic clay-slate o	•		
		production of a		3.	
	n the climat		•		
Purturbur	à, on the ire	on-ore of, 24;	Table I.		
Pyrites.	(See Ores o	f Copper and	Pres of Iron	n.)	
Quality of	f gold, in B	olivia, 860.			
	", in B	razil, 205-6,-14	,-15,-36,-7	,-85,-99, 8	317, 32-3,
		334	4,-8,-58 ; ′1	ables VII	. IX. X.
	", in C	alifornia, 339;	Table XX	Π.	
	,, , in C	nada, 384.			
	,, , in Ir	eland, 633.			
		ew Granada, 36	80.		
		beria, 335,-60.			
		irginia, 374,-83	·.		
		ales, 642.			
		fferent rocks, 8	32,-4-5; T	able X.	
		ifferent parts of			5-6, 332-
	.,		formation,	334	4; Tables I. IX.
 	", at di	fferent depths	,,	334	5-6, 332- l ; Tables I. IX.
		rge masses, infe dies in other lo			of smaller
		ent in crystals			35-6.
		trital deposits s			
		rmations, 359-6	_	· ·	
of		silver-ore in the		nestones o	f Chanar-
	cillo, 98-12				
	•	and of lead-or	e in differ	ent narts	of Sark's
V.	Hone 535			Pour en	U_ ~~

Quartz, an ingredient of rocks, 2, 3, 4, 11, 12, 18, 21,-6, 30,-1,-5-86,-9, 64-5, 78,-9, 85, 161,-4,-71,-2,-4,-7, 178,-80,-5, 211,-14,-17,-19,-21,-4,-5,-39, 244,-5,-7,-8,-50,-5,-9,-65,-97,-8, 800,-1,-2, 803,-4,-11,-12,-13,-15,-16,-21,-2,-4,-71, 881,-5,-6,-7,-91,-2,-8, 402,-8,-11,-14,-74, 475,-91,-8,-4,-5,-7,-8,-9, 501,-11,-18,-17, 563,-74,-7,-94, 602,-8,-7,-15,-86,-45,-8, 663,-7,-8,-9, 713,-14,-41,-2,-50. of metalliferous deposits, 4, 5, 6, 8, 10, 11, ** 12, 18, 19, 20,-3, 33,-7, 86, 162,-8, 165,-76,-7,-8,-80,-1,-2,-8,-4,-94,-6, 214,-23,-8,-32,-3,-4,-9,-40,-1,-9,-55,-7, 263,-98, 300,-2,-7,-11,-12,-13,-15,-17, 819,-20,-1,-2,-4,-9,-82,-4,-7,-41,-71,-2,-3, 374,-5,-6,-7,-8,-9,-81, 412,-20,-8,-4,-31, 460,-8,-4,-5,-8,-9,-79,-92, 522,-31,-3,-54, 555,-6,-7,-64,-5,-7,-75,-6,-7,-96-7, 605, 617, -24, -38, -40, -58, -75, -6, -7, -8, -9, -81, 705,-6,-13,-14,-26,-8,-9,-82,-5,-6,-9,-41,-8. of cross-veins, 226,-32,-86-7, 308,-30,-41, " 608,-82, 716.

Quebrada Seca, on the rocks and metalliferous deposits of, 161-8.

Quetelet, M. L. A. J., on the Magnetic declination at Bilbao, 518.

-----, ,, temperatures observed at small depths in Brussels, 773-5.

Quincy mine, on the expenditure and profits at the, 459; Table XIV. Quito, on the mean temperature of, at the surface, and at inconsiderable depths, 771.

Ralfs, John, Esq., on the identity of diatoms in Sirocco-dust at

Malta, and in the raised and actual beaches
of the Pacific, 160.


Ramgunga, auriferous sands of the, 4, 46.

, detrital matter in the neighbourhood of, but far above its present bed, 45.

Ramgurh, on the iron-ores of, 25.
Rammelsberg, on the iron-pyrites (Sulphur-ore) of, 544,-6.
Ramsay, Prof. A. C., on the Silurian rocks of Merioneth, 635,-6,-7. Copper-turf of ,, 580.
Rampoore, on the conglomerate and iron-ore of, 80; Table II.
Rancié, ventilation effected there by means of a (trompe) water-fall 220.
Raspe, H. R. E., improved operations at <i>Dolcoath</i> , whilst he was employed there, 645-6, 7.
, his presumed authorship of Munchausen's Travels,
Rations, of miners in Chafarcillo, 150; Table V. [645.
, of slaves in Brazil, 292.
, ,, Virginia, 382.
Ratios at which the temperatures of mines increase with their depths, 759.
Rawlings, W. J., Esq., on the silver-ores of West Darlington, 116-117.
, proportions of tin-ore in the vein- stones of Wheal Busy, 472.
Readwin, T. R., Esq., on the Clogau gold-mine, 635,-8.
Reay, W., Esq. (& Walker, Dr. T.), on the proportion of gold contained in clay-slate at Morro Velho, 186,-96.
of (dressing) cleaning gold at Morro Velho, 203.
Redruth, subterranean temperature in the neighbourhood of, 752.
Reduction-works, proprietors of, take their proportions of ore at the mines in Chili, 94.
, produce of ores profitably treated at, in Chili, 94.
, force employed at, in Morro Velho, Table VI.ª
, wages of work-people in ,, , Table VI.ª
, proportions of gold extracted in , , Table VI.
, operations of, in Gongo Soco, Table IX.
Reed, J., Esq., on the rate of blasting vein-stone at Cronebane, 200.
, ,, diminished proportions) 578
or copper in the water)
Rocks, of granite, sprinkled with gold, 4, 175, 811,-20.
—, ,, , tin-ore, 664-5.

Rocks of	talcose and micaceous	alata anri	nkled :	with gold, 177, 210-
mound, or	unidono una mitadoco un	nimeo, ppri		299, 301,-11,-20.
		_		vitreous copper,
,	"	,	"	495.
, of	clay-slate,	•••••	"	gold, 196, 210.
	calcareo-siliceous na		"	gold, 245,—9,
•		•	"	304,-19,-24.
——, of	Itabirite and Jacotin	ga,	"	gold, 214,—15,
				219,—42,—9,
				255,-99, 325.
, of	conglomerate	•••••	"	native copper,
				403,-75.
, of	limestone,	•••••	"	native silver and
				the ores of silver, 76.
, of	the coal-measures,		"	copper-ore, 503,
				506-10.
, of	the New Red Sandst	one,	"	copper-ore, 514.
				(gold, 217,—37,
, C	anga,		") 316.) native copper,
·	•			236,-316.
Rees, Dr.	A., on the silver-ore of	Wheal Je	wel (n	ear Callington), 116.
				e lead-ores of Beer
	Alston, 1	09-10.		
	F., on the temperatu		es in S	Saxony, 757.
	(See Organic Remain			
Rhine, on	the auriferous sands	•		
				, 34 8,- 51 ,- 8 .
 ,	>>			and earnings of the
				loyed on them, 358.
Richards,	Mr. I., on the precipi		opper	at the Devon Con-
	solidated m	•	_	
Kichards,	Capt. J., on the comp			
D: 1 1		gons in m		
	W., Esq., on ventilat			
	neiro, on the mean te			
	o, on the (pyrites) sulp		-	•
D: 1.				
rippie-str	ucture in schistose ro	cks, its rel	ation	to subjacent granite, 259.
			. 1	to shoots (masses) of
	8 z	, ,		gold, 259,-63,-4,-9.
		•		o,,, -, -, ••

Rita, Santa, on the auriferous talc-slate of, 177.
Rivers, their disappearance beneath, and reappearance beyond, beds
of detritus, 48,-4, 858.
, their gradual diminution on the destruction of forests at their sources, 137-41, 348-8.
, existing, unequal to transporting the coarser and heavier detritus, 348.
, which receive much mineral-water, existence of fish in, 354-
855, 717,-81.
Rivot, M. L. E., on the rocks of the Lake Superior \ 891,-2,-4,-5,-6, mining district, \ 897,-8,-9,400, 401,-2,-8,-4.
, ,, metalliferous) ,, 405,-6,-7,-23, 424,-5,-32,-4,-5, 436,-7,-8,-9,-61, 462,-4,-75.
, mode and cost of extracting (native) copper from the mines
, " produce of the mines at , , 488.
, ,, ancient mining works at , , 412-18,-17.
different proportions of silver obtained from the lead- ores of the same mines
, (& Duchanoy, M.), on the proportions of silver and gold in the lead-ores of Hungary, 100.
, (& Zeppenfeld, M.), on the proportions of silver in
the lead-ore of <i>Pontgibaud</i> , 104. Robert, North Wheal, proportion of tin-ore in the vein-stone of, 472.
Roberts, Capt. T., on the proportion of tin-ore in the vein-stone of
Wheal Mary, 472.
Robinson, Henry, Esq., on heaves of the Sulphur-course by joints at Ballygahan, 560.
Robinson, T. W., Esq., on profits made in the Alfred Consolidated mines; 445; Table XIV.
Rockland mine, on the rocks and metalliferous deposits of, 480-1.
Rogers, Prof. W. B., on the rocks of Willis's mountain, in Virginia, 879.
, " matrix of gold, " , 875-
, ,, argentiferous gold of Virginia, 874.



Rule, John, Esq., his plans and sections of the works at Dolcoath, 448. Rutter, Capt. W., on the comparative economy of wheel-barrows and tram-waggons in mines, 144. Sabará, prosperous condition of the gold-mines near, 368. Sabine, Gen. Sir E., on Magnetic declination, 372, 492, 513,—15, 520,-31,-41,-75, 603,-16,-37,-47,-73, 703,-37. Sacramento (Chili), exportation of silver from, 153, Sacred-lakes (Thibet), notice of gold obtained from them, 48. Sahloo, on the rocks and iron-ores of, 19. Salmon, on the abundance of, in New Brunswick, 499. Salmon, H. C., Esq., on the lead districts of East Cornwall, 699,-704. lode of Wheal Mary Ann and Wheal Trelawny, 706,-9,-14. lode of Herod's-foot, 712,-13,-14. silver-ores of Wheal Ludcott, 710. ,, refuse of ancient lead-mines in the Mendips, 625. winding-engines of Carn Brea, 143. Salter, J. W., Esq., on the fossils of the auriferous series in Wales, Sampson, B., Esq., his survey & plans of Wheal Peever, 226. [636. ----, on early use of the plunger-pole, 570. Sand, sounds emitted by it, when in motion, at El Bramador, near Copiapò, 154. -, siliceous, an ingredient in the Carvoeira of Catta Preta, 239. Sandstone, recent, near Caldera, 155. -, of the Sub-Himalaya, 38. —, of Huidobro, 77, 513. –, of Shropshire, 514-15. -, of Bengal, 64,-6,-8. —, of New Brunswick, 501,-2,-3. , of Lake Superior, 392,-3. -, of Kenmare, 612. -, (elastic) of Villa Rica, 172, 210. -, containing the ores of copper, 77, 513-15. of iron, 64,-5,-6,-8, 513. ,, -, affording Petroleum, 516. —, subterranean temperature in, 751,-3,-8. Santiago (Cuba), on the precipitation of copper at, 590-1,-2;

Tables XXXI. XXXIII. XXXV.

Santos, Dr. M. J. dos, on the produce of go Saō Joaō do Morro Grande, detrital gold (aid of a chai	
Sark, on the rocks and metalliferous deposits	s of, 530-8.
, on submarine works at, 584-9.	•
, on subterranean temperature at, 735,-	6,-52.
, on the climate of, 735.	
Sauvage, M., on the proportions of silver coore, 102.	ntained in Spanish lead-
Savings-Bank, established for the slaves at	Gongo Soco, 293.
Saxony, tin-ore an occasional ingredient of	granite in, 77.
, proportions of silver in the ores of	
Scheerer & Marchand, MM., on the specific	
compression	, 480.
Schlagintweit, H. A. H. & R., on Magnetic	declination, 3.
Schorl, an ingredient of the rocks, 2, 175,	
,, of metalliferous depos	its, 240, 675.
	, 174.
Schreiber, M., ,, rocks of Ch	
on the metalliference)	
deposits (
, " ores	,, , 103, 5 23.
, " cross-vein	", 527 .
, ,, plans and sections of	works by, 520.
Schüch, Dr. R., his iron-furnaces at Timbo	
Scotland, proportions of silver in the lead-or	
Seaton, the valley of, injured by water, mu	d, and rubbish from the
mines near Caradon, 353.	
Seccombe, Capt. S., on the positions of the	rocks on opposite sides
of the lodes near Ca	radon, 658.
Sedgwick, The Rev. Prof., on the rocks of	Liskeard, 700.
Seera, on the rocks and copper-ores of, 5.	
Seetzen, M., on the noises emitted by movin	
Selwyn, A. R. C., Esq., on the auriferous g	ranite of Australia, 175.
	etritus of ", 343.
(& Ulrich, G. F. H.,	Esq.), on a large mass of
·	Australian mine-
	gold, Table XXII.
"	, on the chrome-ore
	of Australia, 649.
27	, on the molybdenite
	of Australia, 654.

Seton, West Wheal, on the produce and profits of, 449; Table XIV. Shales of the coal-measures, at Paunchmaulee, 66. , containing near Bathurst, 503—fossil plants, New Brunswick, 509. , so near Bathurst, 509. Shew Brunswick, 509. , so near Bathurst, 509. , so near Bathurst
Shales of the coal-measures, at Paunchmaulee, 66. , containing near Bathurst, 503— fossil plants, New Brunswick, 509. , 503,-9, and the ores of copper, 510. Shanagarry, on the rocks and lead-ores of, 619-21. Sharp Tor, on the different positions of granite and slate on opposite sides of a lode near, 659. Shealgar, on the iron-ores of, 22. Sheep used as beasts of burthen in the Himalaya, 47. Shells, of existing species in the raised beaches of Chili, 156. , n, in recent sandstone near Caldera, 155. , n, burnt for the lime they contain, 155. Sheriff, J. D., Esq., on the elevation of Menheniot, 746. Shingle, gravel, sand, and shells in the raised beaches near Caldera, Shoots of copper and copper-ore, 438, 598, 608. , of gold, 207,-15,-24,-59,-63,-4,-9,-70, 319,-26,-80. , of iron-ores, 32, 558,-64. , of lead-ore, 537. , of silver and silver-ores, 122, 537, , of various metals and ores, from the granite, 32, 122, 207, 259,-63, 438. , confront one another in parallel deposits, 269,-70, 326. , oblique to the strike and dip of
fossil plants, \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \
fossil plants, \ New Brunswick, \ \ 509. \[\begin{align*} \text{ 503,-9, } \\ \text{ 510.} \
fossil plants, \ New Brunswick, \ \ 509. \[\begin{align*} \text{ 503,-9, } \\ \text{ 510.} \
and the ores of copper, \$\) Shanagarry, on the rocks and lead-ores of, 619-21. Sharp Tor, on the different positions of granite and slate on opposite sides of a lode near, 659. Shealgar, on the iron-ores of, 22. Sheep used as beasts of burthen in the Himalaya, 47. Shells, of existing species in the raised beaches of Chili, 156.
and the ores of copper, \$\) Shanagarry, on the rocks and lead-ores of, 619-21. Sharp Tor, on the different positions of granite and slate on opposite sides of a lode near, 659. Shealgar, on the iron-ores of, 22. Sheep used as beasts of burthen in the Himalaya, 47. Shells, of existing species in the raised beaches of Chili, 156.
Sharp Tor, on the different positions of granite and slate on opposite sides of a lode near, 659. Shealgar, on the iron-ores of, 22. Sheep used as beasts of burthen in the Himalaya, 47. Shells, of existing species in the raised beaches of Chili, 156. ———————————————————————————————————
site sides of a lode near, 659. Shealgar, on the iron-ores of, 22. Sheep used as beasts of burthen in the Himalaya, 47. Shells, of existing species in the raised beaches of Chili, 156. ——, ,, in recent sandstone near Caldera, 155. ——, , burnt for the lime they contain, 155. Sheriff, J. D., Esq., on the elevation of Menheniot, 746. Shingle, gravel, sand, and shells in the raised beaches near Caldera, Shoots of copper and copper-ore, 438, 598, 608. ——, of gold, 207,-15,-24,-59,-63,-4,-9,-70, 319,-26,-80. ——, of iron-ores, 32, 558,-64. ——, of lead-ore, 537. ——, of silver and silver-ores, 122, 537, ——, of various metals and ores, from the granite, 32, 122, 207, 259,-63, 438. ——, ,, confront one another in parallel deposits, 269,-70, 326. ——, , oblique to the strike and dip of
Shealgar, on the iron-ores of, 22. Sheep used as beasts of burthen in the Himalaya, 47. Shells, of existing species in the raised beaches of Chili, 156. ——, ,, in recent sandstone near Caldera, 155. ——, ,, burnt for the lime they contain, 155. Sheriff, J. D., Esq., on the elevation of Menheniot, 746. Shingle, gravel, sand, and shells in the raised beaches near Caldera, Shoots of copper and copper-ore, 438, 598, 608. [156. ——, of gold, 207,-15,-24,-59,-63,-4,-9,-70, 319,-26,-80. ——, of iron-ores, 32, 558,-64. ——, of lead-ore, 537. ——, of silver and silver-ores, 122, 537, ——, of various metals and ores, from the granite, 32, 122, 207, 259,-63, 438. ——, ,, confront one another in parallel deposits, 269,-70, 326. ——, , oblique to the strike and dip of
Shealgar, on the iron-ores of, 22. Sheep used as beasts of burthen in the Himalaya, 47. Shells, of existing species in the raised beaches of Chili, 156. ——, ,, in recent sandstone near Caldera, 155. ——, ,, burnt for the lime they contain, 155. Sheriff, J. D., Esq., on the elevation of Menheniot, 746. Shingle, gravel, sand, and shells in the raised beaches near Caldera, Shoots of copper and copper-ore, 438, 598, 608. [156. ——, of gold, 207,-15,-24,-59,-63,-4,-9,-70, 319,-26,-80. ——, of iron-ores, 32, 558,-64. ——, of lead-ore, 537. ——, of silver and silver-ores, 122, 537, ——, of various metals and ores, from the granite, 32, 122, 207, 259,-63, 438. ——, ,, confront one another in parallel deposits, 269,-70, 326. ——, , oblique to the strike and dip of
Sheep used as beasts of burthen in the Himalaya, 47. Shells, of existing species in the raised beaches of Chili, 156. ——, ,, in recent sandstone near Caldera, 155. ——, ,, burnt for the lime they contain, 155. Sheriff, J. D., Esq., on the elevation of Menheniot, 746. Shingle, gravel, sand, and shells in the raised beaches near Caldera, Shoots of copper and copper-ore, 438, 598, 608. ——, of gold, 207,-15,-24,-59,-63,-4,-9,-70, 319,-26,-80. ——, of iron-ores, 82, 558,-64. ——, of lead-ore, 537. ——, of silver and silver-ores, 122, 537, ——, of various metals and ores, from the granite, 82, 122, 207, 259,-63, 488. ——, ,, confront one another in parallel deposits, 269,-70, 326. ——, , oblique to the strike and dip of
Shells, of existing species in the raised beaches of Chili, 156. ———————————————————————————————————
for the lime they contain, 155. Sheriff, J. D., Esq., on the elevation of Menheniot, 746. Shingle, gravel, sand, and shells in the raised beaches near Caldera, Shoots of copper and copper-ore, 438, 598, 608. [156, of gold, 207,-15,-24,-59,-63,-4,-9,-70, 319,-26,-80, of iron-ores, 32, 558,-64, of iead-ore, 537, of silver and silver-ores, 122, 537,, of various metals and ores, from the granite, 32, 122, 207, 259,-63, 438, confront one another in parallel deposits, 269,-70, 326, oblique to the strike and dip of
for the lime they contain, 155. Sheriff, J. D., Esq., on the elevation of Menheniot, 746. Shingle, gravel, sand, and shells in the raised beaches near Caldera, Shoots of copper and copper-ore, 438, 598, 608. [156, of gold, 207,-15,-24,-59,-63,-4,-9,-70, 319,-26,-80, of iron-ores, 32, 553,-64, of lead-ore, 537, of silver and silver-ores, 122, 537,, of various metals and ores, from the granite, 32, 122, 207,
Sheriff, J. D., Esq., on the elevation of Menheniot, 746. Shingle, gravel, sand, and shells in the raised beaches near Caldera, Shoots of copper and copper-ore, 438, 598, 608. [156, of gold, 207,-15,-24,-59,-63,-4,-9,-70, 319,-26,-80, of iron-ores, 82, 558,-64, of lead-ore, 537, of silver and silver-ores, 122, 537,, of various metals and ores, from the granite, 32, 122, 207, 259,-63, 438, confront one another in parallel deposits, 269,-70, 326, oblique to the strike and dip of
Shingle, gravel, sand, and shells in the raised beaches near Caldera, Shoots of copper and copper-ore, 438, 598, 608. [156, of gold, 207,-15,-24,-59,-63,-4,-9,-70, 319,-26,-80, of iron-ores, 32, 553,-64, of lead-ore, 537, of silver and silver-ores, 122, 537,, of various metals and ores, from the granite, 32, 122, 207, 259,-63, 438, , confront one another in parallel deposits, 269,-70, 326, oblique to the strike and dip of
Shoots of copper and copper-ore, 438, 598, 608. [156. ————————————————————————————————————
, of gold, 207,-15,-24,-59,-63,-4,-9,-70, 319,-26,-80, of iron-ores, 32, 553,-64, of lead-ore, 537, of silver and silver-ores, 122, 537,, of various metals and ores, from the granite, 32, 122, 207, 259,-63, 438, , , , , confront one another in parallel deposits, 269,-70, 326, , , , oblique to the strike and dip of
, of iron-ores, 82, 558,-64, of lead-ore, 537, of silver and silver-ores, 122, 537,, of various metals and ores, from the granite, 82, 122, 207, 259,-63, 488, , , confront one another in parallel deposits, 269,-70, 326, , , oblique to the strike and dip of
, of lead-ore, 537, of silver and silver-ores, 122, 537,, of various metals and ores, from the granite, 82, 122, 207, 259,-63, 488, , , , confront one another in parallel deposits, 269,-70, 326, , , oblique to the strike and dip of
, of silver and silver-ores, 122, 537,, of various metals and ores, from the granite, 32, 122, 207, 259,-63, 438, , , , confront one another in parallel deposits, 269,-70, 326, , , oblique to the strike and dip of
259,-63, 488. , confront one another in parallel deposits, 269,-70, 326. , oblique to the strike and dip of
deposits, 269,-70, 326. , oblique to the strike and dip of
deposits, 269,-70, 326. ——-, ,, oblique to the strike and dip of
adicinia
adjoining rocks, 32, 122, 259.
269.
, conform to the rippled (undu-
lated) structural planes of con-
tiguous rocks, 207,-15,-24,
259,-63,-4,-9,-70, 819,-26,
880, 488, 587, -53, -64, -98,
608.
Shropshire, copper-ores in the New Red Sandstone of, 514-15.
, proportion of silver in the lead-ores of, 109.
, subterranean temperature in, 751,-2,-3,-8; Tables

Siberia, on the auriferous rocks of,	
, ,, detrital gold of, 848	
, , alloys of gold in, 88	
deposits, 847.	by parties working auriferous
, " earnings of workpeo	ple in, 848.
Sicily, proportion of silver in the le	
Silliman, Prof. B., on proportions o	
Silurian rocks, of Wicklow, 540-2,	
, of Waterford, 598-	
, of Merioneth, 635-6	
Silver and the cros)	is gold-mine at Pitangui, 222-3.
Silver and the ores of silver, in the Banat,	
, in Chili, 76, 9	
1	Bal-dhu, 114.
	Crennis, 118.
	Cubert (?), 111.
	Dolcoath, 112-13.
	Fowey Consolidated mines, 118.
	Herland, 111-12.
	Levant, 110.
	North Dolcoath, 118.
	North Downs, 114.
•	Trebisken, 120-1.
	Treskerby, 114.
	West Darlington, 116-17.
	Wheal Alfred, 112.
	Wheal Ann, 112.
	Wheal Basset, 113.
, in Cornwall,	Wheal Cock, 110.
	Wheal Duchy (Wh. Brothers), 114-15.
	Wheal Jewel (Calstock), 116.
	Wheal Ludcott, 120, 710-11.
	Wheal Mexico (Perran-zabuloe) 110-11.
	Wheal Mexico (Calstock), 115- 116.
	Wheal Providence (Tremayne), 118.
	Wheal Saint Vincent (East
	Cornwall mines, Wheal
	Langford), 115.
	Wheal Sisters, 115.
	Wilsworthy, 116.

```
Silver and the ores of silver, in France, 103,-4, 525-6.
                 -, in Hungary, 100-1.
                 -, in Ireland, 105, 545,-9,-68.
                 -, in Mexico, 97,-8,-9.
                --, in New Granada. 98.
                 -, in Norway, 100.
                 --, in Peru, 98.
                --, in Sark, 121, 535-7.
                --, in Saxony, 101-2.
                 -, in the United States, 99, 436-8.
                  -, associated with bismuth, 111.
                                    ores of cobalt, 103,-11,-12,-16,
                                                     523.
                                    copper and copper-ore, 99, 104,
                                     112, 378, 436-8, 549,-68, 642.
                                    ores of lead. (See Lead.)
                           ,,
                                     " of nickel, 103, 524.
                           ,,
                                    gold, 99, 104,-5, 206,-86,-86,
                           "
                                           334,-5,-7,-8,-40,-60,-74,
                                           378, 549, --68, 632, --42;
                                           Tables VII. IX. X.
                                     ,, in different rocks, 206,-36,
                                          286, 334-5,-7,-8,-40,-60,
                                           374.
                                     ,, , in the same rocks at differ-
                                           ent depths, 206,-86, 338,
                                           340.
                 --, temperatures of mines affording, 724,-35-6,-56;
                      Tables XXXIII. XXXIV.
Simpkins, I., Esq., on the produce of copper-mines near Lake
                       Superior, 488.
Simul Khet, on the quartz-rocks of, 30.
                   iron-ores of, 29; Table II.
               ,,
                   iron-furnaces of, 57-8.
Siranowski (Altai), on the alloy of gold at, 335.
Sirowlee, on the iron-ores of, Table II.
Sisters, Wheal, on the proportion of silver in the ores of, 115.
Skerrett, Col. J. M. A., on the auriferous) Gongo Soco, 257,-72,-8;
                              deposits of).
                                                       Table XXII.
                                                     , 350.
                              land-slip at
```

<i>Skips</i> , on	their	early use in inclined shafts near Tavistock, 144, 668.
— p	"	later introduction ,, in West Cornwall, 668.
<i>Slags</i> , on	the	proportions of metal in the slags of Brazilian iron-
		furnaces, 261.
Slate. (S	See C	Thlorite, Clay, Hornblende, Mica, Talc, Cleavage,
·	Join	nts.)
Slaves, in	Virg	inia, hire of 382.
	,,	, food of, 382.
, in		il, hire of, 882.
	3 2	, employed alternately in mining and in agriculture,
•	"	299.
 ,	,,	, permitted to search for gold on their own account
,	"	during holidays, 801.
		, general economy of the establishment at Gongo
,	"	Soco, 289–96.
		, their occupations and hours of labour, 291-8.
,	. ,,	, the disposal of their leisure, 298.
,	"	
	"	, their food, clothing, and dwellings, 290-3.
 ,	"	, ,, earnings (dependent on individual skill and
		industry), 292-3.
 ,	"	, " " , how appropriated, 292.
, ,	"	, ", ", sometimes deposited, at interest, in
		a Savings-Bank, 298.
 ,	"	, treatment of old and infirm, 293.
 ,	"	, ,, and instruction of children, 293,-5.
 ,	"	, ,, ,, their pro-
		ficiency at school, 295-6.
 ,	"	, the aptest scholars, often the most skilful workmen,
		296.
 ,	"	, periodical inspection of their gardens, houses,
		furniture, and clothing, 291.
—-,	"	, , , followed by rewards to the
		clean, industrious,
		and careful, 291.
——,	"	, ,, the infliction of
		fines on the dirty, idle, and extravagant, 291.
 ,	"	, the ill-behaved were few in number, 295.
 ,	"	, ,, , nature of and punishments for
-		their offences, 293-5.
,	"	, medical treatment of, 295.
 ,	"	, ,, , rations whilst under, 295.
•	••	
		▼ =

Slaves, in	Braz	il, Di	vine '	worship, ob	Bervance	of, 295.	
,	7)	, nu	ım ber	employed,			•
 -,	"	,	"	who receiv	red their	freedom, 290.	•
,	"	,	37	in Minas (Jeraës, S	62.	
 ,	"	,	"	"	, e	mployed on d deposits, 358.	etrital
Slickensid	e, of	nativ	е сор	per and ea	rthy ma	tter at the Cl	iff mine, 488.
Slide, infl	uence	of a	, on t	he aurifero	as deposi	its of Gonge Se	oco, 252, 66,-70.
dist	olacen	nent c	ccasi	oned by a.	in the	lode of the Cl	
				mite of Car			[439 .
				malaya, 54			•
 ,				, 2 12,-18-1		•	
 ,	,,	, near	Lak	e Superior,			
 ,	silve	r-, in	Chili	, 97.			
 ,	iron-	, earl	y, in	India, 40,	65.		
 ,	copp	er-, ,,	, in	Chili, 154.			
Smyth, R	. B., E	i sq., o	n the	quantities tion of	and prop gold in		as quartz Lustralia, ble XXII.
-		— ,	,,		"	detrital d Australia, Tab	
		 ,	"		nd weigl	hts of masses (nuggets)
						ustralia, Table	
		 ,	"	cost of ext	racting g	gold in Austral	XXII.
		 ,	"	value of n	nining m	nachinery in A Table	Australia, EXXII.
		—,	"	molybdeni	te of A	ustralia, 654.	
Smyth, V	7. W.	, Esq	., on	the Silurian	rocks o	f Clogau, 635	
			, ,			eposit which i	ntersects
					638-9.	of Wicklow, 5	<i>4</i> 0 1 9
			, ,		n of the		±0,-1,-2.
			, ,		lphur-co	weman \ I	
					alliferou		
				pos	it at O	voca,)	
·			, ,	, altitude	of "	, in vario	
			, ,	, width o	of "	, ,,	, 544,-7,
			,	, structu	•••	, 551,-2,-5	
				. composit		. 54750.	_

Smyth, W. W., Req., on the conformable bodies of slate which inter- lie the ore, 551.
, on parallel ranges of Sulphur and copper-ore, 547,-53-4,-6.
, on the occurrence of native copper in joints
and crevices, 556.
, "cross-deposits" and
"cross-fissures," 556-7.
, on displacements of the Sulphur-course, 559—[560.
, on the produce of mines in Wicklow, 561.
, ,, rocks of Bearhaven, 602,-8-4.
, , metalliferous deposits of ,, , 603—4, 610-11.
, , , detrital gold of Wicklow, 628,-9,-30-631,-2,-4.
, " " tin-ore of " , 630-1.
, ,, alleged waste of copper-ore in Cornwall, 690.
, ,, early use of the plunger-pole in Cornwall, 570.
, on wooden linings for pumps conveying acidulous water, 570.
, ,, submarine mines near Whitehaven, 599, 600.
, " subterranean temperature, 727,-53-4,-9, 762-3,-5.
Snow, S. T., Esq., on the proportion of fine copper contained in the crude (native) metal of
Suows, of the Himalaya, 17. [459; Table XIV.
, in Dauphiny; the impediments they offer to mining in elevated regions, 528.
Soanes (miners), in the Himalaya, implements used by, 58-9.
Soares, Senr. Luïz (de Gouvea), on the rocks and auriferous quartz-
formations in the mine of, 182, 247, 327.
Socorro, gold in auriferous sands near, resembles the gold of <i>Gongo Soco</i> , 359.
Soda-amalgam, experiments with, on the ores of Morro Velho, Table XXII.
Soda, salts of, contained in river-water at Copiapò, 140.

Soda, salts of, incrust a great exten	t of surface	ne ar M o		nargo, 9-40.
,, washed out of the so		lilla, bei	ore it	an be
successfully cultiv		_	_	
Soils dissimilar, different qualities				
Socal, on graphite imbedded in clay	-slate, near	he, 41.	[219	,-890.
South Caradon. (See Caradon, S	South.)			
South Cliff mine, on the drift of the	ne, 460.			
, ,, rock, vein-	stones, and	native c	opper o	of the,
		460-3;	Table :	XIII.
Southey, Rob., Esq., on the detrit	al gold of	Brazil,	843,-60) .
, ,, Royalties	payable on		361, -2 ,	
, " restriction	s on the a	,, , {	864,-5.	
, , quantities		", ,e	xported out pa	l with- yment
			of Ro	yalty,
			365,-7	
, ,, debasemen	nt of	,, ,b		gglers,
South Tolgus mine, on the produce				⌈865.
Souza, Senr. M. T. de, on the am				_
Government				•
			go Boci	,
Spain, proportions of silver in the		102-0.		
Speculator, sharp practice of a, 51				
Spix, Dr. J. B. von (& Martius, Dr	. C. F. P. vo			
			Spix,	
Springs of water, how affected by forests, 137-8		ction of	neighb	ouring
Spring-water, precipitation of copp		Crow's-	nest. 68	36.
Squier, E. G., Esq. (&) on ancient r	•		•	
Davis, Dr. E. H.),		- MOME 2	unc ou	416.
	rnaments ar	ıd)		
	ons of coppe		"	, 419.
, on the mixt			er and	copper
-	mplements o	•		
	hworks, 418		nom e	111010110
Stalactitic iron-ore, 225,-46.	nworks, 410	'•		
Otinillaala af fuina)				
the iron	amp-heads t	t the Cl	iff min	e , 42 6.
, weight of	,, a	t <i>Morro</i>	Velho,	Table VII.
, "	,, a	t Gongo	Soco,	Table

Stamping-mills.	weight of stamp-	eads, at Agoa Que	nte. Table XXII.
	_		ta, Table XXII.
	"		, Table XXII.
,	"		, Table XXII.
·	**	, at Copper 1	
,	"	, at Gopper 1	
,	height to which		•
,	heads are life	ed { at more real	ho, Table XXII.
,	"	at Gongo Soc	
 ,	"	•	Table XXII.
,	"	at Copper Fo	ills, 426.
 ,	rate at which theads are work		ho, Table VII.
	"	at Gongo Soc	o, Table IX.
<u> </u>	"	at Copper Fo	
 ,	quantity of wa		•
•	admitted to aid stamping	1	ho, Table VII.
	numbro	at Gongo Soc	o Table IX
,	nranaration of w	in-stone for the S	
,	proparation of v		eweenaw district,
 ,	weight of	_	at <i>Morro Velho</i> , Table VII.
,	"	" "	at Gongo Soco, Table IX.
 ,	"	"	in California, Table XXII.
 ,	3 >	"	in Australia, Table XXII.
 ,	"	" "	at Copper Falls, 426.
·	27	"	at the National mine, 471.
	cost of stamping	" in Californ	nia, Table XXII.
·	"		lia, Table XXII.
Stamp-work. h	ow prepared, 425		,
		ity of, at the Clif	mine, 484-5.
	n	Nat	ional 3471-2.
,	"	, ,, Lake	Superior districts enerally, 482.

Stanniform		amita al	Comme	ll and Devon, 175, 664-6.
Statustics, 8	unu	ry, or C	opper-mi	ning, in Cornwall, 442,-5,-7,-8,-9,
				450,-1,-2,-4,-6,-7,-9, 698,
				694,-5,-8; Table XIV.
 ,	**	,	"	in Devon, 458,-9; Table XIV.
	"	,	"	in Ireland, 442, 601; Table XIV.
	27	,	,,	near Lake Superior, 440,—59,
•	-	•	••	470,-1,-3,-82,-8,-9; Table
				XIV.
	٠,	,	"	in Chili, 154,-66-7.
	"		"	in Cuba, 441; Table XIV.
		of 0		ng, in Wales, 641; Table XXII.
,	"			, in Ireland, 633.
,	"	,	"	, in Australia, Table XXII.
	"	,	"	
 ,	"	,	"	, in Brazil, 179, 208,-35,-47,-83,
				289, 327,-58,-61-8,-66-9;
				Tables VII. IX. X. XIV.
				XXII.
,	"	,	"	, in California, Table XXII.
 ,	,,	,	"	, in Nova Scotia, Table XXII.
,	"	,	"	, in Siberia, 177-8, 346; Table XXII.
	,,	, of	iron-min	ing, near Lake Superior, 389-90.
	• • •	,		, in Ireland (Sulphur-ore), 561-2.
	"	, of); lead-min	ing, in Cornwall, 454, 693, 719,-20;
 ,	"	, 01	IOMI-III	Table XIV.
	"	,	,,	, in Wales, 459,-60.
 ,	"	, of si	ilver-mini	ing, in Cornwall, 111-12,-13,-15,
				116-17,-21, 710-11.
 ,	"	•	"	, in Chili, 95-6, 123-4, 51,-3; Tables IV. V. XIV.
 ,	,,	, of ?	Fin-minin	g, in Cornwall, 442,-3,-4,-6,-7-8,
•		•		452-3,-72, 693; Table XIV.
Steam-engi	nes.	numbe	r and din	nensions of, on the mines of Caradon,
	,			d St. Pinnock, 698, 720.
				ound at the Bearhaven and Tamar
		mine	s, 609.	
				ked with distilled sea-water, in Chili,
				on, quality of, 55. [141.
Stephen, G	. M.	, Es q.,	on the de	etrital gold of Australia, 356,-9.

Stevenson, James, Esq., on the depths of Australian gold-mines, Table XXII.
Stones, sculptured, at Dhoora Devi, in the Himalaya, 19.
Stopes (Steps), substitution of one system of, for another, 145-6, 645-646,-98.
, back, adopted earlier in Germany than in Cornwall, 646.
Stowes mine, rocks and metalliferous deposits of the, 676-7.
Strachey, Capt. R., on beds of graphite near Almora, 41.
, on the disappearance and reappearance of rivers in the Turace, 44.
Strata, dissimilar, their different influence on (masses) shoots of ore
and of vein-stone in the metalliferous deposits which intersect them, 187, 224.
their equality positions in the expects sides
(walls) of metalliferous deposits, 87, 126,-9.
, , , their coincident positions on the opposite walls of
certain cross-veins, 71, 126,-9.
, ,, their different positions on the opposite walls of certain cross-veins
opposite watte of certain 225-0,-36-7, 541,
07088-08888) 042.
, , which merge in
them elsewhere, 226-7,-87, 842.
Streams, action of, on the matrix of tin-ore, 354.
gold, 854.
Stream-tin-ore, of Caradon, 695.
, of Australia associated with gold and platina, 286.
Strêkes, their utility in cleaning (dressing) auriferous sand in Brazil, Tables VII, IX.
Strise, in metalliferous deposits, 207, 483,-69, 552-3, 648,-54.
, in cross-veins, 125, 688.
, crooked and divergent, 469, 683.
Structure. (See Cleavage, Joints, Cross-veins, Metalliferous deposits.)
Submarine mining-works, at Sark's Hope, 584,-9.
,, at Knockmahen, 594,-9-600.
, at Wheal Margery, 599.
,, at Levant, 599.
,, at Botallack, 599.
,, at Whitehaven, 599.
Sulphur-ore (iron-pyrites), of Wicklow, 543-69.
Surface-water intercepted by impermeable limestone at Chanarcillo,

```
Superior, Lake, rocks on the southern shore of, 385-404.
              -, deposits of iron-ore of
                                              , 388-90.
                                              . 419-83.
                            native-copper
                           native silver
                                              , 436.
                     22
               , ancient mines and )
                                              , 412-19.
                            miners


    climate of, 477-8.

              -, subterranean temperature near, 733-4.
              -, ice remaining during summer in some of the mines
                   near, 465,-78, 734.
Surubim, unesteemed fresh-water fish, lives in rivers polluted with
                                          mineral-water, 355.
Swanpool, proportion of silver in the lead-ore of, 120.
Swings of iron-chain numerous near heathen temples in the Hima-
                                                           laya, 28.
Symons, Mr. F. S., on the proportion of gold in (killas) slate at
                      Morro Velho, 186.
Taboleiro, on the manufacture of iron at, 260-2.
        -, ,, mass (nugget) of detrital-gold found near, 283.
Tacquarigua, level of water in the lake of, affected by the condition
                of neighbouring woods, 188.
Talapoongla, on the rocks and copper-ores of, 7.
Talc, an ingredient of the rocks, 4, 5, 6, 7, 9, 19, 20,-3,-5, 171,-2,
                        177,-8,-80, 213,-14,-21,-3,-7,-45,-7,-8,-55,
                        257,-98,-9, 300,-1,-2,-8,-4,-5,-7,-8,-11,-12,
                        814,-16,-21,-2,-4,-7,-72,-5,-7,-9,-87, 602,
                        647; Tables I. VIII.
                    of metalliferous deposits, 6, 23, 177,-80, 215,
                        221,-3,-7,-8,-9,-30,-3,-9,-40,-6,-63,-4,-5,-7,
                       268,-9, 302,-3,-4,-16,-25,-9,-80,-2,-4,-7,
                        876; Tables I. VIII.
                   of cross-veins, 270; Table VIII.
Taldanga, the sandstones, shales, and iron-ores of, 66.
Talho Aberto, (Open-work), extent of, at Catta Preta, 287.
                                     , at Gongo Soco, 256.
                                "
                     "
Tamar mines, proportions of silver in the lead-ores of the, 110.
      ----, erection of a steam-engine underground at the, 609.
Taquaril, flakes and lamine of gold in Jacotinga at, 278.
Tarag-ke-Tal, on the siliceous limestone of, 18, 31.
                     ores of copper at, 18.
                 ,,
```

Taxes on wood,	charcoal, a	mperature in the mining district of, 752. and lime in the Sub-Himalaya, 40. don), on early copper-mining in Cornwall,
		691,-2.
laylor, John, Es	q. (or Care	adon), on the silver-ores of Wheal Ludcott,
	- "	, ,, financial operations at ,, ,719.
	- "	, ,, copper precipitated from spring- water at Crow's-nest, 686.
Taylor, Capt. J.	on the p Balles	roportions of tin-ore in the vein-stones of widden, 472.
Tellurium, alloye		old, in ate of Brazil, 299, 834,-6,-7, 640.
 ,	"	,, , at Descoberta, 299.
,	"	", , at Coelho, 180.
,	75	", , at Catta Branca, 886.
 ,	"	Virginia, 836,-76,-82, 640.
		ld, in the Silurian rocks of Merioneth, 640.
Tennant, Prof.,	on a (nugg	get) mass of gold discovered in Australia, Table XXII.
Tennent, Sir J.	Emerson,	on the discovery of a ruby in a fowl's crop, 356.
Temperate region	ons, subte	rranean temperature in, 761,-2; Tables
-		XXXI. XXXV. XXXVI.
Temperature, su	bterranear	n, in Chili, 724-5,-51; Table XXXI.
 ,	"	, in Brazil, 725-32,-51.
 ,	"	, in the United States, 732-4,-51.
 ,	"	, in the Channel Islands, 784-7,-52.
 ,	,,	, in Ireland, 737-44,-52.
 ,	"	, in Cornwall, 744-9,-52.
 ,	"	, in Shropshire, 750-1,-2.
,	"	, in different rocks, 753-6,-60,-2; Table XXXIII.
 ,	,,	, in mines affording different metals and
		ores, 756-7,-8,-61,-2;
		Table XXXIV.
 ,	"	, ", similar metals and
		ores in different rocks, 757-
		758; Table XXXV.
 ,	,,	, " at different elevations, in tropi-
		cal and in temperate regions, 758,-61;
		Tables XXXI. XXXV. XXXVI.


Temperature, sul	terranean,	changes in, at given spots, by deepening
		and extension of neighbouring mines,
		764-5.
	,, 8	at inconsiderable in Brazil, 767—78; depths, Table XXXVII.
 ,	,, ,	", in New Granada 770.
 ,	,, ,	", in Ecuador, 771
 ,	" 1	,, in India, 772,-8, 80.
 ,	"	", in Belgium, 774-5.
 ,	"	", at Greenwich, 776-7.
,	"	" , near Truro, 779.
	"	" near Edinburgh, 777-
	,,, ,	,, at Upsal, 778. [778.
		d auriferous deposits of, 302-3.
		the peculiar alloy of the stolen gold, 336.
Thomas, Capt. C	haries, on t	he granite of Caradon, 663.
	, "	proportions of tin-ore in the vein-
		stones of Dolcoath and Cook's-
		kitchen, 472.
	 , "	silver & silver-ores of Dolcoath, 113.
	, ,,	progress of works at ,, , 448.
	, "	action of running-water in separating
		tin-ore from its matrix, 354.
	 , "	produce and profits of Dolcoath, 447-
		448; Table XIV.
Thomas, Henry,	Esq., on the	e separation of silver from Spanish lead-
		ores, 103.
	 -, ,,	produce and profits of the Consolidated
	_	Mines, 452; Table XIV.
		e sulphur-ores of Rio Tinto, 543,-6.
	, "	precipitation of , , , 589.
Thomas, Capt. J	osiah, on th	e rate of blasting rocks at Dolcoath, 200.
		the elvans of the Chasewater and Cam-
•	, 1,	borne districts, 662.
	 , ,,	heaves (dimlesoments) of lades in
	, ,,	Cornwall, 183.
	 , , ,,	normandianlanish of the michaet norte
	//	of lodes, 82.
	 , ,,	Gwennen adit 585
	, , ,,	publication of his work on the
	. "	Mining districts of Cornwall, 341.

Thost, C. H. G., Esq., on the mica-slate and serpentine of Corri
Charmaig, 645.
, ,, greenstones & porphyry of Breadal- bane, 651.
, ,, lodes of Breadalbane, 652.
Throw. (See Displacement.)
Thunderstorm at Catta Branca, effects of, 179.
Thury, M. Héricart de, on the rocks of Chalanches, 517,-19.
, ,, metalliferous \ 528.4
deposits)
, " ores they afford, 103, 524.
, " proportion of silver in lead-ore from
Vizille, 103.
Tigrony, on the rocks of, 542.
, ,, sulphur-ores of, 553-4.
Tilpora, on the iron-ores of, Table II.
Tijuco, on the slates and auriferous quartz of, 182.
Timber, quantities of, in the forests near Lake Superior, 390.
, of heavy growth, on the rubbish of ancient mines, 416.
Timbopeba, on the iron-furnaces of, 213.
Tincroft, on the granite and slate of, 658.
, proportion of tin-ore in the vein-stone of, 473.
, ventilation obtained at, by means of a fall of water, 220.
Ting Tang, on the profit realized at, 451; Table XIV.
Tin-mines, subterranean temperature of, 757,-8; Table XXXIV.
Tin-ore, an occasional constituent of rocks, 77, 175, 664-5.
proportion of, in the vein-stones of Cornwall, 472-3, 693.
, on the presumed association of it with fluor, 680.
,, separation of, from its matrix by running-water, 854.
, on proportions extracted by tributers, at Polberro,
- aulled from unbhish by day (909
labourers,
, detrital, mixed with gold, in Wicklow, 630-1.
, ", ", in Banca, 631.
in Anatolia 906
and plating in Ameteric 996
manufact to mine the angle and its 950
Danier and in most line mouth at half of 94K
Titanite, crystals of, imbedded in crystals of quartz in <i>Itabirite</i> , at
Agog Overte 225




Treatment of auriferous vein-stones, Tables VII. IX.
, different modes of, serve to extract different proportions
of gold from similar vein-stones, 198.
Trebilcock, Capt. J., on the auriferous deposits of Gongo Soco, 257.
, ,, of Antonio Pereira,
804.
Trebilcock, Mr. W., on the profit made at Carnon Stream, 452; Table XIV.
Trebisken Green, proportions of silver in the ores of, 121.
Tredinnick, Capt. S., on the proportions of tin-ore in the vein-stone of Great Work, 472.
· · · · · · · · · · · · · · · · · · ·
Treffry, J. T., Esq., on the produce and profits of the Fowey Con-
solidated mines, 456; Table XIV.
Tregoning, Capt. W., on the iron-formation of Gongo Soco, 260.
, , auriferous deposits of , , 255-72.
Trehane, Wheal, proportions of silver in the lead-ores of, 120.
Trelawny, Wheal, on the rocks of, 700-3.
, ,, lode of, 704-15.
————, ,, proportions of silver in the lead-ores of, 120,-708,-9.
, ,, cross-veins (flucans) of, 715-17.
, ,, rate at which the works have been deepened, 748.
marchine and manufacture and a completion and a completion of the complete of
at, 720.
, ,, produce and profits of, 719,-20.
, " subterranean temperature at, 748.
Trelawny, North Wheal, proportions of silver in the lead-ores of,
Trelawny, South, rocks and vein-stones of, 714. [120.
, machinery worked and population employed at,
, subterranean temperature at, 748. [720.
Treloar, Capt. T., on the rocks of Morro Velho,
, , , proportions of gold contained in
certain parts of, 186,-96.
,, , relations between them and the
principal auriferous deposits,
186,-7,-92,-8,-5,-6, 207,-8.
, , , circumstances which affect the
richness of the formations, 189, 190,-5,-6,-7.

Treloar, Capt. T., on the depo at <i>Morro Ve</i>	lho, endlo	ng dip (shoot) of ore,	207,-8.
, ,,	•	of rock which hem, 192,-3.	n sever p	ortions
	crev	of inflamm ices in the au 195,-6.	_	
, on the ex	tent of wa	ter-courses	at <i>Morro</i> Table 2	
-	_	gold obtained North d'el H		
Treloar, Capt. W., on the prestone of		f gold obtaine nara) Gongo		
Trelyon Consolidated Mines		n of tin-ore		
Tremayne (Providence) Wh Tresavean, precipitation of	<i>eal</i> , native	ailver in the		
———, produce and prof Treskerby, profits of, 451;	fits of, 451	; Table XIV		
Tres Puntas, silver exported Trethevy, cromlech of gran	from, 153	•	mation at	660
Trethevy mine, on the hornb	olendic rocl	cs of, 671.		, 000.
Trevandrum, temperatures various (incon	siderable)	depths, near,	} ****,-2,-	_
,	"	•	t different	780.
Trevillion, T., Esq., on the		ns of silver 's-foot, 709.	in lead-	ores at
 , "	ployed <i>Herod</i>	y worked, and; ; the product's-foot, 719,-	ce and pr 20.	ofits of
Treweatha, Wheal, proportion	ons of silve	er in the lead	-ores of,	120.
Treweek, James, Esq., on the		tion of coppe n mine-water		
······································	lim	e, at Wheal	, by the Falmouth	
Tributers (who receive a she on the proportions of			•	
Trollè Col., his use of fire-		• .		-



Valenciana (Chili), extent and produce of works at, 123.


Valenciana (Chili), quantity, quality, and value of silver-ore afforded by, Tables IV. XIV.
Valleys, directions of, in West Cornwall, 718.
Veins. (See Metalliferous deposits.)
Veins, cross. (See Cross-veins.)
Veins, pipe, of lead-ore, 620. (See Pipe-veins.)
Vein-stones. (See Composition of metalliferous deposits.)
Ventilation, natural, by the issue of air from joints in rocks, 250.
, artificial, by pneumatic means, 219-20.
Verneuil, M. E. de (Murchison, Sir R. I., & Keyserling, Count A. von), on the copper-ores of the Uralian coal-measures, 508-9.
, on deposits of gold and platina in the Ural, 340.
—————, on detritus of the Ural traced to its origin, 343. ————, on detrital gold of the Ural, 342.
, on the large (nugget) mass of detrital gold in
Siberia, Table XXII.
Verran, Capt. W., on the <i>shoots</i> of auriferous vein-stone at <i>Morro Velho</i> , 207.
Vicente, Saō, tellurium an alloy of gold in the neighbourhood of, 180.
Victoria (Australia), quantities of gold obtained in and exported from, Table XXII.
detrital gold of traced to its origin 249
numbers and weights of large (numets) magnet
discovered in, Table XXII.
Vincent, Wheal St., proportion of silver in the ore of, 115.
Virginia, on the auriferous rocks of, 372.
, ,, vein-stones of, 322,-75,-7,-81.
, ,, detrital gold of, 383,-4.
, ,, magnetic declination in, 372.
1' 1 f 1 - f - l ! 900 9 FVVVI
Vivian Carn, proportion of silver in the lead ore of, 120.
Vivian, Mr. John, profit in the Saint Ives Consolidated Mines, 443;
Table XIV.
Vivian, Capt. Joseph, proportion of silver in the ore of North Dolcoath, 113.
, on the comparative values of wheel-barrows
and tram-waggons underground, 144.



Walker, Dr. T. (& Reay, W., Esq.), on the processes and products
obtained in (dressing) cleaning gold at Morro Velho, 203.
Walsh, The Rev. Dr., on the gold-mine of Antonio Pereira, 303.
Walton mine, proportion of gold in the vein-stones of the, 373.
Waring, E. P., Esq., on magnetic declination at Chafarcillo, 69.
Water, spring-, diminution of, when the neighbouring surface is
stripped of wood, 137-40, 344.
temperature of, in Brazil, 728.
, , in the Channel Islands, 735,-7.
, , in Ireland, 738,-40.
, in East Cornwall, 745,-6,-8,-9.
(fresh), in mines wrought beneath the sea, 539.
, , , a surface encrusted
with the sulphate of soda, 165.
, , yielding a precipitate of copper, 686.
, prices of, in Chafarcillo, Table V.
, river-, unequal quantities of, at different seasons, 43, 140.
, action of, in separating metallic from earthy minerals,
354.
, now disappearing in gravel of ancient beds, 43-4,
189-40.
, but reappearing on reaching impermeable clay at a
lower level, 44.
, holding in solution salts but depositing them during
of sods and magnesia, its course, 189-41.
, distilled for use by loco-
motives, and sold for
domestic use, 141.
, mine-, unequal quantities of, at different seasons, 477,-573,
584,-5,-8,-90, 608-9.
, impregnated with salts of lime, 717.
, copper, precipitation from,
569-92; Table
XVIII.ª
, , , gradually diminish-
ing, 578.
, , ; lifts lined with
wood, and wooden plungers used for pumping, 569-70.
, holding iron-ore in suspension, preferred to clear water
by horses and mules 854


Water, mine-, fish thrive in, 855, 717-18,-31.
, entire absence of, in the deep works of Chanarcillo,
70,-148.
, prevents extraction and preparation
of much silver-ore at Chanarcillo, 138.
, fall of, applied to ventilating mines, 219-20.
the blasts of forges and iron-furnaces, 219,
Waterford, on the mean temperature at, 739. [528.
Waters, Sampson, Esq., on gold imbedded in grey copper-ore, 878.
Weapons, ancient, of native copper found near Lake Superior, 418-
Weaver, T., Esq., on the clay-slate of Waterford, 593. [19.
, " metalliferous rocks of Wicklow, 540,-1-2.
, " deposits of " , 543,—5,
549-50,-8,-4,-5,-6.
, ,, precipitation of copper at Cronebane, 562,
from mine-water) 573.
, ,, auriferous native silver of ,, , 105.
549.
, ,, detrital gold of Wicklow, 627,-8,-9,-30,-1.
Webb & Geach, Messrs. (See Geach & Webb, Messrs.) [633,-4.
Wellington, Mr. R., on the proportions of tin-ore in the vein-stones
of Condurrow and Ding Dong, 472.
, , (early) profits of Botallack, 442.
Wells, on the temperatures of water in, 735,-7,-8,-40,-5,-8,-9.
Wendron Consolidated Mines, proportion of tin-ore in the vein-stone
of, 472.
Werner, H. A. G., on the relations between metalliferous deposits
and the rocks which adjoin them, 187.
<u></u>
West, William, Esq., on the lodes of the Phanix Mines, Table XXV.
Westmoreland, proportions of silver in the lead-ores of, 109. [Note.
What-Cheer mine, on the metalliferous deposits of the, 465.
Whims, horse-, available only for the shallower parts of mines in
Chanarcillo, 142.
, erected underground, 76.
Whitehall (Virginia), on the deposits of gold and tellurium at, 376.
Whitley, Nicholas, Esq., his Geological map of the Caradon district, 655.
, on the structure of the Cheesewring granite, 672.
, ,, stanniferous granite of Gonamena,

Whitley, Nicholas, Esq., coincidence of the directions of the joints in rocks with the valleys in West Cornwall, 718.
temperature of ground of various kinds near Truro, 779.
Whitney, J. D., Esq., on the sandstones, conglomerates, and traprocks of Lake Superior, 392—404, 423,-38,-9,-75.
, ,, directions and dips of the copper- deposits of Lake Superior, 404,-7,-9.
vein-stones of the copper-deposits of Lake Superior, 424,-5,-61,-2,-3,-4,-7.
, ,, native copper of the copper-deposits of
Lake Superior, 435,-62,-3,-7,-76,-80.
masses at Lake Superior, 427.
, on the drift, and on the occurrence of masses of
native copper in it, at Lake Superior, 460.
, ,, native silver associated with native copper
at Lake Superior, 99, 419,-37.
, ,, prehistoric mines of Lake Superior, 412, and miners 414,—15, 417,-19.
, mining leases granted at , , 487.
, ,, argentiferous gold of California, 99.
Whitney, J. D., Esq., & Foster, J. W., Esq. (See Messrs. Foster & Whitney.)
Whittlesey, C., Esq., on the virgin silver and native copper of Lake Superior, 419.
, ,, prehistoric miners and mining of ,, 414
415,-19. and their ignorance of
smelting, 419.
forests, 416.
, ,, magnetic declination in Upper Michigan,
Whittington, extreme and mean annual temperatures at, 750. [395.
Wicklow, on the metalliferous rocks of, 540-2.
———, ", deposits of, 543-69.




WILLSOIL, W. LL., ESQ., (ou Noyer, G. V., E	sq.), stones of Kenmar 611,-14.
· ,	"	on the fossils the contain, 613.
 ,	"	on the metalliferou
,		nare, 614-15,-16,-18,-21.
Wilsworthy, on the nat		l cobalt, and copper-pyrite
	oportions of silve North of England,	r in the lead-ores of th 107.
F	ower, in, 142. 668	
-	stitution of, for <i>St</i> vall, 146, 645-6,-9	opes in the mines of Corn. 93-4.
Wire-rope, advantages	of, compared with	hempen rope, 143, 646.
Wood, Benj., Esq., on	the produce and	profits of Crennis, 456
		Table XIV.
Wooden pumps, genera	lly used in the gol	ld-mines of Brazil, 288.
Woods, their influence, laws for the pro-		eighbouring springs, 137- zil, 344. [140, 844
Woodville, on the talco	se-slate and aurife	erous deposits of, 375.
		-ores of Cornwall, 118,-19
Workmen in mines, ext		
		lver-ores of, 120, 710,-11.
Wright, Mr. J. E., qua		on of water in the Gwennap
Wyllie, J. F., Esq., on		nd prices of chrome-ore. Charmaig, 649.
		tin-ore in the vein-stones well, and Carnyorth, 472.
Yorkshire, on the displa		by veins in, 72, 227.
Zeppenfeld, M. (& Rivo		he proportions of silver in ach lead-ores, 104.
Zupia, mean temperatur	res at the surface	e, and at small depths in 768-70.

W. CORNISH, PRINTER, GREEN-MARKET, PENZANCE.












. . : •










RELATIONS BETWEEN THE METALLIPEROUS DEPOSIT OF AGUR AND THE ROCKS WHICH ADJOIN IT.

		Bed.		Denth	Composition and annearance of	Nature and enneavence of
Mine.	Direction.	Dip.	Sise.	fas.		adjoining rock.
Guarcoolee S.E. & N.W.	S.E. & N.W.	N.E.	ı	-	Specular (micaocous), brown, and magnetic Talo-clate with a little quarts. iron-ore, with small lumps of the oxide of manganeee. Structure lanellar,	Talo-elate with a little quarts.
Lhusganee	N. & S.	E. 22°-26° 6-8 feet.	6-8 feet.	8	Specular (micaceous) iron-ore, with a little Quartzose talo-slate, dark blue, bedding N. onartz and some talo.	Quartzose talo-slate, dark blue, bedding N. & S., dip E.
Nutos Kanh	E. & W.	zi.	1-9 "	8	Idem. In four distinct beds, interlying the Idem, sometimes blue sometimes drab; bed- slate, but ultimately re-uniting. ding E. & W., dip N.	idem, sometimes blue sometimes drab; bedding E. & W., dip N.
Gallà, S.E. & N.W.	S.E. & N.W.	N.E.		2	Idem.	Beds of homogeneous blue slate, alternating with quartsose talc-slate.
Dhoora Khani S.E. & N.W.	S.E. & N.W.	N.E.	2-7 "	17	Idem, with occasional masses of brown iron-Idem. ore.	ldem.
Capus	N. & S.	E. 48°	1-8 ,,	Surface	Surface Yellowish-brown fron-ore.	
			ł	9	Brown and micaceous specular fron-ore, with Homogeneous, fissile, pale blue, buff, and a little quartz.	Gomogeneous, fissile, pale blue, buff, and reddish-brown slate.
Choocooth 10° W. of N.	10° W. of N. & E. of S.	E. 40°	1	Surface	Surface Brown iron-ore, mixed with the carbonate Decomposed brownish talc-slate, of iron and a little quarts.	Decomposed brownish talc-slate.
Bunns	ı	1	ſ	:	Brown iron-ore and a little quarts.	Homogeneous, fissile, pale blue, passing into buff-coloured slate.
Purturburà	E. & W.	N. 24°	10–12 "	6.5	Some portions of the fron-ore pale others Pale brown and buff talc-slate adjoining the dark brown; masses of slate, in some places slightly impregnated in others vein-little distance.	ale brown and buff talc-slate adjoining the ore; homogeneous dark blue slate at a little distance.



RELATIONS BETWEEN THE METALLIPEROUS DEPOSIT OF KOTELAR AND KHETSAREE AND THE ROCKS WHICH ADJOIN IT.

		Bed.		Depth	Composition and appearance of	Nature and appearance of
Mine.	Direction.	Direction. Dip. Size.		fms.		adjoining rock.
Tilpora*	N. & S. B. 20°-25° 2-5 feet.	E. 20°-25°	2-5 feet.		Compact red iron-ore, with occasional stones(of quarts; drusy cavities lined with earthy	Compact red iron-ore, with occasional stones Clay-slate; reddish-brown, with flakes of of quarts; drusy cavities lined with earthy mica between the lamines; joints curved,
Chitalee	% %	pi 	ı	ı	yellow iron-ore. Clay-slate and slaty clay, quarts, and carbo-(nate of lime, with large irregular masses of scaly red iron-ore.	yellow iron-ore. Clay-slate and slaty clay, quarts, and carbo-Clay-slate; reddish-brown, mottled with nate of lime, with large irregular masses white in some places; in others bluish-of scaly red iron-ore.





DESCRIPTIO IN THE COLORADA LODE.

		Third Limestone.	
Mines,	References to (Section) Pl. II.	Nature of ore.	Approximate quantity of Silver. lbs. Troy.
Manto de Ossa.	_		
Yalenciana.	A B C		
Esperanza.	D E F		
Colorada.	G		
	H K		
	L		
	P		
	Q+		
1	-	pyrites, blende, native silver, and sul- turet of silver.	_
Desempeño.	M	and of shife,	
	Q*		
San Francis- quito.	R† R†		
gano.	8		
Bocona.	T		
San José.	N		
	υ		
San Francisco viejo.	0		
	V		
San Francisco nuevo. Delirio.		enical silver ore, sulphuret of silver, red lver ore, and native silver. enical silver ore, red silver ore, and tive silver.	_




			_		_				М	ru	ATIO	N.							
	Ti	mbe	r.		Coal		Iron é	k St	eel.	ic.	Prov	ende	pr.	Mate	rials	,	Tot	als,	
d. 5	£	s. —	d.	£	8.	d.	£.	8.	d.	d. 1	£ 11	8. 11	d. 5	£ 244	8. 15	d. 1	£ 1047	8. 19	<i>d</i> . 0
0		_			_			_		. 4	4	12	0	200	2	1	944	7	5
1		_			_		31	8	6	: 8	22	1 ,8	1	176	0	8	1103	9	7
2					_		67	16	0	11	119	15	2	382	14	0	1491	4	10
8		_			-		26	0	0	1	127	10	8	319	2	8	1414	9	4
6		_			_	ļ	-	_		7	44	17	0	285	11	8	1255	8	10
8		_		12	12	3	20	3	9	. 1	19	15	8	233	1	9	1160	17	6
3		_		27	4	10	21	19	1	7	17	14	3	161	14	5	996	19	9
7		_		3	8	10	17	16	0	٠ :	11	<u>``</u> .6	7	172	19	1	973	9	1
0		_		2	0	0	19	9	7) :	8	0	0	181	16	11	1145	11	3
		_		34	5	2	-	_		;	3			204	0	11	1031	11	10
	18	10	11		_			_		, .	4	_		828	18	9	1218	2	6
4	18	10	11	79	6	1	204	12	11		387	15	10	2890	18	0	13783	10	11
	_		_						N	fiend	liture	,							
									1	SI 1(11.								
es q	f I	ife).																
o £	:3	1	10)							•		_						
	1	4	4	ļ	per	c₩	t.			į t	. 2 0		2 10)					
ŀ	1	0	5	(ļ t		15	4						
<u>'</u>	0	17 0	6 Տ <u>ե</u>	,	per	lb.					. 5	10	5	Per	CW	t.			
to	0	4	0 0		eac						. 5	10	5	1					
	0	4	0		per	dos	en.				. 5	6	0	,					
	0	1	0			pin					. 0	8	0		ga	llon	1		
	0	0			,,	gal	lon.				. 0	8	0	ead		••			
l	0	Ω	뱒		,,	,				1. 1	ю 0	0	8 1	pe	r co	u,			

. :



96,612 tons.
71,202 ...
381 to adid
40 to 2 to 2.
381 to adid
40 to 2 to 3.
40 to 3.
40 to 3.

Mindle. ١




rom aid in Brazil.					
r hid baiz Export. rasha ama cationid in Paid in	Proceeds of gold dust sold.	Cost.	Loss,	Profit.	

.

.

1







:

ial Brasilian Mining Ast alry Dividend at first set s ¹¹al Brasilian Mining Ass

• -






Name & direction of vein.	. Dip.	Size. feet.	Composition and Appearance of Lode.	Composition and Structure of Rock.
15.	i	0.6—1.3	E S	nde, labradorite, and chlorite.
38.	,, 66°-78° 0·1—2·6	0.1-2.6	rocks, thinly spotted with grains of native-copper. Idem, with grains and small masses of native-copper.	dips N.W. 40'-50'.
8	., 66°-82°	0.6—2.	1.66-820 0.6-2. Calcareous spar, quarts, and chlorite,—with traces of Hornblende, labradorite, and chlorite; of amorphisms and chlorite; of amorphisms and chlorite; of an amorphisms and calculations.	ornblende, labradorite, and chlorite; of
			thick, slightly I	the cavities filled with calcareous
46.		0.1-6.	", 60°-80° 0.1-6. Calcareous-spar, chlorite, quartz, and epidote; enclosing small masses, and sprinkled with granules of native-copper.	ė
			At this place the North-Bartur vein unites with the lode.	
9 6.	" 68°-74°	03-2.5	", 68b-74° 0 3-2.5 Calcareous-spar, chlorite, quarts, and epidote; thinly Hornblende, labradorite, and chlorite, sprinkled with grains of native-copper, and enclosing masses of hornblendic and felspathic rocks.	inde, labradorite, and chlorite.
1	 	1	Neither lode nor branch has been traced in the crystal-Hornblende, labradorite, and chlorite; fine greenstone.	ornblende, labradorite, and chlorite; fine grained and crystalline. (Ante, p. 398.)
26.	N.E. 72	0.8-1.	Calcareous spar, quarts, chlorite, and prehnite; thinly sprinkled with native-copper.	
1 6.	N.E. 75°— 85°	0.2	Calcareous-spar, quartz, prehnite, and chlorite; thinly sprinkled with native-copper.	

 Jackson, Report on the Geological and Mineralogical Survey of Lands in Michigan, 111. pp. 458. Foster & Hill, Ibid, pp. 760—61. Foster & Whitney, Report on the Geology of the Lake Superior Land District, 1. pp. 132,—46—7, Pl. IX. Fig. 20, Whitney, Metallic Wealth of the United States, p. 279. Rivot, Annales des Mines, 6me Série, v11. p. 315.

Mining Magazine (New York, May, 1854), 11. p. 557.



THE CLIFF MINE .- DISTRICT OF KEWEENAW POINT.


Name & direc. Denth	Donth	l	ä		
tion of vein. fms.	fms	ġ A	feet.	Composition and Appearance of Lode.	Composition and Structure of Rock.
Lode 21° W. of N.	Surface.	B. 70°-76°	0.1-0-2	Lode 21° W. of N. Surface. B. 70°-76° 0·1-0-2 Prehnite, calcareous-spar, and quarts, enclosing Hornblende and labradorite (Greenstone); fine-	Iornblende and labradorite (Greenstone); fine-
				capillary red oxide of conner, and thinly stud. defined, and of crystalline structure, in ill-	grained, and of crystalline structure, in ill defined hade which die N N W 959, 200
	-1 20c I	2. 70 -86 I	<u>0-30 II</u>	ne ingredients—as well earthy as metalife—	Idem.
			F	The lergest masses of netice conner concerni-	
			<u>. </u>	occur on or near the lower side (foot wall) of	
		_		the lode.	
			<u></u>	In greenstone the lode is always small and poor.	
_	_				

• Inckson, Report on the Geological and Mineralogical Survey of Lands in Michigan, 111. pp. 469—60. Foster & Whitney, Report on the Goology of the Lake Superior Land District, 1. pp. 127-31,-72. Whitney, Metallic Wealth of the United States, pp. 276-9. Rivot, Annales des Mines, ome Série, vii. pp. 510-14. Lake Superior Miner (26th April, 1867), II. p. 2.

† Ante, p. 439.

Daniel, Mining Journal (2nd September, 1865); xxxv. p. 567.

.



• . · • • . • ł

THE DOUGLASS HOUGHTON OR HENWOOD MINES. - DISTRICT OF ONTONAGON. Fig. 28, 29.

smaller quantities smaller quantities orite; in thick beds oficection and dip. nall irregular veins, te, —which contain occur at intervals.	
Composition and Structure of Rock. Hornblende mixed with smaller quantities of labradorite and chlorite; in thick beds parallel to the lode in direction and dip. Isolated masses, and small irregular veins, of quarts and epidote,—which contain particles of copper,—occur at intervala. Idem. Idem.	" Mechigan, III. pp. 400, 702.
Name & direc. Depth. feet. Composition and appearance of Lode. Composition and Structure of Rock. Lods 40-50° E. of N.W. 60° A mere Renearly sprinkled with grains, and in several places of labradorite and chlorite; in thick beds 2.6 joint to with small meases, of native-copper. CONGLONE. Surface N.W. 40° 1.5 words the south-toest. CONGLONE. Surface N.W. 40° 1.6 words the south-toest. CONGLONE. Surface N.W. 40° 1.6 words the south-toest. CONGLONE. Surface N.W. 40° 1.6 words the south-toest. CONGLONE. Of N.W. 40° 1.6 words the south-toest. CONGLONE. Surface N.W. 40° 1.6 words the south-toest. N.—W. of S. 71. "38° 42° 0.6 — 0.8 A (leader) central vein of calcareous-spar, quarts. CONGLONE. Surface N.W. 80° 1.6 words the words	* sackson, deport on the veological and mineralogical Survey of Lands in michgan, III. pp. 406, 702.
Size. feet. Joint to 2:6 0.3—0.8 0.5—0.8 0.5—0.8	a, Meport
Burface N.W. 50° A me to 13. Surface N.W. 40° Joint 2.6 to 9. 71. "38° -42° 0.5—0 48. N.E. 80° — 0.5	PACKED
Burface to 13.	1
Name & direction fms. LODE 40°—60° E. of to 13. N.—W. of S. CONGLOME 60 9. A0°—60° E. of to 13. CONGLOME 60 9. N.—W. of S. CROSS-VRIN CROSS-VRIN S.E. & N.W.	

destitute of col

Foster & Whitney, Report on the Geology of the Lake Superior Land District, 1. pp. 142—60. Whitney, Metallic Wealth of the United States, pp. 289—90.

Lake Superior Miner, x1. (24th March, 1856), p. 224. Henwood, Report on the Douglas Houghton Mine (Detroit, 1869), pp. 1—9. Ingram, Coulter, Douglass, and Rudolph, Eshibit of the Honsoood Mine (New York, 1864), pp. 10-18.

‡ Ibid, pp. 172,-81,-2; Tables XVIII., XCV. + Henwood, Cornwall Geol. Trans., v. pp. 25, 328; Table LXXXIV.



THE MINES OF CHALANCHES D'ALLEMONT, IN FRANCE.

Name & direc-		Rine		
-8 1	40.—48		granular quarts.	
STR. Hanker. 20° W. of N.— E. of S.	W.88°-42°	1	STR.HELKITH. W.38°-42° 1. Quartz, sabestus, and earthy brown iron-ore. Particles of native-silver at intervals. R. of S.	
Brisks. 86° N. of E.—S. of W.	8. 50°-70°	3.0—4.0	BRISÉR. S. 50°-70° 3·0—4·0 Gneiss, exactly resembling the adjoining Gneiss. Basis of felsper, quarts, and horn- blende, enclosing erystals of hornblende of Glaper, with small masses of cal- careous-spar.	safts, and horn- s of hornblende masses of cal-

The works near this deflection scarcely suffice to prove whether it is, or is not, due to the influence of some other vein.




SARKS-HOPE MINE.—SARK.

•

Copper Lode.			
Composition and structure of Rock.	Felspar and hornblende, mixed with calcareous-spar, occur in ill-defined beds, of massive structure, which dip towards	Bard votince with conservous spaces Bracking generally massive. Pale yellow felipst, chlorite, and born blende, slighty mixed with cal- carous-par. Many joints are faced with steatitic matter. Idem.	Felspar & hornblende, traversed by thin veins of calcarcous spar; in ill-defined beds, of massive structure, which dip towards the N.E. Idem.
Composition and appearance of Lode.	8. Felspar, quarts, and hornblende, veined with calcareous. Felspar and hornblende, mixed spar and spotted with rich argentiferous galena. 2.0—2.6 N.E. Felspar-clay and granular quarts, streaked with ill-defined beds, of massive early brown iron-ore, and spotted with calcareous-spar, structure, which dip towards	iron-ore, and still more thinly with earthy black silver-ore. The same earthy ingredients, enclosing small rich Pale yellow felapar, chlorite, and benches and narrow veins of earthy black silver-ore. S.W. Pale-yellow felapar, chlorite, and calcareous-spar; silghtly mixed with argentiferous galena, earthy black silver-ore and minute crystals of red silver.	4.0—6.0 N.W. or upper side, yellowish felspathic clay, enclosing Relignar & hornblende, traversed masses of calcareous-spar spotted with iron-pyrites. S.E. or lower side, earthy reddish-brown iron-ore and granular grants, enclosing nests filled with felspar-clay, earthy manganese, or calcareous-spar. N.W. Barthy reddish-brown iron-ore, containing isolated masses of felspar-clay, of calcareous-spar, and of iron-pyrites. N.W. vein, felspar-clay, of calcareous-spar, and spotted with iron-pyrites. S.E, felspar-clay, sprinkled with granular quarts, reined with calcareous-spar, and spotted with iron-pyrites.
Sise. feet.	3· 2·0—2·5	6.0	
Dip.	n.e. 85° —	N.W. 70°—76° N.W. 72°	N.W.70° N.W.72° N.W.76°
Depth.	Water's odge. 14.		44 44
Name & direc- Depth. tion of vein.	SILVER LODE. Water's. N.E. 85° 25° E. of N.— edge. W. of S. 14°		Copper Lode, N.E.—S.W.

· The mine is about ninety fathoms deep, and some of its (galleries) levels are extended nearly sixty fathoms beneath the sea.



3175	1.0239	0.0184	0.0308	0-0993	0-0969
2481	0.7284	_	0.0265	0.0776	0.1066
2233	0.6505	_	0.0230	0.0670	0·1029
2767	0.8979	-	0.0287	0.0902	0.1004
3887	0-9995	-	0.0284	0.0731	0•0731
_	_	_	-	_	0-0798
2991	0.6528	0.0441	0.0352	0.0769	0.1178
1279	0-8478	0.0283	0.0277	0.0549	0.0647
1118	1.2288	0.0211	0.0239	0.0718	0.0582

omprises broken machinery and large junks of any kind." ndent of the *Mona* Mine and Smelting Works.






Table XX.


THE MOUNTAIN LODES OF THE (ALLIHIES) BEARHAVEN MINES-COUNTY OF CORK.


Name & direc- Depth. tion of vein. fms.	Depth.	Dip.	Sise. feet.	Composition and appearance of vein.	Composition and structure of Rook.	Main Lode. course.	Cross-
MAIN LODE. 26° N. of E.—	43.	N. 60°-70°	42.	Quarts, enclosing, at intervals, grains, small amasses, & large bodies, of copportpyrites.	Quarts, enclosing, at intervals, grains, small Slate; commonly pale-buff or lilac-coloured, masses, & large bodies, of copper-pyrites.	·	
	140		,	thinly sprinkled with copper-pyrites. As the Mountain lade improves, the Main	(9)	the	
	203.	68°—76°	4 660	Mode accines, in quality. Quarts largely mixed with copper-pyrites.	Buff-coloured, thick lamellar slate; of which	ptps use ptps in silice	
	226.	70°—78°	50.—62. 48.—60.	Idem. Quarts largely charged with copper-pyrites	70°-78° 5062. Idem.	m tne si feet by I resem in positio direction	
	240.	64°72°	4004	lor about 20 fathoms in length. Quartz and quartzose slate, mixed with		orn san 18 ts 18 em 18 w su 18 w su 10 on in 10 mi me tinnounti	
				copper-pyines, and win the carbonate of iron and calcareous-spar in smaller pro- portions.	7,00	o; but ted so ch — s rocks lth th	
				The mass of copper-ore dips or shoots to- wards the N.		At less the following the solution of the solu	
CROSS-COURSE N.E.—S.W.	-88	S.E. 80°	8-4.	The Cross-course consists mostly of quarts, Buff or lilac-coloured slate, where it intersects the Main lode; but	Buff or lilac-coloured slate.	3	
	115.	76°84°	4,6.	mostly of slaty-clay elsewhere.* Idem.	Idem.	· S	


* Captain Henry Pascoe, A.B.G.S C., Manager of the Mines, MSS.











• • -. d







Table XXV.

THE PHENIX MINES, PARISH OF LINKINHORNE.

ı •

Name and Direction of Vein. Lode. Lode Acaved feet L.S.A. B. of S. E. 80le-buff hue; of thick lamellar struction on slight exposure at the surface. Pis on slight exposure at the surface. 5°; but in some places it is nearly the lode in direction and dip, are at and occasionally with crystals of the B. 8structure, traversed by many joints feet L.S.A. E. 80le-planes of cleavage dip S. 8°-10°. Piode.	10
N.—S., 5° W. of N.— B. of S. P is on slight exposure at the surface. 5° but in some places it is nearly the lode in direction and dip, are at and occasionally with crystals of the E. getructure, traversed by many joints E. 80te planes of cleavage dip S. 8°—10°.	10
feet L.S.A. 55 R. 80th planes of cleavage dip S. 8°-10°.	
" E. 76 84'	
F. 68s of cleavage dip S. 8°—10°. Tra- 76r-clay. (c) 1t with one side of the lode,—consists 1.—(1) a central body of galena; entranslucent quarts,—(3) milk-white 1.—(5) yellowish quarts containing mi- 1. of quarts,—(6) galena slightly mixed 1. te, which gradually passes into the 1. of the product of the lode,—consists 1. s. d.	feet
68 R. 7d	
75 R. 78 E. 8 Glende in separate crystals: generally W. etimes of schistose structure. s, dark blue.	
95 E. Sthe planes of cleavage dip S. 10°—	
105 R. 74	
FLUCAN.‡ 30 S.W. 60 E.—N. of W. 40 S.W. 55 S.W. 55	

[•] Continuation of the smaller flucans, which do not reach the surface.
† The ore occurs in smaller flucans, which do not reach the surface.
† The ore occurs in smaller flucans, which do not reach the surface.



		1	•	İ		
	:					
nd hornblendic matter, of crystalline character, but dis-	nd hornblendio rocks, fine-grained, massive, and traversed sints; passing gradually into homogeneous slate.	(a)	(6)		Рама,, VII. (1862), p. 204.	
H	. H A		_ -	•		
ı						
				•		

Ĭ

. . .

ight breese, s. ŀ.B. risk breeze, S.E. E. ight breeze, W. , W. , s. w. risk breeze, E. ight breeze, S.W. risk breeze, E. , N. B.W. , N. ₿. lale, N. W. risk breeze, R. E. ery light breeze, E. ight breeze, E. ₩. , s.w. 1 29 N.W. risk breeze, S.W.

à

349. luy. d. London, E

.

Table XXXI.


TEMPERATURE AT AGOA QUENTE, -- BRAZIL.

(Long. 43° 10' W. Lat. 18° 50' S. about 3,400 feet above the sea). Deduced from 2,398 observations in 1848 and 1849.

.adta		3 A. K. b	ا ۽		6 A.K.			k k k		နိ]	Noon.		4 P.K.	j (6 P.K.	ا پر		8 P.M.a		°	9 P.K.	- (MIDNIGHT.	OHT.		MONTHER EXTREMES & MEANS.	75 26 26 26 26 26 26 26 26 26 26 26 26 26
o)K	Max.	Max. Min. Av. Max. Min.	Av.	Max.		A 7.	Max.	Max. Min. Av.		Max. Min.	<u></u> [Av. Max.	kr. Min.	n. Av.	_	Max. Min.	Av.	Max.	Max. Min. Av.		Max.	Max. Min. Av.		Max. Min.	λ. Αν.		Max. Min.	44
1848.																										•		
Oct 66.	.99	63.6	60.6	63.5 60.6 67.3 56.6	26.56	.663.2	74.5	74.5 58.5 67.8		7.5	3.	·7 7°	-9-	77.5 68. 72.7 79.5 68.5 73.7 75.	75.	÷;	ئ _ە .	73.8	73.8 62. 68.2 73.	8.3		61. 67·6	99	69- 61	°59	79.	5 63.	81. 68.2 79.5 53.5 87.5
Nov	-69	62.	63.1	63.1 70.2 59-		2.99	74.5	63.6 69.		-62	68.2 73.5	.6 82.		67.8 74.4 77.5 66.6 71.3 75.	177.	9.99	71.3	.92	64.3 69.1 74.	39.1		63.2 68.4	<u>*</u>	71. 64.	64.8	82.	62.	8.89
Dec 71. 60. 66·1 72·3 63·	71.	ė	66.1	72.3	œ	9.69	.82	69-2,73-8		83.	3. 7.7	73 77.7 84.		.2 77.8	3 79.	5 71.2	76.2	76.3	73.5 77.8 79.5 71.2 76.2 76.3 69.2 72.9 74.2 63.2 72.2 73.	6.7	74.2	33-2 75	7.5	99	68.2	84.	8	72.6
1849.																												
Jan	72.	61.	61. 66.8 73.	73.	67.4	70-1	2.82	78.5 63.374.2	4.2	4.6 71	9. 78	.1 84	8 72	84.5 70 78.1 84.8 72 79.1 81.	1 81		75.9	8.92	70.8 75.9 76.8 70. 73.4 76. 69. 72.6 73.	13.4	.92	-69	9:6	÷ 64·		68.5 84.8 61.	3 61.	73.2
Feb	5	-09	86.2	66.2 72.8 62.	8.39	-898	78.2	64.6 72.1		1.5 6	81.5 63.8 75.6	9.	.5 64	80.5 64.5 74.7 78.	78.		63.5 72.6 75	.92	63.6 70.9 74.8 63.5 70.3	6-0	74.8	33.6/70	-3 72.	69	. 67.		81.6 60.	70.7
Mar 68.	-88		86.4	57. 65.4 70.2 61.		67.3	76.2	75.2 66.2 71.8	_	0.1	2.2	.3	19 91	. 126.	3 78.	2 64.	72.	74.	80.1 67.6 75.3 80.5 67. 176.2 78.2 64. 72. 74. 64.8 70.6 73.	9-0	73.	65.2 69.8 70.	8.	÷		66.5 80.5 57.	2 67.	70.4
Apr 69. 54 62.6 70 57:5	-69	ż	82.6	2		6-7	.92	64.9 76 62.5 69-	_	6	7. 73	*	99	79. 67. 73.4 80. 65.5 73.5 75. 62.	75.	62.	70.4	72.2	70.4 72.2 60. 68.5 72. 59. 67.7 70.	38.5	<u>~</u> 23:	-69	1.4		67. 64.3	380	80. 64.	68.2
May	64. 42.	43.	52.8 66.2	86.2	\$	6.99	9.99	66.6 54. 61.4	1.4	1.5 6.	2. 67	.1 72	19 24	71.5 62. 67.1 72.7 61.5 67.4 69.	·69	-89		63.9 67. 66.		62.3 66.		55 61.5 64.	.6	. 20		56.6 72.7	7 42	61.
June 60.		43.	53.4 65.	.99	47.66	6.999	.99	52.2 60.5		-02 20	62.3 66.3	.3	1.4 64	73.4 64. 67.4 70.	<u>\$</u>	28		63.3 69.2 55.		61.9	-69	63.6 61.3	.3 66	÷ 48	. 56.8		73.4 43.	60.7
July	54.	46.	47.6 56.		48.20	.299.	9.09	60.5 54.2 56.8		6.4	3.2 64	66.4 63.2 64.7 68.		63.8 66.5 65.	65.	9		61. 63.	€7.	63.	62.2	62.2 56.4 59.4	4-	56. 48.	. 61.8	89 88	46.	9.19
		j	Ī		1	7	Ť	ᅡ	╬	 	+	+	- -	\downarrow	4	1			1	7	1	+	╁	+	4	4	_	_[
Extremes . 72.	72.	42.	:	73.	48.2	:	9.82	52.2	œ :	84.5 62.		84.8 61.5	1.8 61	٠	81.	.89	:	.99 8.92		:	.92	9.89	:	73. 48.	:	84.8	8 42	:
Means	$\overline{:}$: 80.3	80.3	:	:	63.6	:		9.29	<u>:</u>	. 72.4	:	<u>:</u>	Ş	:	<u>:</u>	69.6	:	:	67.8		67.		: :	62.9	:	:	67.

a. The observations at 8 p.m. are not used in deducing the Means.

b. At Midnight, and at 3 a.m., Captains Pengilly, Luke, and Guy were the observers.



ermal springs at Agos Quente (c) have not been reckoned in these averages.

			Ratios.		₹31.4		5 6	È	:	16.3
		MEANS.	Temperatures.	61.	9.09	86.4	73.	73.2	:	82.8
		¥ .	Meen depths. .emt	8	99	122	166	227	:	62
			No. of observations,	41	17	14	63	69	78	:
ſ			Retios.	:	:	:	:	:	:	:
İ		ENGLAND (Salop d).	Temperatures.	o.∳	:	:	:	:	:	54.
		Erro.	Meen depthe.	16	:	:	:	:	:	16
١	İ		No. of observations.	1	:	:	:	:	1	:
ار		•	Ratios		9 9	9	:	:	:	8.9
	j	ENGLAND (Cornwall).	Temperatures.	51%	62.2	8.99	:	:	:	62.5
946		Corny	Mean depths. fms.	33	78	116	:	:	:	8
. B		<u> </u>	No. of observations.	2	4	1	:	:	13	:
Less than 2000 ms. above the sea.	ions.	ė,	Ratios.	_	14:1	_	:	:	:	14-1
200	temperate regions.	IRRLAND, a b	Temperatures.	61.6	60 61.3	2-69	:	:	:	68.4
3	E S	IRRI	Mean depths. fms.	8	9	127	:	:	:	29
	3		observations.	9	8	8	:	:	12	:

T COUNTRIES.

• .

	Retios.		721.4		, é	200	:	16.3
NB.	Temperatures.	61.	60.5	85.4	72.	73.4	:	62-3
Means.	Mean depths. fms.	28	99	122	165	227	:	62
i	No. of observations.	17	11	14	61	61	76	:
•	Ratios.		8.9				:	8.9
Granite, a b	Temporatures.	61.4	:	87.8	:	:	:	69.5
FRAN	Meen depthe.	82	:	126	:	:	:	79
	No. of observations,	67	-:	61	:	:	4	:
ю тв, О	Retios.						:	:
sr, Micacrous, Orloritic Slates, b	Temperatures.	62,8	:	:	:	:	:	62.9
LOOSE, MICA & ORLORI SLATES.	Mean deptha, fms.	18	_ <u>;</u>	:	:	:	:	18
3	observations.	64	•		-	_	1 -	

mines (b),-and of the thermal springs at Agoa Quente (c).




Table XXXIV.

Subterranean Temperature in Mines yielding different Metals and Ores.

	Retios.	_	} 21:4	_	Ġ	.09	:	16.3
MBANS.	Temperatures.	61.	9.09	4.99	72.	73.2	:	62.3
M	Mean depths. fms.	78	99	122	166	227	:	62
	No. of observations.	4	17	14	8	61	78	:
. a .	Ratios.	:	:	:	:	:	:	:
t Tra	Temperatures.	:	:	65.7	:	:	:	2.99
Copper & Tin.	Mean depths. fms.	:	:	114	:	:	:	114
Cor	No. of observations.	:	:	9	:	:	9	:
9	Ratios.	:	32.7	12.6	:	:	:	16.9
COPPER. ab	Temperatures.	52.6	53.7	2.69	:	:	:	53.7
Cor	Meen depths. .emt	22	99	127	:	:	<u>:</u>	
	No. of observations.	16	9	8	:	:	ឌ	:
	Ratios	:	8.7	9.8	:	:	:	8.4
D. a.b	Temperatures.	8.99	60-2	2.99	:	:	:	2.09
LEAD. a,b	Mean depths. fms.	34	20	121	:	:	:	72
	No. of observations.	7	2	87	:	:	11	:
	Retios.	:	38.3	:	14.8	:	:	21.6
SILVER.	Temperatures.	°4.8	:	67.2	:	73.2	:	69-2
SIL	Mean depths.	48	:	138	:	227	<u>:</u>	166
	No. of observations.	н	:	81	:	8	9	:
	Ratios.		Ė	21.8	13.3	:	:	23.
Gогр. во	Temperatures.	9.99	87.6	71.4	72	:	:	9.29
Gor	Mosn depths.	31	49	147	166	:	:	21
	No. of observations.	23	9	67	63	:	25	:
	Depths.	Surface to 50 fms	60 ,, 100 ,,	"160 " " " " " " " " " " " " " " " " " " "	160 ,, 200 ,,	200 ,, and beyond	ala	Means
		Surfa	_	=	Ä	ā	Totals	Kea

In deducing these results, the temperatures—of wells at the surface (a),—of the water pumped out of mines (b),—and of the thermal springs at

Agoa Quento (c) have not been taken into account.

.

IN DIFFERENT ROCES.


han 200 fms, above the sea. egions of the Northern Hemisphere. Corresp and Corresp-ora in Sandstone. biendic Rocks. d

Table XXXVI.

Subterranean Temperature in Mines situate at different elevations above the Sea.

	Ratios.		21.4	_	.9	.09	16-3
	Temperatures.	28 61.	9.09 99	85.4	72.	73.2	62.3
	Mean depths. fms.	28	99	122	165	227	62
	No. of observations.	41	17	14	67	ea	92 :
Мядив.			:	14 122 65.4	:	200 and beyond	
	Depths.	60 fms	60 ,, 100 ,,	100 ,, 150 ,,	200 ,,	d beyon	
	.	Surface to 60 fms.	" 09	100 ,,	150 ,, 200	200 sn	Totals
	Ratios.	٩	9	0.0	:	:	: 6.
ea.	Temperatures.	23 63°	66 57.5	63.8	:	:	61 67
the Sabo	Mean depths. fms.	23	99	119	:	:	61
bove rions.	No of observations.	17	12	91	:	:	88 :
Less than 200 fms. above the Sea. (In temperate regions.) abo	,			:		200 and beyond	Totals
(In tem	Depths.	Surface to 50 fms	60 ,, 100 ,,	100 , 150 ,,	150 ,, 200 ,,	and beyor	
ĭ		Surface	90	100	150	300	Totals Means
	Ratios.	26.6	00 00 10 F	40.0	4.0 .0	3	: &
šea.	Temperatures.	68.7	64 67-6	69-3	72.	227 78-2	67.7 30.
the 20	Meen depths.	32	64	4 143 69-3	166 72		: 99
sbove ions.)	No. of observations.	24	2		61	61	37
More than 200 fms, above the Sea. (In tropical regions.) 22	Depths.	Surface to 50 fms	60 ,, 100 ,,	160 ,, 161	200 " "	200 and beyond	Totals
Mor	1	Surface to	. 09	100 "	160 ,, 200	200 an	Totals

In deducing these averages, the temperatures—of wells at the surface (a),—of the water pumped out of mines (b),—and of the thermal springs at Agoa Quente (c), have not been taken into account.



NTE IN BRAZIL.


							_			
Days.	JULY. UNDERGROUND. lours of observation.						SURPACE. Hours of observation.			each . a
	Noon. Depth,			6 P.M. Depth.						Means of ea
	3 feet 6 feet 9 feet		3 feet 6 feet 9 feet			8 A.K.	Noon.	6 P.K.		
1 .	71°.7		71 [°]	71°.7	••	71 [°]	51°.	63 [.] 9	61 [°]	56°4
2 -	71.8	••	71.	71.7	••	70-9	61 ·	65.4	62·	57·2
3 -	71.8	••	70.8	71.6	••	70.8		63.2		56-9
4	71.4		70.8	71.4	••	71.		63.9	1	56·
5 -	71.5	1	70.8	71.5	••	70-8		65.4		57.7
6	71.5		70.8	71.4	••	70.8	50.2		65.	<i>5</i> 8·
7	71·6 71·2	١.,	70·8		••	70·8 70·8		64·2 65·4	60·	<i>5</i> 9·1
8	71.2	••	70.8	71·2 71·1	••	70·8		65.	1	58· 59·5
9	71.2	1	70.8	71.3	::	70.8			61.0	58.3
10 ₄	71.3	1	70.8	71.3	1	70.8			62.	58-1
12	71.3	,	70.8	71.3	l l	70.8				57·1
13							52.2		·	Ī
1										"
14 (::		••		••	"	''		١
15		••	•••	••	l ''	••		١		١

. : . • • •

.

.

.



DEC 18 0 1888





